

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



PCT/GB 0 1 / 0 2 5 7 2

112



2001

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 04 JUL 2001

WIPO


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation and Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the international application filed on 13 June 2000 under the Patent Cooperation Treaty at the GB Receiving Office. The application was allocated the number PCT/GB00/02302.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or the inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



COCIU

Signed

T A Roberts

Date:

15 June 2001

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

HOME
copy

PCT

REQUEST

The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty.

For receiving Office use only

PCT/GB 00 / 02302

International Application No.

13 JUN 2000

13.06.2000

International Filing Date

United Kingdom Patent Office
PCT International Application

Name of receiving Office and "PCT International Application"

Applicant's or agent's file reference
(if desired) (12 characters maximum)

00111WO

Box No. I TITLE OF INVENTION

COMPOUNDS

Box No. II APPLICANT

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is country) of residence if no State of residence is indicated below.)

ELI LILLY AND COMPANY
LILLY CORPORATE CENTER
INDIANAPOLIS
INDIANA 46285
UNITED STATES OF AMERICA

☐ This person is also inventorTelephone No.
(317) 277 3725Facsimile No.
(317) 276 3861Teleprinter No.
276051 ELI LILLY IND AState (that is, country) of nationality:
United States of AmericaState (that is, country) of residence:
United States of America

This person is applicant for the purposes of: ☐ all designated States ☒ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is country) of residence if no State of residence is indicated below.)

PROTHERICS MOLECULAR DESIGN LIMITED
BEECHFIELD HOUSE
LYME GREEN BUSINESS PARK
MACCLESFIELD
CHESHIRE SK11 0JL
UNITED KINGDOM

This person is:

☒ applicant only
☐ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United KingdomState (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of: ☐ all designated States ☒ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on a continuation sheet.

Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE

The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:

☒ agent ☐ common representative

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.)

HAY, Martin A.
Martin A. Hay & Co.
13 Queen Victoria Street, Macclesfield
Cheshire SK11 6LP
United Kingdom

Telephone No.
+44 (0)1625 500057Facsimile No.
+44 (0)1625 500058

Teleprinter No.

☐ Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

Form PCT/RO/101 (first sheet) (July 1998; reprint January 2000)

See Notes to the request form

Continuation of Box No. III FURTHER APPLICANTS AND/OR (FURTHER) INVENTORS

If none of the following sub-boxes is used, this sheet is not to be included in the request.

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

LIEBESCHUETZ, John Walter
Laburnum Cottage, 42 Bollington Road
Bollington
Cheshire SK10 5EJ GB[▲]

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of: ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

LYONS, Amanda Jane
3 Thistleton Close
Macclesfield
Cheshire SK11 8BE GB[▲]

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of: ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

MURRAY, Christopher William
1 Wheatfield Close, Tytherington
Macclesfield
Cheshire SK10 2TT GB[▲]

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of: ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

RIMMER, Andrew David
9 Stamford Drive
Whittle-le-Woods
Chorley, Lancashire PR6 7HP GB[▲]

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of: ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on another continuation sheet.

Continuation of Box N . III FURTHER APPLICANTS AND/OR (FURTHER) INVENTORS

If none of the following sub-boxes is used, this sheet is not to be included in the request.

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

YOUNG, Stephen Clinton
8 Cranbourne Road
Heaton Moor
Stockport SK4 4LD GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

CAMP, Nicholas Paul
Flat 2, Silver Court, Fosseyway
Nailsea
Avon BS48 2BX GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

JONES, Stuart Donald
17 Oakwood Drive
Prestbury
Cheshire SK10 4HG GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

MORGAN, Phillip John
11 Woodland Avenue
Congleton
Cheshire CW12 1LN GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on another continuation sheet.

See Notes to the request form

Continuation of Box N . III

FURTHER APPLICANTS AND/OR (FURTHER) INVENTORS

If none of the following sub-boxes is used, this sheet is not to be included in the request.

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

RICHARDS, Simon James
39 Vicarage Road
Blackrod
Bolton BL6 5DA GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

☐

all designated States

☐

all designated States except the United States of America

☒

the United States of America only

☐

the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

WYLIE, William Alexander
Flat 4, 39 Station Road
Reddish
Stockport SK5 6LT GB ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United Kingdom

State (that is, country) of residence:
United Kingdom

This person is applicant for the purposes of:

☐

all designated States

☐

all designated States except the United States of America

☒

the United States of America only

☐

the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

MASTERS, John Joseph
12047 Flint Stone Court
Fishers
Indiana 46038 US ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United States of America

State (that is, country) of residence:
United States of America

This person is applicant for the purposes of:

☐

all designated States

☐

all designated States except the United States of America

☒

the United States of America only

☐

the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State(that is country) of residence if no State of residence is indicated below.)

WILEY, Michael Robert
7725 Langwood Drive
Indianapolis
Indiana 46268 US ▲

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
United States OF AMERICA ▲

State (that is, country) of residence:
United States OF AMERICA ▲

This person is applicant for the purposes of:

☐

all designated States

☐

all designated States except the United States of America

☒

the United States of America only

☐

the States indicated in the Supplemental Box

☐ Further applicants and/or (further) inventors are indicated on another continuation sheet.

ADDED
RO/GB

Box No.V DESIGNATION OF STATES

following designations are hereby made under Rule 4.9(a) (mark the applicable check-boxes; at least one must be marked):

Regional Patent

- ☒ AP ARIPO Patent: GH Ghana, GM Gambia, KE Kenya, LS Lasotho, MW Malawi, SD Sudan, SL Sierra Leone, SZ Swaziland, TZ United Republic of Tanzania, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harare Protocol and of the PCT
- ☒ EA Eurasian Patent: AM Armenia, AZ Azerbaijan, BY Belarus, KG Krygyzstan, KZ Kazakhstan, MD Republic of Moldova, RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT
- ☒ EP European Patent: AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany, DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg, MC Monaco, NL Netherlands, PT Portugal, SE Sweden, and any other State which is a Contracting State of the European Patent Convention and of the PCT
- ☒ OA OAPI Patent: BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon, GA Gabon, GN Guinea, GW Guinea-Bissau, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired, specify on dotted line)

National Patent (if other kind of protection or treatment desired, specify on dotted line):

- | | | |
|---|---|---|
| <input checked="" type="checkbox"/> AE United Arab Emirates..... | <input checked="" type="checkbox"/> LR Liberia..... | |
| <input checked="" type="checkbox"/> AL Albania..... | <input checked="" type="checkbox"/> LS Lesotho..... | |
| <input checked="" type="checkbox"/> AM Armenia..... | <input checked="" type="checkbox"/> LT Lithuania..... | |
| <input checked="" type="checkbox"/> AT Austria..... | <input checked="" type="checkbox"/> LU Luxembourg..... | |
| <input checked="" type="checkbox"/> AU Australia..... | <input checked="" type="checkbox"/> LV Latvia..... | |
| <input checked="" type="checkbox"/> AZ Azerbaijan..... | <input checked="" type="checkbox"/> MA Morocco..... | |
| <input checked="" type="checkbox"/> BA Bosnia and Herzegovina..... | <input checked="" type="checkbox"/> MD Republic of Moldova..... | |
| <input checked="" type="checkbox"/> BB Barbados..... | <input checked="" type="checkbox"/> MG Madagascar..... | |
| <input checked="" type="checkbox"/> BG Bulgaria..... | <input checked="" type="checkbox"/> MK The former Yugoslav Republic of Macedonia..... | |
| <input checked="" type="checkbox"/> BR Brazil..... | | |
| <input checked="" type="checkbox"/> BY Belarus..... | <input checked="" type="checkbox"/> MN Mongolia..... | |
| <input checked="" type="checkbox"/> CA Canada..... | <input checked="" type="checkbox"/> MW Malawi..... | |
| <input checked="" type="checkbox"/> CH and LI Switzerland and Liechtenstein..... | <input checked="" type="checkbox"/> MX Mexico..... | |
| <input checked="" type="checkbox"/> CN China..... | <input checked="" type="checkbox"/> NO Norway..... | |
| <input checked="" type="checkbox"/> CR Costa Rica..... | <input checked="" type="checkbox"/> NZ New Zealand..... | |
| <input checked="" type="checkbox"/> CU Cuba..... | <input checked="" type="checkbox"/> PL Poland..... | |
| <input checked="" type="checkbox"/> CZ Czech Republic..... | <input checked="" type="checkbox"/> PT Portugal..... | |
| <input checked="" type="checkbox"/> DE Germany..... | <input checked="" type="checkbox"/> RO Romania..... | |
| <input checked="" type="checkbox"/> DK Denmark..... | <input checked="" type="checkbox"/> RU Russian Federation..... | |
| <input checked="" type="checkbox"/> DM Dominica..... | <input checked="" type="checkbox"/> SD Sudan..... | |
| <input checked="" type="checkbox"/> EE Estonia..... | <input checked="" type="checkbox"/> SE Sweden..... | |
| <input checked="" type="checkbox"/> ES Spain..... | <input checked="" type="checkbox"/> SG Singapore..... | |
| <input checked="" type="checkbox"/> FI Finland..... | <input checked="" type="checkbox"/> SI Slovenia..... | |
| <input checked="" type="checkbox"/> GB United Kingdom..... | <input checked="" type="checkbox"/> SK Slovakia..... | |
| <input checked="" type="checkbox"/> GD Grenada..... | <input checked="" type="checkbox"/> SL Sierra Leone..... | |
| <input checked="" type="checkbox"/> GE Georgia..... | <input checked="" type="checkbox"/> TJ Tajikistan..... | |
| <input checked="" type="checkbox"/> GH Ghana..... | <input checked="" type="checkbox"/> TM Turkmenistan..... | |
| <input checked="" type="checkbox"/> GM Gambia..... | <input checked="" type="checkbox"/> TR Turkey..... | |
| <input checked="" type="checkbox"/> HR Croatia..... | <input checked="" type="checkbox"/> TT Trinidad and Tobago..... | |
| <input checked="" type="checkbox"/> HU Hungary..... | <input checked="" type="checkbox"/> TZ United Republic of Tanzania..... | |
| <input checked="" type="checkbox"/> ID Indonesia..... | <input checked="" type="checkbox"/> UA Ukraine..... | |
| <input checked="" type="checkbox"/> IL Israel..... | <input checked="" type="checkbox"/> UG Uganda..... | |
| <input checked="" type="checkbox"/> IN India..... | <input checked="" type="checkbox"/> US United States of America..... | |
| <input checked="" type="checkbox"/> IS Iceland..... | | |
| <input checked="" type="checkbox"/> JP Japan..... | <input checked="" type="checkbox"/> UZ Uzbekistan..... | |
| <input checked="" type="checkbox"/> KE Kenya..... | <input checked="" type="checkbox"/> VN Viet Nam..... | |
| <input checked="" type="checkbox"/> KG Kyrgyzstan..... | <input checked="" type="checkbox"/> YU Yugoslavia..... | |
| <input checked="" type="checkbox"/> KP Democratic People's Republic of Korea..... | <input checked="" type="checkbox"/> ZA South Africa..... | |
| | <input checked="" type="checkbox"/> ZW Zimbabwe..... | |
| <input checked="" type="checkbox"/> KR Republic of Korea..... | Check-boxes reserved for designating States which have | |
| <input checked="" type="checkbox"/> KZ Kazakhstan..... | become party to the PCT after issuance of this sheet: | |
| <input checked="" type="checkbox"/> LC St Lucia..... | <input checked="" type="checkbox"/> DZ... Algeria | <input checked="" type="checkbox"/> AG Antigua & Barbuda..... |
| <input checked="" type="checkbox"/> LK Sri Lanka..... | <input checked="" type="checkbox"/> MZ... Mozambique..... | |

Precautionary Designation Statement. In addition to the designations made above, the applicant also makes under Rule 4.9(b) all other designations which would be permitted under the PCT except the designation(s) indicated in the Supplemental Box as being excluded from the scope of this statement. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit. (Confirmation (including fees) must reach the receiving Office within the 15-month time limit.)

▲
ADDED
20/68

Supplemental Box *If the Supplemental Box is not used, this sheet need not be included in the request.*

1. If any of the Boxes, the space is insufficient to furnish all the information: in such case, write "Continuation of Box No. ..." [indicate the number of the Box] and furnish the information in the same manner as required according to the captions of the Box in which the space was insufficient, in particular:
- (i) if more than two persons are involved as applicants and/or inventors and no "continuation sheet" is available: in such case, write "Continuation of Box No. III" and indicate for each additional person the same type of information as required in Box No. III; The country of the address indicated in this Box is the applicant's State(that is, country) of residence if no State of residence is indicated below;
 - (ii) if, in Box No. II or in any of the sub-boxes of Box No. III, the indication "the States indicated in the Supplemental Box" is checked: in such case, write "Continuation of Box No. II" or "Continuation of Box No. III" or "Continuation of Boxes No. II and No. III" (as the case may be), indicate the name of the applicant(s) involved and, next to (each) such name, the State(s) (and/or, where applicable, ARIPO, Eurasian, European or OAPI patent) for the purposes of which the named person is applicant;
 - (iii) if in Box No. II or in any of the sub-boxes of Box No. III, the inventor or the inventor/applicant is not inventor for the purposes of all designated States or for the purposes of the United States of America: in such case, write "Continuation of Box No. II" or "Continuation of Box No. III" or "Continuation of Boxes No. II and No. III" (as the case may be), indicate the name of the inventor(s) and, next to (each) such name, the State(s) (and/or, where applicable, ARIPO, Eurasian, European or OAPI patent) for the purposes of which the named person is inventor;
 - (iv) if, in addition to the agent(s) indicated in Box No. IV, there are further agents: in such case, write "Continuation of Box No. IV" and indicate for each further agent the same type of information as required in Box No. IV;
 - (v) if in Box No. V, the name of any State (or OAPI) is accompanied by the indication "patent of addition," or "certificate of addition," or if, in Box No. V, the name of the United States of America is accompanied by an indication "continuation" or "continuation-in-part": in such case, write "Continuation of Box No. V" and the name of each State involved (or OAPI), and after the name of each such State (or OAPI), the number of the parent title or parent application and the date of grant of the parent title or filing of the parent application;
 - (vi) if, in Box No. VI, there are more than three earlier applications whose priority is claimed: in such case, write "Continuation of Box No. VI" and indicate for each additional earlier application the same type of information as required in Box No. VI.
 - (vii) if, in Box No. VI, the earlier application is an ARIPO application; in such case write "Continuation of Box No. VI", specify the number of the item corresponding to that earlier application and indicate at least one country party to the Paris Convention for the Protection of Industrial Property or one member of the World Trade Organisation for which that earlier application was filed.
2. If, with regard to the precautionary designation statement contained in Box No. V, the applicant wishes to exclude any State(s) from the scope of that statement; in such case, write "Designation(s) excluded from precautionary designation statement" and indicate the name or two-letter code of each State so excluded.
3. If the applicant claims, in respect of any designated Office, the benefits of provisions of the national law concerning non-prejudicial disclosures or exceptions to lack of novelty: in such case, write "Statement concerning non-prejudicial disclosures or exceptions to lack of novelty" and furnish that statement below.


Box No. VI PRIORITY CLAIM☒ Further priority claims are indicated in the Supplemental Box

Filing Date of earlier application (day/month/year)	Number of earlier application	Where earlier application is		
		national application: country	regional application:* regional Office	international application: receiving Office
item (4) 14/12/99 14-DECEMBER 1999	9929553.7	United Kingdom		

Box No. PRIORITY CLAIM		<input checked="" type="checkbox"/> Further priority claims are indicated in the Supplemental Box		
Filing Date of earlier application (day/month/year)	Number of earlier application	Where earlier application is		
		national application: country	regional application: regional Office	international application: receiving Office
item (1) 14/06/99 14 JUNE 1999	9913823.2	United Kingdom		
item (2) 02/07/99 02 JULY 1999	60/142064	United States of America		
item (3) 09/08/99 09 AUGUST 1999	9918741.1	United Kingdom		
<input checked="" type="checkbox"/> The receiving Office is requested to prepare and transmit to the International Bureau a certified copy of the earlier application(s) (only if the earlier application was filed with the Office which for the purposes of the present international application is the receiving Office) identified above as item(s) (1)(3)(4) * Where the earlier application is an ARIPO application, it is mandatory to indicate in the Supplemental Box at least one country party to the Paris Convention for the Protection of Industrial Property for which that earlier application was filed (Rule 4.10(b)(ii)). See Supplemental Box.				

Box No. VII INTERNATIONAL SEARCHING AUTHORITY		
Choice of International Searching Authority (ISA) (If two or more International Searching Authorities are competent to carry out the international search, indicate the Authority chosen; the two-letter code may be used): ISA / EPO	Request to use results of earlier search; reference to that search (if an earlier search has been carried out by or requested from the International Searching Authority): Date (day/month/year) Number Country (or regional Office)	

Box No. VIII CHECK LIST; LANGUAGE OF FILING	
This international application contains the following number of sheets: request : 7 description (excluding sequence listing part) : 244 claims : 16 abstract : 1 drawings : 0 sequence listing part of description : 0 Total number of sheets : 268	This international application is accompanied by the item(s) marked below: 1. <input type="checkbox"/> fee calculation sheet 2. <input type="checkbox"/> separate signed power of attorney. 3. <input type="checkbox"/> copy of general power of attorney; reference number, if any 4. <input type="checkbox"/> statement explaining lack of signature 5. <input checked="" type="checkbox"/> priority document(s) identified in Box No VI as item(s) 2 6. <input type="checkbox"/> translation of international application into (language); 7. <input type="checkbox"/> separate indications concerning deposited microorganisms or other biological material 8. <input type="checkbox"/> nucleotide and/or amino acid sequence listing in computer readable form 9. <input checked="" type="checkbox"/> other (specify): Form 23/77 for documents 1, 3, 4 in Box VI, cover letter
Figure of the drawings which should accompany the abstract:	Language of filing of the international application: English

Box No. IX SIGNATURE OF APPLICANT OR AGENT	
Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request)	
	Aged
Martin A. HAY	

For receiving Office use only		2. Drawings: <input type="checkbox"/> received; <input type="checkbox"/> not received;
1. Date of actual receipt of the purported international application:	13 JUN 2000 13.06.2000	
3. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application:		
4. Date of timely receipt of the required corrections under PCT Article 11(2):		
5. International Searching Authority (if two or more are competent): ISA /	6. <input checked="" type="checkbox"/> Transmittal of search copy delayed until search fee is paid	

For International Bureau use only	
Date of receipt of the record copy by the International Bureau:	

ADDED 20/09/02

Compounds

This invention relates to compounds which are inhibitors of serine proteases and to pharmaceutical compositions thereof and their use in the treatment of the human or animal body.

The serine proteases are a group of proteolytic enzymes which have a common catalytic mechanism characterized by a particularly reactive Ser residue. Examples of serine proteases include trypsin, tryptase, chymotrypsin, elastase, thrombin, plasmin, kallikrein, Complement C1, acrosomal protease, lysosomal protease, cocoonase, α -lytic protease, protease A, protease B, serine carboxypeptidase II, subtilisin, urokinase, Factor VIIa, Factor IXa, and Factor Xa. The serine proteases have been investigated extensively over a period of several decades and the therapeutic value of inhibitors of serine proteases is well understood.

Serine protease inhibitors play a central role in the regulation of a wide variety of physiological process including coagulation, fibrinolysis, fertilization, development, malignancy, neuromuscular patterning and inflammation. It is well known that these compounds inhibit a variety of circulating proteases as well as proteases that are activated or released in tissue. It is also becoming clear that serine protease inhibitors inhibit critical cellular processes, such as adhesion, migration, free radical production and apoptosis. In addition, animal experiments indicate that intravenously administered serine protease inhibitors, variants or cells expressing serine protease inhibitors, provide a protective effect against tissue damage.

Serine protease inhibitors have also been predicted to have potential beneficial uses in the treatment of disease in a wide variety of clinical areas such as oncology, neurology, haematology, pulmonary medicine, immunology, inflammation and infectious disease.

In particular serine protease inhibitors may be beneficial in the treatment of thrombotic diseases, asthma, emphysema, cirrhosis, arthritis, carcinoma, melanoma, restenosis, atheroma, trauma, shock and reperfusion injury.

Thus for example an inhibitor of Factor Xa has value as a therapeutic agent as an anticoagulant, e.g. in the treatment and prevention of thrombotic disorders. The use of a Factor Xa inhibitor as an anticoagulant is desirable in view of the selectivity of its effect. Many clinically approved anticoagulants have been associated with adverse events owing to the non-specific nature of their effects on the coagulation cascade.

Also, there are well-known associations of α_1 protease inhibitor deficiency with emphysema and cirrhosis and C1 esterase inhibitor deficiency with angioedema.

It has now been found that certain aromatic compounds carrying bulky lipophilic side chains are particularly effective as inhibitors of serine proteases, especially proteases with negatively charged P1 specificity pockets, and most especially the serine proteases thrombin, and most importantly Factor Xa. The Factor Xa inhibitors of this invention are potentially useful for the prophylaxis or treatment of thrombotic disorders such as amongst others venous thrombosis, pulmonary embolism, arterial thrombosis, myocardial ischaemia, myocardial infarction, and cerebral thrombosis. They potentially have benefit in the treatment of acute vessel closure associated with thrombolytic therapy

and restenosis, e.g. after transluminal coronary angioplasty or bypass grafting of the coronary or peripheral arteries and in the maintenance of vascular access patency in long term hemodialysis patients.

5 Factor Xa inhibitors of this invention may, with benefit, form part of a combination therapy with an anticoagulant with a different mode of action or with a thrombolytic agent.

10 Hence, the invention also comprises certain compounds which have been found to be inhibitors of both Factor Xa and thrombin. These compounds have excellent potential therapeutic value and may synergistically boost Fxa antithrombotic effect.

15 It has been reported in WO99/11658 and WO99/11657 that certain benzamidine and aminoisoquinoline derivatives carrying a bulky lipophilic side chain are excellent inhibitors of serine proteases. Unfortunately, it has since been found that benzamidine compounds of WO 99/11658 in general demonstrate poor oral bioavailability.


20 Surprisingly, it has now been found that certain other aromatic compounds also show inhibitory activity against serine proteases, in particular Factor Xa, despite the lack of the amidino or 1-aminoisoquinoline functionality previously believed to be crucial for activity as a factor
25 Xa inhibitor. Many of these compounds also possess other structural features that further distinguish them from the compounds of WO99/11658 and WO99/11657.

30 Where compounds of the invention have been tested, they have generally demonstrated superior oral bioavailability in comparison with benzamidines disclosed in WO 99/11658. Also, it has been found that the compounds of the invention perform excellently in the prothrombin time assay (PT) when

compared to aminoisoquinolines of similar factor Xa activity and structure. The PT assay is a coagulation assay and it is widely accepted that direct acting Factor Xa inhibitors which perform well in the PT assay are more likely to be good antithrombotics.

In WO99/09053 certain 2-aminobenzamide compounds are disclosed as potential motilin receptor antagonists and in US 3268513 similar 2-aminobenzamide compounds are suggested as potential antibacterial agents. However, the novel compounds of the present invention have not before been suggested as potential serine protease inhibitors.

Thus viewed from an one aspect the invention provides a serine protease inhibitor compound of formula (I)



where R_2 represents a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring atom, optionally being substituted in the 3 and/or 4 position (in relation to the point of attachment of X-X) by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO_2 - or R_1 , or the substituents at the 3 and 4 positions taken together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring optionally substituted by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} , and optionally substituted in the position alpha to the X-X group (i.e. 6 position for a six membered aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido,

aminoalkyl, alkoxy or alkylthio with the proviso that R_2 cannot be aminoisoquinolyl;

each X independently is a C, N, O or S atom or a CO, CR_{1a} , $C(R_{1a})_2$ or NR_{1a} group, at least one X being C, CO, CR_{1a} or $C(R_{1a})_2$;

each R_{1a} independently represents hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl alkoxyalkyl, alkoxy carbonyl, alkylaminocarbonyl, alkoxy carbonylamino, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

R_1 is as defined for R_{1a} , provided that R_1 is not unsubstituted aminoalkyl;

L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group;

Y (the α -atom) is a nitrogen atom or a CR_{1b} group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group, preferably containing 5 to 10 ring atoms and optionally substituted by groups R_{3a} or phenyl optionally substituted by R_{3a} ;

each R_{3a} independently is R_{1c} , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkyl imidazolyl, thiazolyl, alkyl thiazolyl, alkyl oxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, haloalkoxy and haloalkyl;

Lp is a lipophilic organic group;

D is a hydrogen bond donor group; and n is 0, 1 or 2;

and

R_{1b} , R_{1c} , and R_{1j} are as defined for R_{1a} ,

or a physiologically tolerable salt thereof, e.g. a halide, phosphate or sulphate salt or a salt with ammonium or an organic amine such as ethylamine or meglumine.

Compounds of formula I as defined above, but in which
5 R_1 is an unsubstituted aminoalkyl group are claimed in a co-pending application.

In the compounds of the invention, where the alpha atom is carbon it preferably has the conformation that would result from construction from a D- α -aminoacid
10 $NH_2-CR_{1b}(Cy)-COOH$ where the NH_2 represents part of X-X. Likewise the fourth substituent R_{1b} at an alpha carbon is preferably a methyl or hydroxymethyl group or hydrogen.

In the compounds of the invention, unless otherwise indicated, aryl groups preferably contain 5 to 10 ring atoms
15 optionally including 1, 2 or 3 heteroatoms selected from O, N and S; alkyl, alkenyl or alkynyl groups or alkylene moieties preferably contain up to 6 carbons, e.g. C_{1-6} or C_{1-3} ; cyclic groups preferably have ring sizes of 3 to 8 atoms; and fused multicyclic groups preferably contain 8 to
20 16 ring atoms.

Examples of particular values for R_{1a} are: hydrogen, methyl or ethyl. R_{1a} is preferably a hydrogen atom.

The linker group from the R_2 group to the alpha atom is preferably selected from $-CH=CH-$, $-CONH-$, $-CONR_{1a}-$, $-NH-CO-$,
25 $-NH-CH_2-$, $-CH_2-NH-$, $-CH_2O-$, $-OCH_2-$, $-COO-$, $-OC=O-$ and $-CH_2CH_2-$. Preferably, the X moiety nearest to the alpha atom is an NH or O atom, most preferably a NH group. The X moiety alpha to the aromatic ring is preferably a carbon based group such as CH_2 or CO, preferably CO. Thus a
30 particularly preferred linker X-X is $-CONH-$. In an alternative embodiment the linker is preferably a $-OCH_2-$ group.

Examples of particular values for R_{1b} are: hydrogen, (1-4C)alkyl, such as methyl or hydroxy(1-4C)alkyl, such as hydroxymethyl. R_{1b} is preferably a hydrogen atom.

The alpha atom (Y) is preferably a CH or C(CH₃) group,
5 especially CH.

The linker group from the alpha atom to the lipophilic group is preferably CO, CH₂NH, CONR_{1d}(CH₂)_m, (CH₂)_mN(R_{1d})CO(CH₂)_m, (CH₂)_{m+2}, CO(CH₂)_m, (CH₂)_mCO, (CH₂)_mOC=O, (CH₂)_mO, CH=CH(CH₂)_m, SO₂, SO₂NR_{1d}, SO₂(CH₂)_m,
10 (CH₂)_mSO₂ or (CH₂)_mSO₂NR_{1d} (where each m is independently 0 or 1 and R_{1d} is as defined for R_{1a}).

Examples of particular values for R_{1d} are: hydrogen; for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-6C)alkyl, such as methyl
15 or ethyl, or aryl(1-6C)alkyl, such as benzyl or phenylethyl; for aminoalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: (2-6C)carboxamido, such as carboxamidomethyl; for hydroxyalkyl optionally substituted by hydroxy,
20 alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-6C)carboxyalkyl, such as carboxymethyl, carboxyethyl or carboxypropyl; for alkoxyalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-
25 5C)alkoxycarbonyl(1-6C)alkyl, such as methoxycarbonylmethyl, methoxycarbonylethyl, methoxycarbonylpropyl, ethoxycarbonylmethyl, ethoxycarbonylethyl and ethoxycarbonylpropyl.

R_{1d} is preferably a hydrogen atom.

30 The linker may be optionally branched, for example, to incorporate a polar functionality.

Examples of particular values for L are CO, CONH, CH₂NHCO and CONHCH₂.

It will be appreciated by those skilled in the art that a diverse range of organic groups are lipophilic, and that it is therefore impractical to define with precision each and every structure that may be incorporated into a serine protease inhibitor according to the invention. Accordingly, it is being assumed that the addressee of this specification will not require an exhaustive computer listing of structures of lipophilic groups, but will instead make use of the structures of lipophilic groups disclosed in the specification, especially those exemplified; the test systems described herein for identifying serine protease inhibitors; and common general knowledge of the lipophilicity, synthesis and stability of organic compounds, to obtain novel serine protease inhibitor compounds of formula (I).

The lipophilic group may be, for example, an alkyl, alkenyl, carbocyclic or heterocyclic group, or a combination of two or more such groups linked by a spiro linkage or a single or double bond or by C=O, O, S, SO, SO₂, CONR_{1e}, NR_{1e}-CO-, NR_{1e} linkage (where R_{1e} is as defined for R_{1a}), optionally substituted by one or more oxo or R₃ groups in which R₃ is as defined for R_{3a}.

By way of illustration, representative lipophilic groups include methylcyclohexyl, methylcyclohexylmethyl, methylphenylmethyl, phenylethyl, benzylpiperidinyl, benzoylpiperidinyl, bispiperidinyl and phenylpiperazinyl.

Phenylethyl is an example of a combination of an alkyl group and a carbocyclic group linked through a single bond.

Benzylpiperidinyl is an example of a combination of an alkyl group, a carbocyclic group and a heterocyclic group linked by single bonds.

5 Benzoylpiperidinyl is an example of a combination of a carbocyclic group and a heterocyclic group linked through C=O.

Methylcyclohexylmethyl is an example of a combination of an alkyl group (methyl) and a carbocyclic group (cyclohexyl) linked by a single bond and having a
10 substituent R_3 (methyl) on cyclohexyl. It will be appreciated that this group could alternatively have been regarded as a combination of two alkyl groups and a carbocyclic group. However, in order to provide clarity, in this specification any terminal alkyl group in Lp will be
15 treated as a substituent R_3 .

When the lipophilic group comprises an alkyl group, this may be, for example, a (1-3C) alkyl group, such as methyl, ethyl or propyl. Preferably an alkyl group is unsubstituted.

20 When the lipophilic group comprises a carbocyclic group, this may be, for example, a non-aromatic or aromatic, mono or polycyclic hydrocarbon group containing up to 25, more preferably up to 10 carbon atoms. The carbocyclic group may thus be, for example, a cycloalkyl, polycycloalkyl, phenyl or naphthyl group, or a cycloalkyl group fused with a
25 phenyl group.

Examples of particular values for a cycloalkyl group are (3-6C) cycloalkyl groups, such as cyclopentyl and cyclohexyl. A cycloalkyl group is preferably unsubstituted
30 or substituted by one group R_3 , preferably amino or an alkyl group, such as methyl.

Examples of particular values for a polycycloalkyl group are (6-10C) polycycloalkyl groups, such as bicycloalkyl, for example decalinyl, norbornyl or adamantyl. A polycycloalkyl group is preferably unsubstituted.

- 5 A phenyl group is preferably unsubstituted or substituted by one or two R_3 groups. More preferably it is substituted by one or two R_3 groups.

A naphthyl group is preferably unsubstituted or substituted by one R_3 group.

- 10 Examples of a cycloalkyl or cycloalkenyl group fused with a phenyl group are indanyl and tetrahydronaphthyl. This group is preferably unsubstituted.

- When the lipophilic group comprises a heterocyclic group, this may be, for example, a non-aromatic or aromatic,
15 mono or polycyclic group containing one or two oxygen, nitrogen or sulfur atoms in the ring system, and in total up to 25, more preferably up to 10 ring system atoms.

- Examples of a heterocyclic group when it is a non-aromatic monocyclic group are azacycloalkyl groups, such as
20 pyrrolidinyl and piperidinyl; azacycloalkenyl groups, such as pyrrolinyl; diazacycloalkyl groups, such as piperazinyl; oxacycloalkyl groups, such as tetrahydropyranyl; and thiacycloalkyl groups, such as tetrahydrothiopyranyl. A non-aromatic monocyclic group preferably contains 5, 6 or 7 ring
25 atoms and is preferably unsubstituted or substituted by one group R_3 , preferably alkyl, such as methyl or ethyl, or hydroxyalkyl, such as hydroxymethyl.

- Examples of a heterocyclic group when it is a non-aromatic polycyclic group are bicyclic groups, such as
30 azacycloalkyl fused with phenyl, for example dihydroindolyl, dihydroisoindolyl, tetrahydroquinolyl and tetrahydroisoquinolyl; and tricyclic groups, such as

azacycloalkyl fused with indolyl, for example tetrahydropyrido[3,4-b]indole. This group is preferably unsubstituted.


5 Examples of a heterocyclic group when it is a aromatic monocyclic group are furyl, pyrrolyl, thienyl, imidazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, oxazolyl, oxadiazolyl (such as 1,3,4-oxadiazolyl), thiadiazolyl (such as 1,3,4-thiadiazolyl) and thiazolyl. This group is preferably unsubstituted or substituted by one R_3 .

10 Examples of a heterocyclic group when it is an aromatic polycyclic group are bicyclic groups such as benzofuryl, quinolinyl, isoquinolinyl and benzothienyl. This group is preferably unsubstituted or substituted by one R_3 .

15 The lipophilic group preferably comprises a cycloalkyl, azacycloalkyl, diazacycloalkyl, phenyl, naphthyl, adamantyl, bicycloalkyl, mono- or diazabicycloalkyl, mono- or bicyclo heteroaromatic or a linear or branched alkyl or alkenyl group all optionally substituted by one or more groups R_3 , or a combination of at least two such groups linked by a
20 spiro linkage or a single or double bond or by C=O, O, S, SO, SO₂, CONR_{1e}, NR_{1e}-CO- or NR_{1e} linkage (where R_{1e} is as defined for R_{1a}).


Where Lp comprises a combination of at least two groups, it preferably comprises a combination of two or
25 three such groups. The groups are preferably linked by a single bond, C=O, O or NR_{1e}.

Of particular interest are compounds of formula I in which Lp comprises an azacycloalkyl or diazacycloalkyl group of formula



in which r is 1 or 2, one of X_a and X_b is N and the other is CH or N, provided that when r is 1, X_a and X_b are not both N.

- 5 Preferred compounds comprising this group are those in which L_p is a group of formula:



in which:

r is 1 or 2;

- 10 one of X_a and X_b is N and the other is CH or N provided that when r is 1, X_a and X_b are not both N;

s , t and u are each 0 or 1;

L_a and L_b are each independently selected from a single bond, $C=O$, O and NR_{1e} , in which R_{1e} is hydrogen or (1-


- 15 6C)alkyl;

G is (1-6C)alkanediyl; and

R_{10} is (1-6C)alkyl; (3-6C)cycloalkyl which is unsubstituted or substituted by (1-6C)alkyl; indanyl; pyridyl;

tetrahydropyranyl; tetrahydrothiopyranyl; phenyl which is

- 20 unsubstituted or substituted by one or two R_3 groups; pyrrolinyl; or a group of formula



in which v is 1, 2 or 3; one of X_c and X_d is N and the other is CH or N, provided that when v is 1, X_c and X_d are not

both N; and R_{11} is hydrogen, (1-6C)alkyl or when X_d is CH, hydroxy(1-6C)alkyl; provided that when t is 0, the sum of s and u is 1; when X_b is N, L_a is a bond or C=O; when X_c is N, L_b is a bond or C=O; when X_b and X_c are both N, t is 1; and
 5 when $(L_a)_s-(G)_t-(L_b)$ represents an alkyl group and X_b and X_c both represent N, the alkyl group contains at least two chain carbon atoms.

It will be appreciated that the provisos exclude compounds having two heteroatoms bonded directly together or
 10 separated by an alkyl group having only one carbon atom in the chain.

When X_a is N, L is preferably CO or CH_2CO .

When X_a is CH, L is preferably CONH, $CONHCH_2$ or CH_2NHCO .

15 Examples of values for G are CH_2 , $(CH_2)_2$ and $(CH_2)_3$.

Examples of values for R_{11} are hydrogen, methyl, ethyl or 2-propyl, or when X_d is CH, hydroxymethyl.

Examples of particular values for R_3 are:-
 hydrogen;

20 hydroxyl;

for alkoxy: methoxy or ethoxy;

for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-6C)alkyl, such as methyl, ethyl, propyl, 2-propyl, butyl, 2-butyl, t-butyl,

25 pentyl, 2-pentyl or 3-pentyl, (1-6C)alkylamino(1-6C)alkyl, such as isopropylaminomethyl, dimethylamino-methyl, diethylaminomethyl or dimethylaminoethyl, or (1-6C)alkanoyl, such as acetyl;

for hydroxyalkyl optionally substituted by hydroxy,

30 alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-

- 6C)hydroxyalkyl, such as hydroxymethyl or hydroxyethyl,
carboxy or carboxy(1-5C)alkyl;
for alkoxyalkyl: methoxymethyl;
for alkoxycarbonyl: methoxycarbonyl or ethoxycarbonyl;
5 for alkylaminocarbonyl: methylaminocarbonyl or
dimethylaminocarbonyl;
for aminoalkyl optionally substituted by hydroxy,
alkylamino, alkoxy, oxo, aryl or cycloalkyl: aminomethyl,
aminocarbonyl or aminocarbonyl(1-5C)alkyl;
10 for alkylamino optionally substituted by hydroxy,
alkylamino, alkoxy, oxo, aryl or cycloalkyl: methylamino,
dimethylamino, ethylamino, formylamino or acetylamino;
amino;
for halo: fluoro or chloro;
15 cyano;
nitro;
thiol;
for alkylthio: methylthio;
for alkylsulphonyl: methylsulphonyl, ethylsulphonyl or
20 isopropylsulphonyl;
for alkylsulphenyl: methylsulphenyl;
for triazolyl: 1,2,4-triazol-2-yl, 1,2,4-triazol-4-yl or
1,2,3-triazol-4-yl;
for imidazolyl: 1,3-imidazol-1-yl or 1,3-imidazol-4-yl;
25 for tetrazolyl: tetrazol-1-yl or tetrazol-5-yl;
for alkylsulphonamido: methylsulphonamido, ethylsulphonamido
or propylsulphonamido;
for alkylaminosulphonyl: methylaminosulphonyl,
ethylaminosulphonyl or propylaminosulphonyl;
30 aminosulphonyl;
for haloalkoxy: trifluoromethoxy; and
for haloalkyl: trifluoromethyl or trichloromethyl.

Examples of particular values for R_{1e} are hydrogen and (1-6C)alkyl, such as methyl or ethyl.

Examples of values for R_{10} are:

for (1-6C)alkyl: methyl, ethyl, 2-propyl and 3-pentyl;

5 for (3-6C)cycloalkyl which is unsubstituted or substituted by (1-6C)alkyl: cyclopentyl, 3-methylcyclopentyl, cyclohexyl and 4-methylcyclohexyl;

for indanyl: 2-indanyl;

for pyridyl: pyrid-2-yl, pyrid-3-yl and pyrid-4-yl;

10 for tetrahydropyranyl: tetrahydropyran-4-yl;

for tetrahydrothiopyranyl: tetrahydrothiopyran-4-yl;


for phenyl which is unsubstituted or substituted by one or two R_3 groups: phenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-(methylthio)phenyl, 2-ethylphenyl, 2-

15 methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-methanesulphonylphenyl, 3-methanesulphonylphenyl, 4-methanesulphonylphenyl, 4-fluoro-2-methanesulphonylphenyl, 4-amino-2-methanesulphonylphenyl, 4-amido-2-

20 2-aminosulphonylphenyl, 2-methylaminosulphonylphenyl, 2-dimethylaminosulphonylphenyl, 2-methylsulphonylamino-phenyl, 2-carboxamidophenyl and 2-acetamidophenyl;

for pyrrolinyl: pyrrolin-1-yl; and

for a group of formula




25

piperidin-1-yl, 4-methyl-piperidin-1-yl, piperidin-4-yl, 1-methylpiperidin-4-yl, 1-(2-propyl)piperidin-4-yl, pyrrolidin-1-yl, 3-methylpyrrolidin-1-yl, pyrrolidin-3-yl, 1-methyl-pyrrolidin-3-yl, 1-(2-propyl)pyrrolidin-3-yl, 1-

methyl-piperazin-4-yl, 1-ethylpiperazin-4-yl, 1-(2-propyl)piperazin-4-yl, hexahydro-1,4-diazapin-1-yl and 4-methyl-hexahydro-1,4-diazapin-1-yl.

A preferred sub-group of compounds of formula I is that
5 in which -L-Lp(D)_n is



q is 1 or 2;


(a) Q is a direct bond; and R_q is piperidin-4-yl which may bear a C₁₋₃alkyl substituent at the 1-position; or R_q is
10 NR_aR_b in which each of R_a and R_b independently is hydrogen or C₁₋₃alkyl; or one of R_a and R_b is hydrogen or methyl and the other of R_a and R_b is -CH₂-R_c or -CH₂-R_d in which R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, CONH₂, SO₂NH₂, methylaminosulphonyl,
15 dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and in which R_d is isopropyl or cyclopentyl, or NR_aR_b is pyrrolidino, piperidino, morpholino, piperazino, or tetrahydro-1,4-diazepino in which a pyrrolidino or piperidino may be a 3,4-didehydro
20 derivate and in which a pyrrolidino, piperidino, piperazino, or tetrahydro-1,4-diazepino may bear a methyl group at the 4-position (preferably R_q is piperidin-4-yl which may bear a (1-3C)alkyl substituent at the 1-position);

(b) Q is -O- or -NH-; and R_q is R_c which is
25 defined as above (R_c is preferably pyrid-2-yl, pyrid-3-yl or pyrid-4-yl); or

(c) Q is methylene; and R_q is NR_aR_b which is defined as above.

q is preferably 2.

Another sub-group of compounds is that in which
-L-Lp(D)_n is




- 5 in which R_r is -(CH₂)_c-R_c, -CHR_eR_f, -CH₂-CHR_eR_f, or R_g in which c is 1 or 2 and R_c is defined as above; each of R_e and R_f independently is hydrogen or C₁₋₃alkyl; or CHR_eR_f is cyclopentyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position), cyclohexyl (which may
10 bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position), tetrahydropyran-4-yl, tetrahydrothiopyran-4-yl, pyrrolidin-3-yl (which may bear a 1-methyl substituent), piperidin-4-yl (which may bear a 1-methyl substituent), or indan-2-yl; and R_g is 2-methylsulphonylphenyl which may bear
15 a 4-fluoro substituent or R_g is λ⁶-1,1-dioxobenzo[b]thiophen-7-yl.

Preferably c is 2.

Preferably R_c is pyrid-2-yl, pyrid-3-yl or pyrid-4-yl.

- Another sub-group of compounds of formula I is that in
20 which -L-Lp(D)_n is




in which q is 1 or 2;

s is 0 or 1; and

- R_s is -(CH₂)_c-R_c, -CHR_eR_f, or -CH₂-CHR_eR_f each of which
25 is defined as above.


Preferably s is 1.

Yet another sub-group of compounds of formula I is that in which $-L-Lp(D)_n$ is



- 5 in which R_t is piperidin-4-yl, piperidin-3-yl or pyrrolidin-3-yl (especially piperidin-4-yl), any of which may bear a C_{1-3} alkyl substituent at the 1-position (preferably methyl, ethyl or, more preferably, 2-propyl); or R_t is phenyl (which phenyl may bear a fluoro, chloro, C_{1-4} alkyl, methoxy or
- 10 methylsulphonyl substituent).

A further sub-group of compounds of formula I is that in which $-L-Lp(D)_n$ is




- 15 in which Het is a divalent 5 membered heteroaromatic group containing 1, 2 or 3 heteroatoms selected from O, N and S and having the two ring atoms at which it is connected separated by one ring atom;

h is 0 or 1; and

- 20 R_h is phenyl which may bear one or more R_3 substituents, for example independently selected from, for an ortho or a para substituent: C_{1-5} alkyl, fluoro, chloro, difluoromethyl, trifluoromethyl, methoxy, dimethylamino, methylsulphonyl, and C_{1-2} acyl, and for a meta substituent: fluoro, chloro and methyl.

- 25 Within this sub-group, a particularly preferred group of compounds is that in which $-L-Lp(D)_n$ is














in which R_h is phenyl which may bear one or two R_3 substituents, for example an ortho and/or a para substituent independently selected from, for an ortho: methyl, fluoro, chloro, methylsulphonyl and acetyl, and for a para substituent: methyl, fluoro, chloro, methoxy and dimethylamino;

Z_1 is S, Z_2 is CH, h is 0; or


Z_1 is NH, Z_2 is N, h is 1.


Most preferably, the lipophilic group Lp is selected from






5








5




10









5









10
























5





10











wherein R₃ is as hereinbefore defined;

5 m represents 0 or 1;

R₄ represents hydrogen, (CH₂)_wCOOH or (CH₂)_wCONH₂;

w represents an integer from 0 to 4; and


X represents CH or N.

Where two or more X atoms are present in a ring,
10 preferably at least one is CH.

When R₃ is present as a substituent on an aromatic ring, it is preferably selected from hydrogen, alkylsulphonyl, aminosulphonyl, alkylaminosulphonyl, alkylaminocarbonyl, amino, amido, alkoxycarbonyl,
15 acetylamino, chloro, fluoro, cyano, methoxy, ethoxy, nitro, hydroxy, alkylsulphonylamino, triazolyl and tetrazolyl.


When R₃ is present as a substituent on a saturated ring, it is preferably selected from hydrogen, hydroxy, amino, (1-3C)alkoxy, (1-3C)hydroxyalkyl, (1-3C)alkyl,
20 carboxy, methoxycarbonyl and ethoxycarbonyl.

One group of lipophilic groups Lp is that of formula





in which L_x represents O or NH.

For example specific lipophilic groups include




5





where R_8 is as defined for R_3 (preferably as defined for a
 5 substituent on an aromatic ring), especially where R_8
 represents H, OMe, SO_2Me , F, cyano, amido, amino, NO_2 , Cl or
 OH; and R_1 is hydrogen or (1-6C)alkyl (such as methyl, ethyl
 or 2-propyl).

Another highly preferred lipophilic group is of formula
 10 (DP)




(DP)

wherein A represents N or CH (preferably N) and R_3 is as
 15 hereinbefore defined. When the lipophilic group is (DP) it
 is preferred that the group L represents CO, CH_2 or SO_2 .
 Also, it is preferred if the R_3 groups in the formula DP are
 hydrogen.

(J)

Another highly preferred lipophilic group is based on the formula (K)



(K)

Preferably X₂ is hydrogen or fluoro. Compounds in which the lipophilic group is based on the formula (K) or (J) have been found to perform relatively well in the prothrombin time assay, when compared with corresponding aminoisoquinolines of WO99/11657.

The hydrogen bond donor group which may be attached to the lipophilic group preferably has a nitrogen or oxygen atom as the hydrogen bearing donor atom and conveniently is a hydroxyl group, a primary, secondary or tertiary amine, or a primary or secondary imine group (as part of an amidine or guanidine) or a saturated or unsaturated heterocyclic group containing a ring nitrogen, preferably a group containing 5 to 7 ring atoms. Where the donor atom is a ring nitrogen, the remote portion of the heterocyclic ring may be part of the lipophilic group.

The cyclic group attached to the alpha carbon is preferably an optionally R_{3a} substituted phenyl, pyridyl (such as pyrid-2-yl, pyrid-3-yl or pyrid-4-yl), thienyl (such as thien-2-yl or thien-3-yl), thiazolyl (such as thiazol-2-yl, thiazol-4-yl or thiazol-5-yl), naphthyl (such as naphth-1-yl), piperidiny (such as piperidin-4-yl) or cycloalkyl, such as a cyclohexyl group.

Examples of particular values for R_{3a} are:-

- hydrogen;
- 10 hydroxyl;
- for alkoxy: methoxy or ethoxy;
- for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or ethyl, or alkylaminoalkyl, such as methylaminomethyl or
- 15 dimethylaminomethyl;
- for hydroxyalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: hydroxymethyl or carboxy;
- for alkoxyalkyl: methoxymethyl;
- 20 for alkoxycarbonyl: methoxycarbonyl or ethoxycarbonyl;
- for alkylaminocarbonyl: methylaminocarbonyl or dimethylaminocarbonyl;
- for aminoalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: aminomethyl;
- 25 $CONH_2$ or CH_2CONH_2 ;
- for alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-6C)alkanoylamino, such as acetylamino;
- for alkoxycarbonylamino: methoxycarbonylamino, ethoxycarbonylamino or t-butoxycarbonylamino;
- 30 amino;
- for halo: fluoro or chloro;

cyano;

nitro;

thiol;

for alkylthio: methylthio;

5 for alkylsulphonyl: methylsulphonyl or ethylsulphonyl;

for alkylsulphenyl: methylsulphenyl;

for alkylsulphonamido: methylsulphonylamido or
ethylsulphonylamido;

10 for alkylaminosulphonyl: methylaminosulphonyl or
ethylaminosulphonyl;

aminosulphonyl;

for haloalkoxy: trifluoromethoxy; and

for haloalkyl: trifluoromethyl.

Examples of particular values for R_{1C} are:

15 hydrogen;

hydroxyl;

for alkoxy: methoxy or ethoxy;

for alkyl optionally substituted by hydroxy, alkylamino,
alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or

20 ethyl, or alkylaminoalkyl, such as methylaminomethyl or
dimethylaminomethyl;

for hydroxyalkyl: hydroxymethyl;

for alkoxyalkyl: methoxymethyl;

for alkoxycarbonyl: methoxycarbonyl or ethoxycarbonyl;

25 for alkylaminocarbonyl: methylaminocarbonyl or
dimethylaminocarbonyl;

for alkoxycarbonylamino: methoxycarbonylamino,
ethoxycarbonylamino or t-butoxycarbonylamino;

for alkylamino optionally substituted by hydroxy,

30 alkylamino, alkoxy, oxo, aryl or cycloalkyl: (1-
6C)alkanoylamino, such as acetylamino; and

for aminoalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: aminomethyl, CONH_2 or CH_2CONH_2 .


Preferably R_{3a} is hydrogen, hydroxyl, methoxy, methyl, amino, fluoro, chloro, ethylsulphonylamino, amido or methylaminocarbonyl.

Examples of particular values for Cy are phenyl, 4-aminophenyl, 4-amidophenyl, 4-(N-methyl)amidophenyl, 4-(N,N-dimethyl)amidophenyl, 2-chlorophenyl, 2-methylphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-methoxyphenyl, 4-methoxyphenyl, 4-carboxyphenyl, 3-ethylsulphonylamino, thien-2-yl, thien-3-yl, thiazol-4-yl, thiazol-5-yl, 2-methylthiazol-4-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, piperidin-4-yl, 1-methylpiperidin-4-yl, cyclohexyl and naphth-1-yl.

Referring to the group R_2 , examples of a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring atom are phenyl; pyrrolyl, such as 2-pyrrolyl; pyridyl, such as 3-pyridyl; pyrazinyl, such as 2-pyrazinyl; furyl, such as 2-furyl; and thienyl, such as 2-thienyl or 3-thienyl. Preferably the ring is interrupted (i.e. a carbon atom is replaced) by at most one heteroatom. More preferably the ring is phenyl, 2-thienyl or 2-pyrrolyl. Most preferably, the ring is phenyl.

When the ring is phenyl, the group R_2 may be a group of

formula



in which R₅ is amino, hydroxy or hydrogen, and R₆ and R₇ which may be the same or different represent halo, nitro, thiol, cyano, haloalkyl, haloalkoxy, amido, hydrazido, amino, alkylthio, alkenyl, alkynyl or R₁ or taken together
5 form a 5 or 6 membered fused carbocyclic ring or 5 membered heterocyclic ring, which may itself be substituted by R_{1j}, amino, halo, cyano, nitro, thiol, alkylthio, haloalkyl, haloalkoxy.

When the substituents at the 3 and 4 positions taken
10 together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring, examples of the resultant bicyclic ring are naphthyl, such as 2-naphthyl; benzimidazolyl, such as benzimidazol-5-yl or benzimidazol-6-yl; isoquinolinyl, such as isoquinolin-7-yl; indolyl, such
15 as indol-2-yl, indol-5-yl or indol-6-yl; indazolyl, such as indazol-5-yl; indazol-6-yl; 3,4-methylenedioxyphenyl; dihydroindolyl, such as 2,3-dihydroindol-6-yl; benzothiazolyl, such as benzothiazol-2-yl or benzothiazol-6-yl; benzo[b]thiophenyl, such as benzo[b]thiophen-2-yl;
20 benzofuryl, such as benzofur-2-yl; imidazo[1,2-a]pyrimidinyl, such as imidazo[1,2-a]pyrimidin-2-yl; tetrahydroimidazo[1,2-a]pyrimidinyl, such as tetrahydroimidazo[1,2-a]pyrimidin-2-yl; and benzisoxazolyl, such as benzisoxazol-5-yl.

25 R₂ preferably represents:

(i) phenyl optionally being substituted in the 3 and/or 4 position by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or
30 difluoromethoxy, carboxy, acyloxy, MeSO₂- or R₁, and optionally substituted at the 6 position by amino, hydroxy,

halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio;

(ii) naphth-2-yl optionally substituted at the 6 or 7 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} and optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or benzisoxazol-5-yl optionally substituted at the 3 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} ;

(iv) benzimidazol-5-yl or benzothiazol-6-yl optionally substituted at the 2 position by amino;

(v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_1 ;

(vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;

(vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

(viii) pyrazol-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_1 ;

(ix) pyrid-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_1 ;

(x) pyrid-3-yl optionally substituted at the 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_1 ;

(xi) benzofur-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by alkyl and optionally substituted at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(xiii) indol-6-yl substituted at the 5 position by amino, hydroxy, halo (such as fluoro or chloro), alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio and optionally substituted at the 3 position by halo (such as chloro), haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}.

Examples of particular values for substituents that may be present on R₂ are:

for halo: fluoro, chloro, bromo or iodo;
nitro;

thiol;

for haloalkoxy: difluoromethoxy or trifluoromethoxy;
hydrazido;

for alkylhydrazido: methylhydrazido;
amino;

cyano;

for haloalkyl: trifluoromethyl;

for alkylthio: methylthio;

- for alkenyl: vinyl;
for alkynyl: ethynyl;
for acylamino: acetyl amino;
carboxy;
5 for acyloxy: acetoxy;
hydroxy;
for alkyl: methyl or ethyl;
amido (CONH_2);
for aminoalkyl: aminomethyl; and
10 for alkoxy: methoxy or ethoxy.
- Examples of particular values for R_1 are:
- hydrogen;
hydroxy;
for alkoxy: methoxy or ethoxy;
15 for alkyl optionally substituted by hydroxy, alkyl amino,
alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or
ethyl, alkyl aminoalkyl, such as dimethyl aminomethyl, or
alkanoyl, such as acetyl;
for hydroxyalkyl: hydroxymethyl;
20 for alkoxyalkyl: methoxymethyl;
for alkoxy carbonyl: methoxy carbonyl;
for alkyl aminocarbonyl: methyl aminocarbonyl;
for alkyl amino: methyl amino, ethyl amino or dimethyl amino;
for hydroxyalkyl substituted by hydroxy, alkyl amino, alkoxy,
25 oxo, aryl or cycloalkyl: carboxyl or carboxymethyl; and
for aminoalkyl substituted by hydroxy, alkyl amino, alkoxy,
oxo, aryl or cycloalkyl: amido (CONH_2) or amidomethyl.

Examples of particular values for R_{1j} are:

- hydrogen;
30 hydroxy;
for alkoxy: methoxy or ethoxy;

for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or ethyl, or alkanoyl, such as acetyl;

for hydroxyalkyl: hydroxymethyl;

5 for alkoxyalkyl: methoxymethyl;

for alkoxycarbonyl: methoxycarbonyl;

for alkylamino: methylamino, ethylamino or dimethylamino;

for hydroxyalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: carboxyl or carboxymethyl; and

10 for aminoalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: amido (CONH₂) or amidomethyl.

More preferably R₂ represents:

(i) phenyl optionally being substituted in the 3 and/or 4 position by fluoro, chloro, bromo, iodo, nitro,

15 difluoromethoxy, trifluoromethoxy, amino, cyano, trifluoromethyl, methylthio, vinyl, carboxy, acetoxy, MeSO₂-, hydroxy, methoxy, ethoxy, methyl, methoxycarbonyl, methylamino, ethylamino or amido, and optionally substituted at the 6 position by amino, hydroxy, fluoro,

20 methoxycarbonyl, cyano or aminomethyl (preferably phenyl substituted in the 4 position by chloro, amino, vinyl, methylamino, methyl or methoxy, optionally at the 3 position with amino or hydroxy, and optionally at the 6 position with amino or hydroxy);

25 (ii) naphth-2-yl optionally substituted at the 6, position by hydroxy and optionally substituted at the 3 position by amino or hydroxy;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or

30 benzisoxazol-5-yl optionally substituted at the 3 position by chloro, bromo, amino, methyl or methoxy (preferably

indol-6-yl optionally substituted at the 3 position by chloro, bromo, methyl or methoxy);

(iv) benzimidazol-5-yl or benzothiazol-6-yl optionally substituted at the 2 position by amino;

5 (v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by methylthio, methyl or acetyl;

(vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;

10 (vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

(viii) pyrazol-2-yl substituted at the 5 position by methyl;

(ix) pyrid-2-yl optionally substituted at the 6
15 position by chloro;

(x) pyrid-3-yl optionally substituted at the 4 position by chloro;

(xi) benzofur-2-yl optionally substituted at the 3 position by chloro, methyl or methoxy, at the 5 or 6
20 position by methyl and at the 6 position by methoxy;

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by methyl and optionally substituted at the 5 or 6 position by fluoro, chloro, bromo, methyl or methoxy;

(xiii) indol-6-yl substituted at the 5 position by
25 chloro, fluoro or hydroxy and optionally substituted at the 3 position by chloro or methyl; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the 3 position by fluoro, chloro or methyl, and optionally substituted at the 5 or 6 position by fluoro, chloro,
30 methyl, hydroxy, or methoxy.

Examples of particular values for R_2 are:

(i) phenyl, 2-aminophenyl, 3-aminophenyl, 2-amino-3-fluorophenyl, 2-amino-4-fluorophenyl, 2-amino-4-chlorophenyl, 2-amino-3-bromophenyl, 2-amino-3-nitrophenyl, 2-amino-4-nitrophenyl, 3,4-dimethoxy-5-aminophenyl, 2-amino-4-methylphenyl, 2-amino-3-methylphenyl, 2-amino-3-methoxyphenyl, 3,4-diaminophenyl, 3,5-diaminophenyl, 3-amino-4-fluorophenyl, 3-amino-4-chlorophenyl, 3-amino-4-bromophenyl, 3-amino-4-hydroxyphenyl, 3-amino-4-carboxymethylphenyl, 3-amino-4-methylphenyl, 3-amino-4-methoxyphenyl, 2-fluorophenyl, 4-fluoro-3-cyanophenyl, 3-chlorophenyl, 3-chloro-4-hydroxyphenyl, 3-chloro-5-hydroxyphenyl, 4-chlorophenyl, 4-chloro-2-hydroxyphenyl, 4-chloro-3-hydroxyphenyl, 4-chloro-3-methylphenyl, 4-chloro-3-methoxyphenyl, 4-bromophenyl, 4-bromo-3-methylphenyl, 4-iodophenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 3-cyano-5-aminophenyl, 2-hydroxyphenyl, 2-hydroxy-4-methoxyphenyl, 3-hydroxyphenyl, 3-hydroxy-4-methylphenyl, 2,4-dihydroxyphenyl, 3,4-dihydroxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-difluoromethoxyphenyl, 4-trifluoromethoxyphenyl, 4-trifluoromethylphenyl, 4-methylthiophenyl, 4-methoxycarbonylphenyl, 4-acetoxyphenyl, 4-methanesulfonylphenyl, 3-methylphenyl, 3-methyl-5-aminophenyl, 4-methylphenyl, 4-vinylphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-methoxy-3-chlorophenyl, 4-methoxy-3-methylphenyl, 3-methylaminophenyl, 4-methylaminophenyl, 4-ethylaminophenyl or 2-aminomethylphenyl;

(ii) naphth-2-yl, 3-aminonaphth-2-yl, 3-hydroxynaphth-2-yl or 6-hydroxynaphth-2-yl;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, 3-chloroindol-6-yl, 3-bromoindol-6-yl, 3-methylindol-6-yl, 3-methoxyindol-6-yl, indazol-5-yl, 3-aminoindazol-5-yl, indazol-6-yl, benzothiazol-6-yl, 3-aminobenzisoxazol-5-yl;

(iv) benzimidazol-5-yl, 2-aminobenzimidazol-5-yl, or benzothiazol-6-yl;

(v) thien-2-yl, 5-methylthien-2-yl, 5-methylthio-thien-2-yl, 5-acetylthien-2-yl or thien-3-yl;

5 (vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;

(vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

10 (viii) 5-methylpyrazol-2-yl;

(ix) 5-chloropyrid-2-yl;

(x) pyrid-3-yl, 6-chloropyrid-3-yl;

(xi) benzofur-2-yl, 5-chlorobenzofur-2-yl, 3-methylbenzofur-2-yl, 5-methylbenzofur-2-yl, 6-


15 methoxybenzofur-2-yl;

(xii) indol-2-yl, 5-fluoroindol-2-yl, 5-chloroindol-2-yl, 5-methylindol-2-yl, 5-methoxyindol-2-yl, 6-methoxyindol-2-yl and 1-methyl-indol-2-yl;

(xiii) 5-fluoroindol-6-yl; or

20 (xiv) benzo[b]thiophen-2-yl, 5-chlorobenzo[b]thiophen-2-yl or 6-chlorobenzo[b]thiophen-2-yl.

In one embodiment the aromatic R₂ group is an optionally substituted phenyl, naphthyl, indolyl or isoindolyl group and accordingly, preferred compounds of the
25 invention are of formula (II)



(II)

wherein R_5 is amino, hydroxy or hydrogen, and R_6 and R_7 which may be the same or different represent halo, nitro, thiol, cyano, haloalkyl, haloalkoxy, amido, hydrazido, amino, alkylthio, alkenyl, alkynyl or R_1 or taken together
5 form a 5 or 6 membered fused carbocyclic ring or 5 membered heterocyclic ring, which may itself be substituted by R_{1j} , amino, halo, cyano, nitro, thiol, alkylthio, haloalkyl, haloalkoxy;

Ar is an unsubstituted or substituted aryl group,
10 preferably phenyl;

X-X is -CONH-, -CH₂CH₂-, CH₂O-, -COO-, -CH₂NH-, -OCH₂- or -NHCH₂-, especially -CONH-;







L_1 is a valence bond or an organic linker group containing 1 to 4 backbone atoms selected from C, N, O and
15 S;


Lp_1 is a cycloalkyl, azacycloalkyl, diazacycloalkyl, phenyl, naphthyl, adamantyl, decaliny, bicycloalkyl, mono- or diazabicycloalkyl, mono- or bicyclo heteroaromatic or a linear or branched alkyl, alkylene, alkenyl or alkenylene
20 group all optionally substituted by a group R_3 , or a combination of at least two such groups linked by a spiro linkage or a single or double bond or by C=O, O, S, SO, SO₂, CONR_{1e}, NR_{1e}-CO-, NR_{1e} linkage (for example, representative lipophilic groups include a methyl-cyclohexyl,
25 methylcyclohexylmethyl, bispiperidiny, methylphenylmethyl, phenylethyl, benzylpiperidiny, benzoylpiperidiny or phenylpiperaziny and those as hereinbefore described);

D is a hydrogen bond donor group;

and n is 0, 1 or 2.

30 Suitable R_2 groups may be





5


wherein R_5 is hydrogen, amino or hydroxy and R_3 (in relation to R_2) is halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} .

In a particularly favoured embodiment the R_2 group is an indole as marked by a * above in which R_5 is hydrogen and R_3 is a hydrogen or halogen present at the 3 position.

It is preferred that at least one of R_6 and R_7 be other than hydrogen and that R_6 , if present, is preferably a substituent containing one or more polar hydrogens such as hydroxy, amino, alkylamino, alkylaminoalkyl, aminocarbonyl, alkylaminocarbonyl, hydrazo and alkylhydrazo; alternatively R_6 and R_7 are joined together in the formation of a naphthyl or indolyl or azaindolyl or diazaindolyl group.

It is especially preferred that R_6 be amino and R_7 be chloro, bromo, methyl, methoxy or vinyl; or that R_6 and R_7 taken together form an indolyl ring with the NH at the 6-position or taken together form a naphthyl ring.

In a further preferred embodiment the compounds of the invention are of formula (A)



(A)

- 5 (wherein R_5 , R_6 , R_7 , Ar , $X-X$, Lp_1 , D_n are as hereinbefore defined; L_2 is a valence bond or an organic linker group containing 1 to 3 backbone atoms selected from C, N, O and S and R_{8a} and R_{9a} are hydrogen or taken together with the carbon atom to which they are attached form a carbonyl group). Again, in an alternative embodiment the phenyl derivative forming part of the R_2 functionality may instead be a nitrogen heterocyclic group, e.g. pyridine.


15 In one embodiment, L_2 comprises the backbone of an alpha amino acid, the lipophilic group being the side chain of the amino acid.

In one preferred embodiment R_{8a} and R_{9a} are hydrogen and L_2 is a $OC=O$ or $NHC=O$ group.

20 In a preferred embodiment, L_2 represents a valence bond and the lipophilic group is bound directly to a carbonyl alpha to the alpha atom via a nitrogen atom which forms part of the lipophilic group. Suitable lipophilic groups in this case therefore include piperidinyl, pyrrolidinyl and piperazinyl. In a preferred embodiment the piperidine or piperazinyl group is further substituted by a phenyl, benzyl, phenoxy, piperidine, pyridine or benzoyl group, optionally substituted on the phenyl ring by one or more R_3 groups. In a more preferred embodiment a piperazine is

substituted with a phenyl group substituted at the 2-position with an electron withdrawing group such as fluoro, nitro, triazolyl, cyano, alkoxycarbonyl, aminocarbonyl, aminosulphonyl, alkylaminosulphonyl and, especially preferred, alkylsulphonyl; and, at the 4-position, with hydrogen, fluoro, alkoxy or hydroxy. In another more preferred embodiment a piperidine is substituted at the 4-position with 4-piperidine which itself may be substituted on nitrogen by alkyl or aminocarbonylalkyl or alkylaminocarbonyl alkyl.

In a further embodiment, the lipophilic group has attached a group of the formula $-COOR_{1g}$ or $-CON$ -aminoacid or ester derivative thereof (where R_{1g} is as defined for R_{1a}). Particularly preferred compounds are those of formula (G)




(G)

(wherein Ar, R_6 and R_7 are as hereinbefore defined, R_5 represents hydrogen or amino and



represents a cyclic group) or of formula (H)




(H)

(wherein R_6 and R_7 are as hereinbefore defined, and R_5 represents hydrogen or amino). In a preferred embodiment R_6 is amino and R_7 a halogen, especially chlorine.

Again, in an alternative embodiment the phenyl derivative forming part of the R_2 functionality in formulae (G) and (H) may instead be a nitrogen heterocyclic group, e.g. pyridine, indole.

In another embodiment the group binding the alpha carbon atom to the lipophilic group comprises a heterocyclic group. Accordingly, preferred compounds of the invention also include those of formula (III)



(III)

(wherein R_5 , R_6 , R_7 , Ar , $X-X$, Lp_1 , D_n are as hereinbefore defined;

m is 0, 1 or 2;


Het is a 5 or 6-membered heterocyclic group interrupted by 1, 2 or 3 heteroatoms selected from O, N and S optionally substituted by a group R_{3b} where R_{3b} is as defined for R_3).

Again, in an alternative embodiment the phenyl derivative forming part of the R_2 functionality may instead be a nitrogen heterocyclic group, e.g. pyridine.


Where Het is a five membered ring, the two ring atoms at which it is connected are preferably separated by one ring atom. Where Het is a six-membered ring, the two ring atoms at which it is connected are preferably separated by one or two ring atoms. Representative heterocyclic groups include thiazole, oxazole, oxadiazole, triazole, thiadiazole or imidazole. Where the heterocyclic group is substituted by R_{3b} this is preferably a COOH or COOR_{1h} connected to the heterocycle via a valence bond or alkylene chain (where R_{1h} is as defined for R_{1a}).

In a further embodiment, the lipophilic group has attached a group of the formula $-\text{COOR}_{1g}$ or $-\text{CON-aminoacid}$ or ester derivative thereof.

In an alternative embodiment, the main aromatic R_2 ring in the compounds of the invention is a five membered aromatic ring leading to compounds of formula (IV) or (IVa)




(IV)




(IVa)


(wherein R_5 , R_6 , R_7 , $X-X$, Ar , L_1 , Lp_1 , D and n are as hereinbefore described for formula (II) and Z represents N , O or S). It is preferred that at least one of R_6 and R_7 be other than hydrogen, or that R_6 and R_7 taken together enable the formation of an indolyl, or azaindolyl group or diazaindolyl group. Preferences for other substituents are as for formula (A) above. Examples of possible fused systems are given below.

10




15







Hence in a preferred embodiment the compounds of the invention are of formula C or D




5



(wherein R_5 , R_6 , R_7 , Ar , $X-X$, Z , R_8 , R_9 , L_2 , Lp_1 , D_n are as
 10 hereinbefore defined) preferences for Ar , $X-X$, R_{8a} , R_{9a} , L_2 ,
 Lp_1 , D_n are as for formula (A) above; or compounds of
 formula E or F:



51




(F)

wherein Lp_1 is connected to the carbonyl via a nitrogen atom, R_6 , R_7 , Ar , Z , Lp_1 , D_n are as hereinbefore defined and R_5 is hydrogen or amino) preferences for Ar , Lp_1 , D_n are as
 5 for formula (A) above.


Particularly preferred are the compounds of formula I of Examples 35, 63, 66, 73, 100, 318 and 320, and physiologically tolerable salts thereof.

As previously mentioned, a number of compounds of the
 10 invention have been found to be excellent mixed inhibitors in that they inhibit both the serine proteases Factor Xa and thrombin. Such mixed inhibitors are preferably based on the formula (L)




(L)

15 wherein R' represents




X_3 represents hydrogen or a polar group such as amino or CONH_2 , especially CONH_2 ; and

- 5 R'' represents a cyclic group bound to the carbonyl by a nitrogen atom or an optionally substituted group of formula



- The compounds of the invention may be prepared by
 10 conventional chemical synthetic routes or by routes as
 illustrated by the following examples, e.g. by amide bond
 formation to couple the aromatic function to the alpha atom
 and to couple the lipophilic function to the alpha atom.
 Where the alpha atom is a carbon, the cyclic group-alpha
 15 atom combination may conveniently derive from an alpha amino


acid with the aromatic deriving from for example an acid derivative of a compound based on R_2 , e.g. o-amino-benzoic acid. Amide formation from such reagents (in which any amino or hydroxyl function may if desired be protected during some or all of the synthesis steps) yields a compound of formula (V).



(where Cy and R_2 are as defined above).

The lipophilic group (and optionally simultaneously the hydrogen bond donor) may then conveniently be introduced by reaction of a compound of formula (V) (or another analogous carboxylic acid) optionally after transformation into an activated form, e.g. an acid chloride or active ester, with a lipophilic group carrying an amine, hydroxylamine, hydrazine or hydroxyl group, e.g. to produce compounds with linkages of $\text{-CO-NR}_{1d}\text{-}$, $\text{-CO-NR}_{1d}\text{-O-}$, $\text{-CO-NR}_{1d}\text{-NR}_{1d}\text{-}$ and -CO-O- from the alpha atom (where it is a carbon) to the lipophilic group. Cyclisation can be base induced via nucleophilic attack of the alpha atom on a leaving group on the active side chain. If necessary the amide linkage can be reduced using an appropriate reducing agent employing the necessary protection depending on whether concurrent reduction of the carboxylic acid moiety is also desired. Alternatively a compound of formula V or another analogous carboxylic acid may be transformed into an alcohol by reaction with isobutylchloroformate and reduction with sodium borohydride.

Such an alcohol, e.g. of formula VI



can be reacted to introduce the lipophilic group by reactions such as:

alkylation with an alkyl halide in the presence of a
5 base;

under Mitsunobu conditions, such as reaction with diethyl azodicarboxylate/triphenylphosphine and a hydroxylated aryl compound;

10 by reaction with an activated carboxylic acid (e.g. an acid chloride) or with a carboxylic acid and diethylazodicarboxylate/triphenylphosphine;

by reaction with an isocyanate; and

15 by treatment with methanesulphonyl chloride or trifluoromethanesulphonic anhydride and reaction with an amine, or with a thiol optionally followed by oxidation, e.g. with potassium metaperiodate or hydrogen peroxide.

Alternatively, the reactions described above may be performed on a corresponding compound of formula (VI) in which R_2 is replaced with a protecting group, such as t-butoxycarbonyl (Boc), followed by deprotection and
20 introduction of the group R_2 .

In this way compounds with linkages of $-CH_2-O-$, $-CH_2-O-CO-$, $-CH_2-O-CO-NR_{1d}-$, $-CH_2-NR_{1d}-$, $-CH_2-S-$, $-CH_2-SO-$ and $-CH_2-SO_2-$ between the alpha carbon and the lipophilic
25 group may be produced.

Alternatively the alcohol can be oxidized to form a corresponding aldehyde (e.g. by oxidation with manganese dioxide or DMSO/oxalyl chloride or DMSO/ SO_3 or Dess-Martin reagent) which may be reacted to introduce the lipophilic
30 group by reactions such as:

reaction with Wittig reagents or Horner-Emmons reagents, optionally followed by reduction of the resulting carbon:carbon double bond using H_2/Pd -carbon;

reaction with an organometallic, eg a Grignard reagent,
 5 optionally followed by reaction on the resulting hydroxyl group, such as oxidation (eg with MnO_2 , DMSO/oxalyl chloride or Dess-Martin reagent), alkylation (eg with an alkyl halide in the presence of a base in a solvent such as DMF), arylation (eg with diethylazo dicarboxylate/triphenyl
 10 phosphine and a hydroxyaryl compound), ester formation (eg with an acid chloride or with a carboxylic acid and diethylazido dicarboxylate/triphenyl phosphine), or carbamate formation (eg with an isocyanate);

by reaction with an amine followed by reduction, e.g.
 15 with sodium cyanoborohydride;
 by reaction with a hydrazine; or
 by reaction with a carbazide.

In this way compounds with linkages of $-CH=CR_{1d}-$,
 $-CH_2-CHR_{1d}-$, $-CHOH-$, $-CHR_{1d}-O-$, $-CHR_{1d}-O-CO-$,
 20 $-CHR_{1d}-O-CO-NR_{1d}-$, $-CO-$, $-CH_2-NR_{1d}-$, $-CH=N-NR_{1d}-$ and
 $-CH=N-NR_{1d}-CO-NR_{1d}-$ between the alpha carbon and the lipophilic group may be produced.

The transformation of alcohol to amine referred to above may be used to produce an amine reagent for lipophilic
 25 group introduction, e.g. a compound




Such an amine reagent may be reacted to introduce the lipophilic group, e.g. by acylation with an acid halide or activated ester, by reaction with isocyanate, by reaction
 30 with an isothiocyanate, or by reaction with a sulphonyl chloride. In this way compounds with linkages of $-CH_2NR_{1d}-CO-$, $-CH_2-NR_{1d}-CO-NR_{1d}-$, $-CH_2NR_{1d}-CS-NR_{1d}-$ and $-CH_2NR_{1d}-SO_2-$

between the alpha carbon and the lipophilic groups may be produced.

The transformation of acid to amide referred to above may be used to produce an amide reagent for introduction of the lipophilic group, e.g. a compound




Such amides may be reacted to introduce lipophilic groups, e.g. by reaction with a haloketone (e.g. phenacyl bromide). This provides a linkage




from alpha carbon to lipophilic group.


Analogously the amide may be transformed to a thioamide by reaction with Lawesson's reagent and then reacted with a haloketone to form a linkage




The amide reagent may likewise be transformed to a nitrile reagent by dehydration, e.g. with trifluoroacetic anhydride. The nitrile reagent may be reacted with hydrazine then with acyl halide and then cyclized, (e.g. with trifluoroacetic anhydride) to produce a linkage




Alternatively it may be treated with hydroxylamine then reacted with acyl halide and cyclized (e.g. with trifluoroacetic anhydride) to produce a linkage



The hydrazide produced by reaction of a carboxylic acid reagent with hydrazine discussed above may likewise be used as a reagent for lipophilic group introduction, e.g. as a
 5 compound of formula




Thus the hydrazide reagent can be reacted with an acyl halide and cyclized, e.g. with trifluoroacetic anhydride to yield a linkage




10

or reacted with an acyl halide or an isocyanate to yield linkages $\text{-CO-NR}_{1d}\text{-NR}_{1d}\text{-CO-}$ and $\text{-CO-NR}_{1d}\text{-NR}_{1d}\text{-CO-NR}_{1d}\text{-}$ respectively.

Alternatively the hydrazide may be transformed by
 15 reaction with Lawesson's reagent and then reacted with an acyl halide and cyclized (e.g. with trifluoroacetic anhydride) to produce the linkage



An alternative route to these compounds is to carry out
 20 any of the above chemical reactions to incorporate the lipophilic group (and optional H bond donor) into a protected intermediate such as a compound of formula (VII).



PG = Protecting group

The protecting group may then be removed before coupling of the for example o-amino benzoic acid (optionally protected).

The protection of amino and carboxylic acid groups is described in McOmie, Protecting Groups in Organic Chemistry, Plenum Press, NY, 1973, and Greene and Wuts, Protecting Groups in Organic Synthesis, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of carboxy protecting groups include C₁-C₆ alkyl groups such as methyl, ethyl, t-butyl and t-amyl; aryl(C₁-C₄)alkyl groups such as benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, benzhydryl and trityl; silyl groups such as trimethylsilyl and t-butyldimethylsilyl; and allyl groups such as allyl and 1-(trimethylsilylmethyl)prop-1-en-3-yl.

Examples of amine protecting groups (PG) include acyl groups, such as groups of formula RCO in which R represents C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, phenyl C₁₋₆ alkyl, phenyl, C₁₋₆ alkoxy, phenyl C₁₋₆ alkoxy, or a C₃₋₁₀ cycloalkoxy, wherein a phenyl group may be optionally substituted, for example by one or two of halogen, C₁-C₄ alkyl and C₁-C₄ alkoxy. Preferred amino protecting groups include benzyloxycarbonyl (CBz), t-butoxycarbonyl (Boc) and benzyl.

Compounds of the type (VII) made be prepared (for example) by one or more of the following methods.

(i) from aryl or heteroaryl aldehydes via the Strecker synthesis or modifications thereof, via Bucherer-Bergs

hydantoin synthesis, or via the Ugi methodology (Isonitrile Chemistry, Ugi I. Ed.; Academic: New York, 1971; pp145-199) with removal and replacement of protecting groups;

(ii) from styrenes via Sharpless methodology (J. Am. Chem. Soc. 1998, 120, 1207-1217)

(iii) from aryl boronic acids via Petasis methodology (Tetrahedron, 1997, 53, 16463-16470) with removal and replacement of protecting groups;


(iv) from aryl and heteroaryl acetic acids - via Evan's azidation (Synthesis, 1997, 536-540) or by oximation, followed by reduction and addition of protecting groups; or

(v) from existing aryl glycines by manipulation of functional groups, for example, alkylation of hydroxy groups, palladium assisted carbonylation of triflates derived from hydroxy groups and further manipulation of the carboxylic esters to give carboxylic acids by hydrolysis, carboxamides by activation of the carboxylic acid and coupling with amines, amines via Curtius reaction on the carboxylic acid or

(vi) from aliphatic, carbocyclic and non-aromatic heterocyclic aldehydes and ketones using a Horner-Emmons reaction with N-benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester (Synthesis, 1992, 487-490).

A starting reagent for lipophilic group introduction where the alpha atom is nitrogen may be produced for example by reaction of a beta protected hydrazine (such protection to be chosen as to be compatible with the subsequent reagents to be employed) with phosgene, diphosgene, triphosgene or N,N'carbonyl

diimidazole to give a reactive compound of the type:



PG = Protecting group


This intermediate may be used as has been described above for the carboxylic starting reagents where the alpha atom is carbon.

Removal of the protecting group by standard methods and coupling with an activated aryl carboxylic acid will give compounds of the type



(where R_2 , X, Y, Cy, L, Lp and D are as defined above).

Thus viewed from a further aspect the invention provides a process for the preparation of a compound according to the invention which process comprises coupling a lipophilic group to a compound of formula (VIII)




(wherein R_2 , X, Y and Cy are as defined above and Z_1 is a reactive functional group), and optionally subsequently coupling a hydrogen bond donor group to said lipophilic group.

Instead of introducing the group $\text{L}-\text{Lp}(\text{D})_n$ as the final stage process step, the compounds of formula I may alternatively be prepared by a process in which the group R_2 is introduced in the final process step.


Thus viewed from another aspect the invention provides a process for the preparation of a compound according to the invention which process comprises coupling a lipophilic group to a compound of formula (IX)

5



(wherein Y, Cy, L, Lp D, and n are as defined above and Z_2 is HX or a reactive functional group), or a protected derivative thereof, with a compound of formula (X)

10



(wherein R_2 is as defined above and Z_3 is XH or an appropriate reactive group), or a protected derivative thereof, followed if necessary by the removal of any protecting groups.

15


Thus, for a compound of formula I in which X-X represents CONH, a compound of formula (IX) in which Z_2 is H_2N may be reacted with a compounds of formula (X) in which Z_3 is COOH or a reactive derivative thereof, such as a acyl halide or an anhydride, for example as described in the Examples herein.

20


Where the lipophilic group Lp comprises more than one group, it may generally be formed by coupling these groups together at an appropriate stage in the preparation of the compound of formula I using conventional methods or as described in the Examples.

25

For a compound of formula I in which Lp comprises an azacycloalkyl or diazacycloalkyl group of formula



in which X_b is N and each of s and u is 0, alkylating the amino group of a corresponding compound in which the corresponding residue is of formula




5

using a conventional alkylating method. The alkylation may be carried out using any conventional method; however, generally preferred is a reductive alkylation using the appropriate aldehyde or ketone, for example as described in the Alkylation Methods in the Examples.

10

Thus, a particular starting material for the alkylation is one of formula




in which X_a is N and L is CO or X_a is CH and L is CONH,

15


CONHCH₂ or CH₂NHCO.

For a compound of formula I in which L_p comprises an azacycloalkyl or diazacycloalkyl group of formula



20

in which R_{10} is a group of formula




in which X_d is N and R_{11} is (1-6C)alkyl, alkylating the amino group of a corresponding compound of formula I in which R_{11} is hydrogen using a conventional method.

- 5 Generally preferred is a reductive alkylation using the appropriate aldehyde or ketone, for example as described in the Alkylation Methods in the Examples.


For a compound of formula I in which Lp comprises an azacycloalkyl or diazacycloalkyl group of formula

10



in which X_b is CH and $(L_a)_s-(G)_t-(L_b)_u$ is O and R_{10} is phenyl or pyridyl, coupling a corresponding compound containing a


15 group of formula




with phenols or 3-hydroxypyridine using Mitsunobu conditions, eg. DEAD (diethyl azodicarboxylate) / Ph_3P or 2-triphenylphosphonium 4,4-dimethyl-tetrahydro-1,2,5-thiadiazole to give aryloxy or 3-pyridoxy substituted

20 piperidines or pyrrolidine. Alternatively the hydroxy group may be reacted with sodium hydride and 2-chloro or 4-chloropyridine to give 2-pyridoxy or 4-pyridoxy substituted piperidines or pyrrolidines.


For a compound of formula I in which $-L-Lp(D)_n$ is




in which Q is a direct bond, reductively alkylating an amine
 5 of formula $H-Q$ using a corresponding compound in which the
 corresponding residue is a ketone of formula



For a compound of formula I in which $-L-Lp(D)_n$ is



10 in which Q is methylene, reductively alkylating an amine of
 formula $H-NR_aR_b$ using a corresponding compound in which the
 corresponding residue is an aldehyde of formula



The intermediates used in the process according to the
 15 invention may generally, when not commercially available, be
 prepared by conventional methods or as described in the
 Examples herein.

For example, methyl 1-acetyl-3-formylindole-6-
 carboxylic acid may be converted to the 3-formate by the

method of Merour et al (Synthesis, 1994, 411) and then reacted with trimethyl orthoformate to give methyl 1-acetyl-3-methoxyindole-6-carboxylate which is then hydrolysed to methyl 1-acetyl-3-methoxyindole-6-carboxylate.

- 5 5-Fluoroindole-6-carboxylic acid may be prepared from 4-fluoro-3-methoxyaniline by the following method. 4-Fluoro-3-methoxyaniline is treated with glyoxal-1,1-dimethyl acetal and then hydrogenated over Pd/C. The product is N-protected with methanesulphonyl chloride and then cyclised using
- 10 titanium tetrachloride in toluene. Demethylation with BBr₃ to the phenol followed by reaction with triflic anhydride and then palladium carbonylation in methanol gives the methyl ester, which is then converted to 5-fluoro-1-methanesulphonylindole-6-carboxylic acid by hydrolysis with
- 15 lithium hydroxide. This 'benzoyl' component may be reacted as previously described and deprotected by hydrolysis with sodium hydroxide at 100°C.

The intermediates disclosed herein, including the novel intermediates of formulae (V), (VI), (VII), (VIII) and (IX)

20 are provided as further aspects of the invention.

The compounds of the invention may be administered by any convenient route, e.g. into the gastrointestinal tract (e.g. rectally or orally), the nose, lungs, musculature or vasculature or transdermally. The compounds may be

25 administered in any convenient administrative form, e.g. tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches etc. Such compositions may contain components conventional in pharmaceutical preparations, e.g. diluents,

30 carriers, pH modifiers, sweeteners, bulking agents, and further active agents. Preferably the compositions will be

sterile and in a solution or suspension form suitable for injection or infusion. Such compositions form a further aspect of the invention.

The following are examples of pharmaceutical
5 compositions of compounds according to the invention.

Formulation 1

Hard gelatin capsules are prepared using the following
10 ingredients:

		Quantity (mg/capsule)
Active Ingredient		250
Starch, dried		200
Magnesium stearate		<u>10</u>
Total		460 mg

The above ingredients are mixed and filled into hard
25 gelatin capsules in 460 mg quantities.

Formulation 2

Tablets each containing 60 mg of active ingredient are made as follows:

5		
	Active Ingredient	60 mg
	Starch	45 mg
	Microcrystalline cellulose	35 mg
10	Polyvinylpyrrolidone	4 mg
	Sodium carboxymethyl starch	4.5 mg
	Magnesium stearate	0.5 mg
	Talc	<u>1 mg</u>
15	Total	150 mg

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

It is believed that the compounds of the invention will have excellent oral bioavailability.

30 Viewed from this aspect the invention provides a pharmaceutical composition comprising a serine protease inhibitor according to the invention together with at least

one pharmaceutically acceptable carrier or excipient. The pharmaceutical composition may also optionally comprise at least one further antithrombotic and/or thrombolytic agent.

Viewed from a further aspect the invention provides the use of a serine protease inhibitor according to the invention for the manufacture of a medicament for use in a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat (i.e. treat or prevent) a condition responsive to said inhibitor.

Viewed from a further aspect the invention provides a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat a condition responsive to a serine protease inhibitor (e.g. a condition such as a thrombotic disorder responsive to a factor Xa inhibitor), said method comprising administering to said body an effective amount of a serine protease inhibitor according to the invention.

The dosage of the inhibitor compound of the invention will depend upon the nature and severity of the condition being treated, the administration route and the size and species of the patient. However in general, quantities of from 0.01 to 100 $\mu\text{mol/kg}$ bodyweight will be administered.

All publications referred to herein are hereby incorporated by reference.

The invention will now be described further with reference to the following non-limiting Examples.

Experimental

Abbreviations used follow IUPAC-IUB nomenclature.

Additional abbreviations are Hplc, high-performance liquid
5 chromatography; DMF, dimethylformamide; DCM,
dichloromethane; HAOT, 1-hydroxy-7-azabenzotriazole; HATU,
[O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium
hexafluorophosphate]; Fmoc, 9-Fluorenylmethoxycarbonyl;
HOBT, 1-hydroxybenzotriazole; TBTU, 2-(1H-(benzotriazol-1-
10 yl)-1,1,3,3-tetramethyluroniumtetrafluoroborate; EDCI, 1-(3-
Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride;
DIPEA, diisopropylethylamine; Boc, tertiary
butyloxycarbonyl; DIPCI, diisopropylcarbodiimide; DBU, 1,8-
diazabicyclo[5.4.0]undec-7-ene; TEA, triethylamine; Rink
15 linker, p-[(R,S)- α -[1-(9H-Fluoren-9-yl)methoxyformamido]-
2,4-dimethoxybenzyl]phenyl acetic acid; TFA, trifluoroacetic
acid; MALDI-TOF, Matrix assisted laser desorption ionisation
- time of flight mass spectrometry, RT, retention time.
Amino acid derivatives, resins and coupling reagents were
20 obtained, for example, from Novabiochem (Nottingham, UK) and
other solvents and reagents from Rathburn (Walkerburn, UK)
or Aldrich (Gillingham, UK) and were used without further
purification. All solution concentrations are expressed as
%Vol./%Vol. unless otherwise stated.

25

Purification: Purification was by gradient reverse phase
Hplc on a Waters Deltaprep 4000 at a flow rate of 50 ml/
min. using a Deltapak C18 radial compression column (40 mm x
210 mm, 10-15 mm particle size). Eluant A consisted of
30 aqTFA (0.1%) and eluant B 90% MeCN in aq TFA(0.1%) with
gradient elution (Gradient 1, 0 min. 20%B then 20% to 100%
over 36 min., Gradient 2, 0 min. 5%B for 1 min. then 5%B to

20%B over 4 min., then 20% to 60% over 32 min. or Gradient 3, 0 min. 20%B then 20% to 100% over 15 min.). Fractions were analysed by analytical Hplc and MALDI-TOF before pooling those with >95% purity for lyophilisation.

5

Analysis: Analytical Hplc was on a Shimadzu LC6 gradient system equipped with an autosampler, a variable wavelength detector at flow rates of 0.4 ml/ min. Eluents A and B as for preparative Hplc. Columns used were Techogell15 C18 (2x150mm) (Hplc Technology), Magellan C8 column (2.1x150 mm, 5µm particle size) and Luna C18 (2.1x150 mm, 5µM particle size). (Phenomenex)) Purified products were further analysed by MALDI-TOF and nmr. NMR denotes an ¹HNMR consistent with the structure was obtained.

15

Synthesis of inhibitors

Method 1: Using a solid phase strategy on a Protein Technologies, Symphony Multiple Peptide Synthesiser by attachment of bis amino compounds to Peg-trityl chloride resin: Trityl chloride resin was typically treated with greater than 2 fold excess of the di-amine in dry DCM .The resin was further modified by the attachment of acids. Activation of Fmoc protected amino acid (2-5eq) was by TBTU/ DIPEA, all couplings (minimum 120 min.) were carried out in DMF. Deprotection of the Fmoc group was achieved with 20% piperidine in DMF. In the next stage other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU or HATU/EDCI with or without Boc protection of amino groups. Cleavage of the products from the resin was by treatment (30 min., ambient) with 10% triethylsilane in

30

TFA, filtration, evaporation and trituration with diethylether.

Synthesis using the Symphony Multiple Peptide Synthesiser.

5

The Symphony Multiple Peptide Synthesiser is charged with DMF, DCM, TBTU in DMF (450 mM), DIPEA in DMF (900 mM), 20% piperidine in DMF. Resins are held in plastic reaction vessels that allow the introduction of reagents and solvents and nitrogen for agitation or air drying.

10

A typical synthesis cycle on the Symphony is as follows:-

The reaction vessel containing the resin (0.1 mmol) is charged with the Fmoc protected amino acid (0.5 mmol) and then this is dissolved in DMF (2.5ml), treated with TBTU (0.56 mmol, 1.25ml) and DIPEA (1.1 mmol, 1.25ml) and agitated with nitrogen for 2 hours (agitation times may vary). After coupling the resin is washed with DMF (6x 5ml) then deprotected with 20% piperidine in DMF (2x 5ml for 1 min.each, then 1x 5ml for 8 min.) the resin is then washed with DMF (6x 5ml).

20

Example 1.

25 1-(2-Amino-4-chlorobenzoyl-D-phenylglyciny1)-4,4'-bispiperidine

4,4-Bipiperidine.dihydrochloride (4mmol,1g) was dissolved in water (5ml) and 2M sodium hydroxide solution (10mmol, 5ml) added. The solution was extracted with ethylacetate (2x 50ml) the combined extracts were washed with water, dried over anhydrous sodium carbonate, filtered and evaporated to give the 4,4 bipiperidine (0.35g) as a white solid. The 4,4

30

bipiperidine was dissolved in dry DMF (2ml) and added to
Peg-tritylchloride resin (0.95 mmol/g, 1.5g) pre swollen in
dry DCM (10ml). After 2h the resin was washed with DCM
(6x5ml), DMF (6x5ml) and DCM (6x5ml). The resin was then air
5 dried to allow aliquots to be taken.

The 4,4 bipiperidine trityl resin (0.1 mmol) was treated
with Fmoc-D-Phenylglycine (0.5 mmol, 187mg), DMF (2.5ml),
TBTU in DMF (1.25ml of a 450mM solution) and DIPEA in DMF
10 (1.25ml of a 900 mM solution). The mixture was agitated with
nitrogen for 2 hours. Deprotection and washing as above.

A solution of 4-chloroanthranilic acid (87mg 0.5mmole) in
dry dimethylformamide (DMF) was treated successively with
15 HOAt (102mg 0.75mmole) and EDCI (115mg 0.6mmole) and stirred
at room temperature for 10min. The mixture was transferred
to the reaction vessel on the Symphony and agitated for 2
hours with nitrogen. The resin was washed with DMF (6x5ml),
DCM (6x5ml) and air dried. The product was cleaved from the
20 resin with 10% triethylsilane in TFA (10ml) for 30 minutes,
the resin filtered off and the TFA solution evaporated to
dryness and triturated with diethyl ether to give the crude
product. The crude product was dissolved in water (10ml),
filtered and purified by preparative reverse phase Hplc.

25

^1H nmr (CD_3CN) 7.30 (6H,m); 6.60 (1H,s); 6.55 (1H,d); 5.85
(1H, s); 4.40 (1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 1.60
(4H, m); 1.10 (6H, m) MS TOF 456 ($\text{M}+1^+$). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 11.77 min.

30

Example 2.

1-(2-Amino-5-bromobenzoyl-D-phenylglyciny1)-4,4'-
bispiperidine

1H nmr (CD₃CN) 7.30 (7H,m); 6.50 (1H,d); 5.85 (1H, s); 4.40
5 (1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 1.60 (4H, m); 1.10
(6H, m) MS TOF 500 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 11.31 min.

Example 3.

10 1-(2-Amino-4-methylbenzoyl-D-phenylglyciny1)-4,4'-
bispiperidine

1H nmr (CD₃CN) 7.30 (6H,m); 6.50 (1H,s); 6.45 (1H,d); 5.80
(1H, s); 4.40 (1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 2.05
(3H,s); 1.60 (4H, m); 1.10 (6H, m) MS TOF 436 (M+1⁺). Hplc
15 (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 9.22
min.

Example 4.

20 1-(2-Amino-5-methylbenzoyl-D-phenylglyciny1)-4,4'-
bispiperidine

1H nmr (CD₃CN) 7.30 (7H,m); 6.50 (1H,d); 5.85 (1H, s); 4.40
(1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 1.60 (4H, m); 1.10
(6H, m). MS TOF 436 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 8.74 min.

25

Example 5.

1-(2-Amino-5-methoxybenzoyl-D-phenylglyciny1)-4,4'-
bispiperidine

1H nmr (CD₃CN) 7.55 (6H,m); 7.30 (1H,d); 6.95 (1H,m); 6.15
30 (1H, s); 4.40 (1H,m); 3.75 (1H, m); 3.60 (3H, s); 2.30-2.95
(6H, m); 2.20 (3H, s); 1.60 (4H, m); 1.10 (6H, m) MS TOF 452

(M+1⁺).. Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 8.20 min.

Example 6.

5 1-(3-Methylbenzoyl-D-phenylglyciny1)-4,4'-bispiperidine

¹H nmr (CD₃CN) 7.40 (2H,m); 7.30 (7H,m); 5.85 (1H, s); 4.40
(1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 2.20 (3H, s); 1.60
(4H, m); 1.10 (6H, m) MS TOF 421 (M+1⁺). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 10.68 min.

10

Example 7.

1-(4-Methylbenzoyl-D-phenylglyciny1)-4,4'-bispiperidine

¹H nmr (CD₃CN) 7.55 (2H,m); 7.30 (5H,m); 7.10 (2H,m); 5.85
(1H, s); 4.40 (1H,m); 3.75 (1H, m); 2.30-2.95 (6H, m); 2.20
15 (3H,s); 1.60 (4H, m); 1.10 (6H, m) MS TOF 420 (M+1⁺). Hplc
(Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 10.61
min.

Example 8.

20 1-(3-Amino-2-naphthoyl-D-phenylglyciny1)-4,4'-bispiperidine

¹H nmr (CD₃CN) 7.90 (1H,d); 7.60 (1H,d); 7.40 (1H,m); 7.30
(6H,m); 7.05 (1H,m); 6.90 (1H,s); 5.85 (1H, s); 4.40 (1H,m);
3.75 (1H, m); 2.30-2.95 (6H, m); 1.60 (4H, m); 1.10 (6H, m)
MS TOF 471 (M+1⁺). Hplc (Magellan C8, Gradient 3,
25 water/acetonitrile/TFA) rt 9.87 min.

Example 9.

1-(3-Aminobenzoyl-D-phenylglyciny1)-4,4'-bispiperidine

MS TOF 421 (M+1⁺). Hplc (Magellan C8, Gradient 3,
30 water/acetonitrile/TFA) rt 9.06 min.

Example 10.

1-(2-Aminobenzoyl-D-phenylglyciny)-4,4'-bispiperidine
MS TOF 421 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 9.00 min.

5

Example 11.

1-(2-Amino-4-fluorobenzoyl-D-phenylglyciny)-4,4'-
bispiperidine

MS TOF 440 (M+1⁺). Hplc (Magellan C8, Gradient 3,
10 water/acetonitrile/TFA) rt 9.23 min.

Example 12.

1-(2-Amino-5-fluorobenzoyl-D-phenylglyciny)-4,4'-
bispiperidine

15 MS TOF 440 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 9.14 min.

Example 13.

1-(2-Amino-4-nitrobenzoyl-D-phenylglyciny)-4,4'-
20 bispiperidine

MS TOF 467 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 10.59 min.

Example 14.

25 1-(2-Amino-5-nitrobenzoyl-D-phenylglyciny)-4,4'-
bispiperidine

MS TOF (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 10.57 min.

30 Example 15.

1-(2-Amino-4,5-dimethoxybenzoyl-D-phenylglyciny)-4,4'-
bispiperidine

MS TOF 481 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.67 min.

Example 16.

5 1-(Benzoyl-D-phenylglyciny)-4,4'-bispiperidine

MS TOF 407 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 9.88 min.

Example 17.

10 1-(4-Chlorobenzoyl-D-phenylglyciny)-4,4'-bispiperidine

MS TOF 441 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 10.89 min.

Example 18.

15 1-(2-Hydroxybenzoyl-D-phenylglyciny)-4,4'-bispiperidine

MS TOF 423 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 8.97 min.

Method 2: By solution phase strategy: Typically an activated
20 amino acid was treated with an amine (primary or secondary) or alcohol (1eq.). Activation of the protected amino acid (Boc or Cbz protection) was by HATU/DIPEA (1:2) by TBTU/DIPEA (1:2), by HOBt or HOAt and a carbodiimide (EDCI or DCC), or by diethyl cyanophosphonate and triethylamine or
25 DIPEA, all couplings (minimum 120min.) were carried out in DMF without or without dichloromethane as co-solvent. After an aqueous work up the deprotection of the Boc group was achieved with TFA. Other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU, EDC
30 or DIPCI with or without Boc protection of amino groups. The final products were purified by preparative reverse phase Hplc.

Examples 19-126

The compounds of Examples 19-126 were prepared by the method described below, but using the appropriate starting materials.

Boc D-phenylglycine (251 mg, 1 mmol.) was dissolved in DMF (3ml) with HATU (380 mg., 1 mmol.) and DIPEA (350 μ l., 2 mmol.). To this mixture was added 4-methylbenzylamine (121mg., 1 mmol.) and DIPEA (170 μ l., 1 mmol.). The mixture was stirred overnight. The mixture was then taken up into ethylacetate and washed with water, sodium carbonate solution, water, 10% hydrochloric acid solution and water. The ethylacetate was evaporated without drying and treated immediately with TFA for 30 min. The TFA was then evaporated to dryness and the product triturated with diethylether. TEA (1ml) was added and evaporated to dryness. A solution of 3-hydroxymethylbenzoic acid (76mg, 0.5mmole) in dry dimethylformamide (DMF) was treated with TBTU (161mg., 0.5mmol.) and DIPEA (1.5 mmol.). The mixture was then added to the D-phenylglycine-4-methylbenzylamide (0.5mmol.) and stirred overnight. The crude product was dissolved in water/acetonitrile (20ml), filtered and purified by preparative Hplc to yield pure product.

^1H nmr (CD_3CN) 7.75 (1H, m); 7.65 (2H, m); 7.30 (7H, broad m); 6.80 (3H, m); 5.40 (1H, s); 4.45 (2H, s); 4.10 (2H, m); 2.10 (3H, s). MS TOF 389 ($\text{M}+1^+$). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.51 min.

Compounds made by the above method:-

Example 19.

1-(2-Aminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (DMSO) 7.65 (3H, m); 7.45 (1H, m); 7.35 (5H, m); 7.15 (1H, m); 6.65 (1H, d); 6.55 (1H, m); 6.05 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 511 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.43 min.

Example 20.

10 1-(2-Amino-4-chlorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (DMSO) 7.55 (3H, m); 7.45 (1H, m); 7.35 (5H, m); 7.15 (1H, m); 6.75 (1H, s); 6.55 (1H, d); 6.05 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 546 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.18 min.

Example 21.

1-(2-Amino-5-fluorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

20 1H nmr (CDCl₃) 7.75 (1H, m); 7.60 (1H, m); 7.25 (6H, m); 7.15 (1H, m); 6.90 (1H, m); 6.75 (1H, m); 5.85 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 529 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.87 min.

25 Example 22.

1-(2-Amino-4-methylbenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

30 1H nmr (DMSO) 7.55 (3H, m); 7.45 (2H, m); 7.35 (5H, m); 6.65 (1H, s); 6.35 (1H, d); 6.05 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m) 2.15 (3H, s);. MS TOF 525 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.12 min.

Example 23.

1-(2-Amino-5-methylbenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 7.75 (1H, m); 7.60 (1H, m); 7.25 (6H, m); 7.15 (1H, m); 6.90 (1H, m); 6.75 (1H, m); 5.85 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m) 2.30 (3H, s). MS TOF 525 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.84 min.

10 Example 24.

1-(2-Amino-4-nitrobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 7.75 (2H, m); 7.55 (1H, m); 7.35 (7H, m); 7.25 (1H, m); 5.80 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 556 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.35 min.

Example 25.

20 1-(2-Amino-5-nitrobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 8.25 (1H, d); 7.85 (1H, m); 7.55 (1H, m); 7.25 (7H, m); 7.05 (1H, m); 5.80 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 556 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.08 min.

25

Example 26.

1-(2-Amino-5-cyanobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.65 (4H, m); 7.25 (6H, m); 6.65 (1H, d); 5.80 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 536 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.89 min.

Example 27.

1-(2,5-Diaminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 5 ^1H nmr (CDCl_3) 7.70 (1H, d); 7.45 (7H, m); 6.85 (1H, s); 6.55 (1H, m); 6.55 (1H, m); 5.90 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 526 ($\text{M}+1^+$). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.82 min.

10 Example 28.

1-(2-Amino-4,5-dimethoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- ^1H nmr (CD_3CN) 7.65 (2H, m); 7.35 (2H, m); 7.25 (5H, m); 6.75 (1H, d); 6.15 (1H, d); 5.80 (1H, s); 3.60 (3H, s); 3.50 (3H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 571 ($\text{M}+1^+$). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.84 min.

Example 29.

- 20 1-(Benzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- ^1H nmr (CD_3CN) 7.75 (2H, m); 7.70 (1H, m); 7.40 (10H, m); 6.05 (1H, s); 3.15 (3H, s); 3.00-2.00 (8H, m). MS TOF 496 ($\text{M}+1^+$). Hplc (Magellan C8, Gradient 3, 25 water/acetonitrile/TFA) rt 12.84 min.

Example 30.

1-(3-Aminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 30 ^1H nmr (CD_3CN) 7.85 (1H, m); 7.60 (1H, m); 7.50 (2H, m); 7.30 (7H, m); 7.05 (1H, d); 6.05 (1H, s); 3.15 (3H, s); 3.00-

2.00 (8H,m). MS TOF 511 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.32 min.

Example 31.

5 1-(4-Aminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
¹H nmr (CDCl₃) 7.95 (1H, d); 7.80-7.45 (10H, broad m); 7.35 (1H,d); 6.20 (1H, s); 3.15 (3H,s); 3.00-2.00 (8H,m). MS TOF 511 (M+1⁺). Hplc (Magellan C8, Gradient 3,
10 water/acetonitrile/TFA) rt 12.05 min.

Example 32.

1-(3,4 Diaminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
15 ¹H nmr (CDCl₃) 7.75 (1H, d); 7.40-7.15 (9H, broad m); 6.55 (1H,d); 6.00 (1H, s); 3.15 (3H,s); 3.00-2.00 (8H,m). MS TOF 540 (M+1⁺). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 11.30 min.

20 Example 33.

1-(3-Chlorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
¹H nmr (CD₃CN) 7.85 (1H, m); 7.80 (1H, s); 7.60 (2H, m); 7.30 (8H, m); 6.00 (1H, s); 3.20 (3H,s); 3.00-2.00 (8H,m).
25 MS TOF 531 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.40 min.

Example 34.

1-(4-Chlorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylph nyl)piperazine
30 ¹H nmr (CD₃CN) 7.95 (1H, m); 7.75 (2H, m); 7.60 (1H, m); 7.40 (8H, m); 6.05 (1H, s); 3.25 (3H,s); 3.00-2.00 (8H,m).

MS TOF 531 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 16.54 min.

Example 35.

5 1-(3-Amino-4-chlorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 8.05 (1H, m); 7.80 (1H, m); 7.70 (1H, s); 7.20-7.60 (8H, broad m); 6.05 (1H, s); 3.25 (3H, s); 3.00-2.00 (8H, m). MS TOF 546 (M+1⁺). Hplc (Magellan C8, Gradient
10 3, water/acetonitrile/TFA) rt 14.53 min.

Example 36.

1-(4-Bromobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

15 ¹H nmr (CD₃CN) 7.85 (1H, m); 7.65 (2H, m); 7.60 (2H, d); 7.45 (2H, d); 7.30 (5H, m); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.00 (8H, m). MS TOF 576 (M+1⁺). Hplc (Magellan C8, Gradient
3, water/acetonitrile/TFA) rt 15.94 min.

20 Example 37.

1-(4-Iodobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN)) 7.75 (2H, m); 7.65 (1H, m); 7.55 (2H, d); 7.45 (2H, d); 7.30 (5H, m); 5.95 (1H, s); 3.20 (3H, s); 3.00-
25 2.00 (8H, m). MS TOF 622 (M+1⁺). Hplc (Magellan C8, Gradient
3, water/acetonitrile/TFA) rt 15.96 min.

Example 38.

1-(3-Amino-4-methylbenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

30 ¹H nmr (CDCl₃) 7.95 (1H, s); 7.60 (1H, d); 7.45 (1H, d); 7.40-7.15 (8H, broad m); 6.00 (1H, s); 3.15 (3H, s); 3.00-

2.50 (8H,m) 2.20 (3H, s). MS TOF 525 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.71 min.

Example 39.

5 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
1H nmr (CD₃CN) 7.85 (2H, d); 7.65 (1H, m); 7.50 (2H, m);
7.40 (5H, m); 6.80 (2H, d); 6.00 (1H, s); 3.80 (3H, s); 3.20
(3H,s); 3.00-2.00 (8H,m). MS TOF 526 (M+1⁺). Hplc (Magellan
10 C8, Gradient 3, water/acetonitrile/TFA) rt 14.63 min.

Example 40.

1-(3-Amino-4-methoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
15 1H nmr (CDCl₃) 7.90 (1H, m); 7.75 (1H, d); 7.60 (2H, m);
7.40-7.15 (6H, broad m); 7.45 (1H, d); 6.10 (1H, s); 3.95
(3H, s); 3.35 (3H,s); 3.00-2.50 (8H,m). MS TOF 541 (M+1⁺).
Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
11.78 min.

20

Example 41.

1-(3,4-Dihydroxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
1H nmr (CDCl₃) 7.55 (1H, m); 7.45 (1H, d); 7.25 (2H, m);
25 7.15 (5H, m); 7.00 (1H, d); 6.60 (1H, d); 5.80 (1H, s); 3.05
(3H,s); 3.00-2.50 (8H,m). MS TOF 541 (M+1⁺). Hplc (Magellan
C8, Gradient 3, water/acetonitrile/TFA) rt 11.78 min.

Example 42.

30 1-(Naphth-2-oyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 8.35 (1H, s); 8.00 (1H, d); 7.85 (5H, m); 7.45 (4H, m); 7.25 (4H, m); 6.10 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 546 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 16.66 min.

5

Example 43.

1-(3-Aminonaphth-2-oyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 8.15 (1H, d); 8.00 (1H, s); 7.75 (2H, m); 7.65 (1H, d); 7.30 7.60 (9H, m); 6.10 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 561 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.90 min.

Example 44.

15 1-(Thiophene-3-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 8.15 (1H, s); 7.95 (1H, m); 7.85 (1H, m); 7.60 (8H, m); 6.30 (1H, s); 3.45 (3H, s); 2.00-2.50 (8H, m). MS TOF 502 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.28 min.

Example 45.

1-(Thiophene-2-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

25 ¹H nmr (CDCl₃) 7.65 (2H, m); 7.45 (1H, s); 7.30 (2H, m); 7.20 (5H, m); 6.95 (1H, m); 6.00 (1H, s); 3.05 (3H, s); 3.00-2.50 (8H, m). MS TOF 502 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.52 min.

Example 46.

30 1-(5-Methylthiophene-2-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CDCl₃) 7.70 (1H, m); 7.45 (2H, m); 7.35 (6H, m); 6.65 (1H, m); 6.00 (1H, s); 3.05 (3H, s); 3.00-2.50 (8H, m) 2.45 (3H, s). MS TOF 516 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.98 min.

5

Example 47.

1-(Isoquinolin-7-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 9.50 (1H, s); 8.75 (1H, s); 8.55 (1H, d); 8.30 (1H, d); 8.10 (2H, m); 7.65 (1H, m); 7.45 (2H, m); 7.35 (5H, m); 6.10 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 547 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.39 min.

15 **Example 48.**

1-(Pyridin-3-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 9.00 (1H, s); 8.70 (1H, d); 8.35 (1H, d); 8.10 (1H, m); 7.65 (2H, m); 7.45 (1H, m); 7.30 (5H, m); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 497 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.99 min.

Example 49.

25 **1-(Indol-6-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine**

¹H nmr (CD₃CN) 7.95 (2H, m); 7.60 (2H, m); 7.50 (3H, m); 7.35 (5H, m); 6.45 (1H, s); 6.05 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 535 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.44 min.

30

Example 50.

1-(2,5-Diaminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

MS TOF 526 (M+1⁺). Hplc (Magellan C8, Gradient 3,

5 water/acetonitrile/TFA) rt 11.89 min.

Example 51.

1-(4-Methylaminobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

10 ¹H nmr (CD₃CN) 7.65 (3H, m); 7.50 (2H, m); 7.35 (5H, m);
6.60 (2H, d); 6.05 (1H, s); 3.30 (3H, s); 3.00-2.50 (8H, m);
2.80 (3H, s). MS TOF 525 (M+1⁺). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 13.17 min.

15 Example 52.

1-(3-Methyl-4-chlorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.90 (1H, s); 7.85 (1H, s); 7.80 (1H, s);
7.55 (6H, m); 6.25 (1H, s); 3.45 (3H, s); 3.00-2.50 (8H,
20 m); 2.60 (3H, s). MS TOF 545 (M+1⁺). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 16.39 min.

Example 53.

1-(4-Vinylbenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

25 ¹H nmr (CD₃CN) 7.75 (2H, d); 7.60 (1H, m); 7.45 (4H, m);
7.35 (5H, m); 6.75 (1H, m); 6.05 (1H, s); 5.90 (1H, d); 5.30
(1H, d); 3.00-2.50 (8H, m); 2.80 (3H, s). MS TOF 522 (M+1⁺).
Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
30 15.45 min.

Example 54.

1- (3-Amino-4-hydroxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.60 (1H, m); 7.50-7.10 (9H, m); 7.35 (1H, d); 5.95 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 527 (M+1⁺). Hplc (Magellan C8, Gradient 2, water/acetonitrile/TFA) rt 15.46 min.

Example 55.

10 1-(4-Methylthiobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.85 (2H, d); 7.80 (1H, m); 7.60 (2H, m); 7.50 (5H, m); 7.40 (2H, d); 6.15 (1H, s); 3.40 (3H, s); 3.10-2.70 (8H, m); 2.60 (3H, s). MS TOF 542 (M+1⁺). Hplc
15 (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 16.67 min.

Example 56.

20 1-(3-Carboxamidobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 8.25 (1H, s); 7.95 (2H, d); 7.70 (1H, m); 7.55 (3H, m); 7.40 (5H, m); 6.05 (1H, s); 3.30 (3H, s); 3.00-2.50 (8H, m). MS TOF 539 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.83 min.

25

Example 57.

1-(3-Amino-4-methylbenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.90 (1H, d); 7.70 (1H, m); 7.55 (2H, m);
30 7.45 (5H, m); 7.20 (1H, s); 6.95 (1H, d); 6.05 (1H, s); 3.80 (3H, s); 3.30 (3H, s); 3.00-2.50 (8H, m). MS TOF 569 (M+1⁺).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.49 min.

Example 58.

5 1-(3-Methyl-4-bromobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.65 (3H, m); 7.45 (3H, m); 7.30 (5H, m); 6.00 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m); 2.40 (3H, s). MS TOF 589 (M+1⁺). Hplc (Magellan C8, Gradient 3,

10 water/acetonitrile/TFA) rt 16.67 min.

Example 59.

1-(4-Ethoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

15 ¹H nmr (CD₃CN) 7.75 (2H, d); 7.60 (1H, m); 7.50 (2H, m); 7.35 (5H, m); 6.85 (2H, d); 6.00 (1H, s); 4.00 (2H, m); 3.20 (3H, s); 3.00-2.50 (8H, m); 1.30 (3H, t). MS TOF 540 (M+1⁺).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 16.58 min.

20

Example 60.

1-(Indol-5-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

25 ¹H nmr (CD₃CN) 8.15 (1H, s); 7.95 (1H, m); 7.65 (2H, m); 7.60-7.35 (7H, m); 6.60 (1H, s); 6.10 (1H, s); 3.30 (3H, s); 3.00-2.60 (8H, m). MS TOF 535 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.88 min.

Example 61.

30 1-(Benzimidazo-5-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 8.75 (1H, s); 8.25 (1H, s); 7.75 (2H, m);
7.60 (1H, m); 7.50 (2H, m); 7.35 (5H, m); 6.60 (2H, d); 6.05
(1H, s); 3.30 (3H, s); 3.00-2.50 (8H, m). MS TOF 536 (M+1⁺).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
5 10.08 min.

Example 62.

**1-(3-Aminobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine**

10 ¹H nmr (CD₃CN) a mixture of conformers only one recorded
here 7.65 (1H, m); 7.35 (5H, m); 7.05 (1H, m); 6.95 (2H, m);
5.85 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m);
2.90-2.40 (8H, m); 2.55 (3H, s); 1.60 (2H, m); 1.30 (2H, m);
1.00 (2H, m). MS TOF 435 (M+1⁺). Hplc (Magellan C8,
15 Gradient 3, water/acetonitrile/TFA) rt 7.65 min.

Example 63.

**1-(3-Amino-4-chlorobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine**

20 ¹H nmr (CD₃CN) a mixture of conformers only one recorded
here 7.75 (1H, m); 7.30 (5H, m); 7.20 (1H, m); 6.95 (1H, m);
5.85 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m);
2.90-2.40 (8H, m); 2.55 (3H, s); 1.60 (2H, m); 1.30 (2H, m);
1.00 (2H, m). MS TOF 469 (M+1⁺). Hplc (Magellan C8,
25 Gradient 3, water/acetonitrile/TFA) rt 9.58 min.

Example 64.

**1-(3-Amino-4-methylbenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine**

30 ¹H nmr (CD₃CN) a mixture of conformers only one recorded
here 7.75 (1H, m); 7.35 (5H, m); 7.05 (2H, m); 5.85 (1H, s);
4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m);

2.65 (3H, s); 2.15 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 449 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 8.03 min

5 Example 65.

1-(3-Aminonaphth-2-oyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded here 7.95 (1H, m); 7.65 (1H, d); 7.45 (2H, m); 7.30 (5H, m); 10 7.15 (1H, m); 6.95 (1H, s) 5.95 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m); 2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 485 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 9.94 min.

15

Example 66.

1-(Indol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded here 7.78 (2H, s); 7.50 (1H, d); 7.25 (7H, m); 6.34 (1H, s); 20 6.82 (1H, s); 4.40 (1H, m); 3.83 (1H, m); 3.35 (2H, t); 2.9-2.4 (8H, m) and 2.65 (3H, s) masked by water in solvent; 1.60 (2H, m); 1.40 (2H, m); 1.08 (2H, m). MS TOF 459 (M+1⁺). Hplc (Luna2 C18, Gradient 3, water/acetonitrile/TFA) rt 10.01 25 min.

Example 67.

1-(3-Amino-4-fluorobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (d₄ methanol) a mixture of conformers only one recorded here 7.4 (6H, m); 7.1 (1H, m); 7.0 (1H, t); 6.0 (1H, s); 4.63 (1H, m); 4.02 (1H, m); 3.30 (2H, m); 2.90-2.40

(8H, m); 2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 453 (M+1⁺).

Hplc (Symmetry C8, Gradient 3, water/acetonitrile/TFA) rt 5.03 min.

5

Example 68.

1-(3-Amino-4-bromobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

10 here 7.75 (1H, m); 7.35 (5H, m); 7.05 (1H, m); 6.80 (1H, m); 5.85 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m) and 2.65 (3H, s) masked by water in solvent; 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 513 and 515 (M+1⁺).

15 (Symmetry C8, Gradient 3, water/acetonitrile/TFA) rt 5.70 min.

Example 69.

1-(3-Amino-4-methoxybenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

20 here 7.70 (1H, m); 7.30 (5H, m); 7.0 (2H, m); 6.72 (1H, d); 5.80 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.70 (3H, s); 3.30 (2H, m); 2.9-2.4 (8H, m) masked by water in solvent;

25 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 465 (M+1⁺).
Hplc (Luna2 C18, Gradient 3, water/acetonitrile/TFA) rt 7.55 min.

Example 70.

30 1-(4-(Methylamino)benzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded here 7.70 (3H, m); 7.35 (5H, m); 6.60 (2H, d); 5.90 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.40 (2H, m); 2.9-2.4 (8H, m); 2.70 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m).

5 MS TOF 465 (M+1⁺).

Hplc (Luna2 C18, Gradient 3, water/acetonitrile/TFA) rt 8.52 min.

Example 71.

10 1-(4-Ethylaminobenzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.65 (3H, m); 7.45 (2H, m); 7.35 (5H, m); 6.60 (2H, d); 6.00 (1H, s); 3.20 (3H, s); 3.10 (2H, q); 3.00-2.50 (8H, m); 1.15 (3H, t). MS TOF 539 (M+1⁺). Hplc

15 (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.57 min.

Example 72.

1-(3-Methylaminobenzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

20 ¹H nmr (CD₃CN) 7.75 (1H, d); 7.60 (1H, d); 7.35 (7H, m); 7.15 (1H, t); 7.00 (1H, m); 6.70 (1H, d); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m); 2.70 (3H, s). MS TOF 525 (M+1⁺).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.07 min.

25

Example 73.

1-(4-Chloro-3-aminobenzoyl-D-phenylglyciny)-4-(2-methylsulphonylphenyl)piperazine

¹H nmr (CD₃CN) 7.95 (1H, d); 7.60 (1H, m); 7.45 (10H, m);

30 7.00 (1H, d); 6.00 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m).

MS TOF 527 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.56 min.

Example 74.

1-(4-Trifluoromethoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 5 1H nmr (CD₃CN) 7.85 (3H, m); 7.65 (1H, d); 7.45 (2H, m);
7.35 (6H, m); 6.00 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m).
MS TOF 580 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 16.01 min.

10 Example 75.

1-(4-Difluoromethoxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 1H nmr (CD₃CN) 7.85 (3H, m); 7.45 (2H, d); 7.30 (5H, m);
7.15 (2H, d); 6.80 (1H, t); 6.00 (1H, s); 3.20 (3H, s);
15 3.00-2.50 (8H, m). MS TOF 562 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 14.99 min.

Example 76.

- 1-(4-Trifluoromethylbenzoyl-D-phenylglyciny1)-N-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.85 (2H, d); 7.70 (2H, d); 7.45 (2H, m);
7.35 (6H, m); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m).
MS TOF 564 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 15.00 min.

25

Example 77.

1-(Indol-3-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 1H nmr (CD₃CN) 8.05 (1H, s); 7.85 (1H, d); 7.70 (1H, m); 7.50
30 (2H, m); 7.35 (6H, m); 7.20 (2H, m); 6.15 (1H, s); 3.20 (3H,
s); 3.00-2.50 (8H, m). MS TOF 535 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 14.25 min.

Example 78.

1-(4-Chloro-3-aminobenzoyl-L-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 5 1H nmr (CD₃CN) 7.75 (1H, d); 7.60 (1H, d); 7.45 (8H, m);
6.90 (1H, d); 5.95 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m).
MS TOF 545 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 14.53 min.

10 Example 79.

1-(2-Carboxybenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 1H nmr (CD₃CN) 7.75 (1H, d); 7.60 (1H, d); 7.50 (1H, d);
7.25-7.50 (9H, m); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H,
15 m). MS TOF 540 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 12.19 min.

Example 80.

1-(2-Fluorobenzoyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 20 1H nmr (CD₃CN) 7.85 (1H, m); 7.60 (1H, d); 7.25-7.50 (10H,
m); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF
514 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 13.29 min.

25

Example 81.

1-(3-Bromoindol-6-carbonyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- 1H nmr (CD₃CN) 7.85 (2H, m); 7.70-7.20 (10H, m); 6.05 (1H,
30 s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 614 (M+1+).
Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
16.16 min.

Example 82.

1-(3-Chloroindol-6-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

5 1H nmr (CD₃CN) 7.95 (2H, m); 7.70-7.30 (10H, m); 6.05 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 570 (M+1+).
Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 16.18 min.

10 Example 83.

1-(2-Cyanobenzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.25-7.80 (12H, m); 6.05 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 521 (M+1+). Hplc (Magellan
15 C8, Gradient 3, water/acetonitrile/TFA) rt 14.85 min.

Example 84.

1-(2-Aminomethylbenzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

20 1H nmr (CD₃CN) 7.95 (2H, m); 7.80-7.35 (10H, m); 6.15 (1H, s); 4.30 (2H, s); 3.15 (3H, s); 3.00-2.50 (8H, m). MS
TOF 525 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.21 min.

25 Example 85.

1-(4-Carboxy-3-aminobenzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.75 (1H, d); 7.60 (1H, d); 7.45 (7H, m); 7.15 (1H, s); 6.85 (1H, d); 5.95 (1H, s); 3.25 (3H, s);
30 3.00-2.50 (8H, m). MS TOF 554 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.00 min.

Example 86.

1- (1H-Indazol-6-carbonyl-D-phenylglycinyI)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 8.05 (2H, m); 7.85 (1H, d); 7.70 (1H, d); 7.55 (2H, m); 7.45 (5H, m); 5.95 (1H, s); 3.30 (3H, s); 3.00-2.50 (8H, m). MS TOF 545 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 13.44 min.

Example 87.

10 1-(4-Methylcarboxybenzoyl-D-phenylglycinyI)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.95 (2H, m); 7.80 (2H, m); 7.45 (2H, m); 7.35 (6H, m); 6.00 (1H, s); 3.90 (3H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 554 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.90 min.

Example 88.

1-(4-Acetoxybenzoyl-D-phenylglycinyI)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

20 1H nmr (CD₃CN) 7.75 (3H, m); 7.60 (1H, d); 7.45 (2H, m); 7.35 (5H, m); 7.10 (2H, d); 6.00 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m); 2.20 (3H, s). MS TOF 554 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.53 min.

Example 89.

1-(5-Methylpyrazin-2-carbonyl-D-phenylglycinyI)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

30 1H nmr (CD₃CN) 8.90 (1H, s); 8.35 (1H, s); 7.55 (1H, m); 7.40 (2H, m); 7.25 (5H, m); 5.85 (1H, s); 3.10 (3H, s); 3.00-2.50 (8H, m); 2.40 (3H, s). MS TOF 512 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.17 min.

Example 90.

1-(1,3-Benzodioxol-5-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

5 1H nmr (CD₃CN) 7.55 (2H, m); 7.35 (2H, m); 7.25 (6H, m);
6.70 (1H, d); 5.85 (2H, s); 5.80 (1H, s); 3.10 (3H, s);
3.00-2.50 (8H, m). MS TOF 540 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 14.28 min.

10 Example 91.

1-(4-(Methylsulphonyl)benzoyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.95 (3H, m); 7.60 (1H, m); 7.50 (2H, m); 7.35
(6H, m); 6.05 (1H, s); 3.25 (3H, s); 3.10 (3H, s); 3.00-2.50
15 (8H, m). MS TOF 574 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 13.62 min.

Example 92.

1-(2,3-Dichloroindol-6-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

20 1H nmr (CD₃CN) 7.90 (1H, d); 7.85 (1H, s); 7.55 (2H, m); 7.40
(2H, m); 7.25 (5H, m); 6.05 (1H, s); 3.30 (3H, s); 3.00-2.50
(8H, m); 2.40 (3H, s). MS TOF 614 (M+1+). Hplc (Magellan
C8, Gradient 3, water/acetonitrile/TFA)
25 rt 16.35 min.

Example 93.

1-(3-Chloro-2-oxo-(1H)indol-6-carbonyl-D-phenylglyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

30 1H nmr (CD₃CN) 7.90 (1H, d); 7.55 (1H, m); 7.25-7.50 (9H, m);
5.95 (1H, s); 5.20 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m).

MS TOF 585 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.38 min.

Example 94.

- 5 1-(3,3-Dichloro-2-oxo-(1H)indol-6-carbonyl-D-phenylglyciny)-
4-(4-fluoro-2-methylsulphonylphenyl)-piperazine

¹H nmr (CD₃CN) 7.90 (1H, d); 7.65 (2H, m); 7.55 (1H, m); 7.45 (2H, m); 7.35 (5H, m); 5.95 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m). MS TOF 619 (M+1+). Hplc (Magellan C8, Gradient 3,

- 10 water/acetonitrile/TFA) rt 15.13 min.

Example 95.

1-(3-Methylindol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine

- 15 ¹H nmr (CD₃CN) a mixture of conformers only one recorded
here 7.85 (2H, m); 7.40 (3H, m); 7.30 (3H, m); 7.05 (1H, s);
5.95 (1H, s); 4.55 (1H, m); 3.85 (1H, m); 3.30 (2H, m);
2.90-2.40 (8H, m); 2.55 (3H, s); 2.20 (3H, s); 1.60 (2H, m);
1.30 (2H, m); 1.00 (2H, m). MS TOF 473 (M+1+). Hplc
20 (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.40
min.

Example 96.

1-(2,3-Dihydroindol-6-carbonyl-D-phenylglyciny)-1'-methyl-
25 4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded
here 7.75 (1H, m); 7.30 (7H, m); 5.85 (1H, s); 4.45 (1H, m);
3.85 (1H, m); 3.65 (2H, t); 3.30 (2H, m); 3.10 (2H, t);
2.90-2.40 (8H, m); 2.55 (3H, s); 1.60 (2H, m); 1.30 (2H, m);

- 30 1.00 (2H, m). MS TOF 461 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 8.68 min.

Example 97.

1-(1H-indazol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

5 here 7.95 (1H, m); 7.85 (2H, m); 7.65 (1H, m); 7.45 (2H, m);
7.30 (3H, m); 5.95 (1H, s); 4.55 (1H, m); 3.95 (1H, m); 3.30
(2H, m); 2.90-2.40 (8H, m); 2.55 (3H, s); 1.60 (2H, m); 1.30
(2H, m); 1.00 (2H, m). MS TOF 460 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 9.72 min.

10

Example 98.

1-(Benzimidazol-5-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

15 here. 8.05 (1H, s); 7.90 (1H, m); 7.75 (2H, m); 7.30 (5H, m);
5.95 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m);
2.90-2.40 (8H, m); 2.75 (3H, s); 1.60 (2H, m); 1.30 (2H, m);
1.00 (2H, m). MS TOF 460 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 8.80 min.

20

Example 99.

1-(Benzthiazol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

25 here 8.40 (1H, s); 7.95 (3H, m); 7.30 (5H, m); 5.85 (1H, s);
4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m);
2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS
TOF 477 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 9.58 min.

30

Example 100.

1-(3-Chloroindol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

5 here 7.85 (2H, m); 7.30 (7H, m); 5.85 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m); 2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 493 (M+1+).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.22 min.

10

Example 101.

1-(3-Bromoindol-6-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CD₃CN) a mixture of conformers only one recorded

15 here 7.85 (2H, m); 7.30 (7H, m); 5.85 (1H, s); 4.45 (1H, m); 3.85 (1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m); 2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 539 (M+1+).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.45min.

20

Example 102.

1-(3-Amino-4-chlorobenzoyl-L-phenylglyciny)-1'-methyl-4,4'-bispiperidine

¹H nmr (CDCl₃) a mixture of conformers only one recorded

25 here 7.65 (1H, m); 7.30 (6H, m); 7.00 (1H, m); 5.85 (1H, s); 4.65 (1H, m); 3.80 (1H, m); 3.55 (2H, m); 2.90-2.40 (8H, m); 2.65 (3H, s); 1.60 (2H, m); 1.30 (2H, m); 1.00 (2H, m). MS

TOF 469 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 10.71min.

30

Example 103.

1-(4-Vinylbenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine

1H nmr (CD₃CN) a mixture of conformers only one recorded
5 here 7.85 (1H, m); 7.70 (2H, m); 7.40 (6H, m); 6.75 (1H, m);
6.00 (1H, s); 5.85 (1H, d); 5.50 (1H, d); 4.55 (1H, m); 3.95
(1H, m); 3.30 (2H, m); 2.90-2.40 (8H, m); 2.65 (3H, s); 1.60
(2H, m); 1.30 (2H, m); 1.00 (2H, m). MS TOF 446 (M+1+).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
10 11.21min.

Example 104.

1-(3-Amino-4-chlorobenzoyl-D-phenylglyciny)-4-(4-amino-2-
methylsulphonylphenyl)piperazine

15 1H nmr (CD₃CN) 7.55 (1H, m); 7.45 (3H, m); 7.35 (5H, m);
7.10 (1H, d); 6.90 (1H, d); 6.10 (1H, s); 3.20 (3H, s);
3.00-2.50 (8H, m). MS TOF 542 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 12.02 min.

20 Example 105.

1-(3-Aminobenzoyl-D-phenylglyciny)-4-(4-amino-2-methyl
sulphonylphenyl)piperazine

1H nmr (CD₃CN) 7.55 (2H, m); 7.45 (3H, m); 7.35 (5H, m);
7.10 (1H, d); 6.90 (1H, d); 6.10 (1H, s); 3.10 (3H, s);
25 3.00-2.50 (8H, m). MS TOF 508 (M+1+). Hplc (Magellan C8,
Gradient 3, water/acetonitrile/TFA) rt 9.35 min.

Example 106.

1-(3-Amino-4-chlorobenzoyl-D-phenylglyciny)-4-(4-carboxamido-
30 2-methylsulphonylphenyl)piperazine

1H nmr (CD₃CN) 8.05 (1H, d); 7.80 (1H, m); 7.35-7.60 (8H, m);
7.10 (1H, d); 6.10 (1H, s); 3.25 (3H, s); 3.00-2.50 (8H, m).

MS TOF 570 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.24 min.

Example 107.

- 5 1-(3-Amino-4-chlorobenzoyl-D-phenylglyciny1)-4-(4-nitro-2-methylsulphonylphenyl)piperazine
- 1H nmr (CD3CN) 8.70 (1H, s); 8.45 (1H, d); 7.55 (1H, m); 7.45 (5H, m); 7.30 (2H, m); 7.10 (1H, d); 6.10 (1H, s); 3.40 (3H, s); 3.00-2.50 (8H, m). MS TOF 572 (M+1+). Hplc (Magellan
- 10 C8, Gradient 3, water/acetonitrile/TFA) rt 14.25 min.

Example 108.

- 15 1-(3-Amino-4-chlorobenzoyl-D-4-aminophenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
- 1H nmr (CD3CN) 7.65 (1H, d); 7.45 (4H, m); 7.25 (2H, m); 7.15 (2H, d); 7.05 (1H, d); 6.10 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 560 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.90 min.

20 Example 109.

- 1-(3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine
- 1H nmr (CD3CN) 7.70 (2H, d); 7.55 (1H, d); 7.45 (2H, d); 7.25 (2H, m); 7.20 (2H, d); 6.90 (1H, d); 6.10 (1H, s); 3.20 (3H, s); 3.00-2.50 (8H, m). MS TOF 588 (M+1+). Hplc
- 25 (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 12.18 min.

Example 110.

- 30 1-(3-Amino-4-chlorobenzoyl-D-4-(methylcarboxamido)phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

1H nmr (CD3CN) 7.70 (2H, d); 7.55 (1H, d); 7.45 (2H, d);
7.25 (2H, m); 7.20 (2H, d); 6.90 (1H, d); 6.10 (1H, s); 3.20
(3H, s); 2.70 (3H, s); 3.00-2.50 (8H, m). MS TOF 602 (M+1+).

Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt
5 12.70 min.

Example 111.

3-Amino-4-chlorobenzoyl-D-phenylglycine 4-methylbenzylamide

1H nmr (CD3CN) 7.55 (1H, m); 7.35 (7H, m); 7.00 (4H, m); 5.45
10 (1H, s); 4.25 (2H, m); 2.20 (3H, s). MS TOF 408 (M+1+). Hplc
(Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.61
min.

Example 112.

15 3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglycine R,S -2-
methylcyclohexylamide

1H nmr (CD3CN) mixture of isomers only one recorded here
7.75 (2H, d); 7.60 (2H, m); 7.30 (2H, m); 7.10 (1H, d); 5.55
(1H, s); 3.90 (1H, m); 3.25 (1H, m); 1.00-2.00 (8H, m) 0.50
20 (3H, m). MS TOF 443 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 9.18 min

Example 113.

3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglycine 2-
25 indanamide

MS TOF 463 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 12.58 min.

Example 114.

30 3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglycine (S)-N -
benzyl-alpha-methylbenzylamide

MS TOF 541 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 15.34 min.

Example 115.

- 5 3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglycine 1-(S)-1-naphthylethylamide

MS TOF 5013 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.00 min.

- 10 Example 116.

3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglycine 3-(1-(R,S)-hydroxyethyl)benzamide

MS TOF 443 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 11.81 min.

15

Example 117.

3-Amino-4-chlorobenzoyl-D-phenylglycine cis,trans-2-aminocyclohexylamide

- MS TOF 401 (M+1+). Hplc (Magellan C8, Gradient 3,
20 water/acetonitrile/TFA) rt 11.00 min.

Example 118.

1-(3-Amino-4-chlorobenzoyl-D,L-(4-piperidinyl)glyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- MS TOF 552 (M+1+). Hplc (Magellan C8, Gradient 3,
25 water/acetonitrile/TFA) rt 11.00 min.

Example 119.

1-(3-Amino-4-chlorobenzoyl-D,L-(4-N-methylpiperidinyl)-glyciny)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

- MS TOF 566 (M+1+). Hplc (Magellan C8, Gradient 3,
30 water/acetonitrile/TFA) rt 10.83 min.

Example 120.

1- (3-Amino-4-chlorobenzoyl-D,L- (4-N-trifluoroacetyl-
piperidinyl)glyciny1-4- (4-fluoro-2-methylsulphonylphenyl) -
5 piperazine

MS TOF 649 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 12.63 min.

Example 121.

10 3-Amino-4-chlorobenzoyl-D-phenylglycine (2-chloro-5-
carboxamido)benzenesulphonamide

MS TOF 521 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 10.23 min.

15 Example 122.

1- (4-Cyanobenzoyl-D-phenylglyciny1) -1'-methyl-4,4'-
bispiperidine

MS TOF 445 (M+1+). Hplc (Magellan C8, Gradient 3,
water/acetonitrile/TFA) rt 10.13min.

20

Example 123.

1- (3-Cyanobenzoyl-D-phenylglyciny1) -1'-methyl-4,4'-
bispiperidine

MS TOF 445 (M+1+). Hplc (Magellan C8, Gradient 3,
25 water/acetonitrile/TFA) rt 10.23min.

Example 124.

1- (4-Chlorobenzoyl-D-phenylglyciny1) -4- (4-pyridyl) -piperazine

MS TOF 435 (M+1+). Hplc (Magellan C8, Gradient 3,
30 water/acetonitrile/TFA) rt 12.11 min.

Example 125.

1-(4-Methoxybenzyl-D-phenylglyciny1)-4-(4-fluoro-2-methylsulphonylphenyl)piperazine

MS TOF 512 (M+1+). Hplc (Magellan C8, Gradient 3,
5 water/acetonitrile/TFA) rt 11.91 min.

Example 126.

1-N-(3-Amino-4-chlorobenzoyl)-2-N-(4-methoxybenzoyl)-1,2-diamino-1-phenylethane

10 ¹H nmr (CD₃OH) 7.45 (2H, m); 7.35 (3H, m); 7.20 (2H, m); 7.10 (3H, m); 6.75 (2H, d); 4.80 (1H, m); 4.25 (2H, m); 3.70 (3H, s). MS TOF 424 (M+1+). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.05 min.

15 Examples 127 to 136.

Preparation of Starting Materials

4-methoxybenzoyl-D-phenylglyciny1-R,S-3-hydroxypyrrolidine
D-phenylglyciny1-R,S-3-hydroxypyrrolidine (3.42g, 15.5mmol)
20 was dissolved in dichloromethane (100ml) and placed under argon. Triethylamine (2.27ml, 16.28mmol) was added followed by 4-methoxybenzoyl chloride (2.78g, 16.3mmol) and the mixture stirred at room temperature for 3.5h. The organic solution was washed with 0.5% hydrochloric acid (50ml), sat.
25 sodium bicarbonate solution (50ml) and brine (50ml). The organic solution was dried (MgSO₄) and evaporated to an off-white solid, 4-methoxybenzoyl-D-phenylglyciny1-R,S-3-hydroxypyrrolidine, (5.49g, 100%)

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
30 11.7min

LCMS M+1 355 Nmr.

4-methoxybenzoyl-D-phenylglyciny-4-hydroxypiperidine

By a similar method D-phenylglyciny-4-hydroxypiperidine was converted to 4-methoxybenzoyl-D-phenylglyciny-4-hydroxypiperidine.

- 5 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 11.9min

LCMS M+1 369 Nmr

Example 127

- 10 **1-(4-Methoxybenzoyl-D-phenylglyciny)-3-(R,S)-(2-fluorophenoxy)pyrrolidine**

To a solution of 4-methoxybenzoyl-D-phenylglyciny-R,S-3-hydroxypyrrolidine (400mg, 1.13mmol) in benzene (10ml) at 10°C was added 2-triphenylphosphonium 4,4-dimethyl-

- 15 tetrahydro-1,2,5-thiadiazolidine 1,1-dioxide (Reference: J. Castro et al. J. Org. Chem. 1994, 59, 2289-2291) (696mg, 1.69mmol) and 3-methoxyphenol (210mg) and the mixture allowed to warm to room temperature overnight. The reaction mixture was diluted with ether (30ml) and washed with dilute
20 sodium bicarbonate solution. The organic solution was dried (MgSO₄) and concentrated. The residue was purified by reverse phase preparative chromatography to give 1-(4-methoxybenzoyl-D-phenylglyciny)-3-(R,S)-(3-methoxyphenoxy)pyrrolidine.

- 25 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 11.75min.

LCMS M+1 461 Nmr (mixture of diastereomers).

Example 128.

- 30 **1-(4-Methoxybenzoyl-D-phenylglyciny)-3-(R,S)-(3-methoxyphenoxy)pyrrolidine**

From 4-methoxybenzoyl-D-phenylglyciny1-R,S-3-hydroxypyrrolidine and 3-methoxyphenol:
Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 11.75min.

5 LCMS M+1 461 Nmr (mixture of diastereomers).

Example 129.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(3-methoxyphenoxy)piperidine

10 From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 3-methoxyphenol:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 16.09min

LCMS M+1 475. Nmr

15

Example 130.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(4-methoxyphenoxy)piperidine

20 From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 4-methoxyphenol:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 15.8min.

LCMS M+1 475. Nmr.

25 Example 131.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(3-fluorophenoxy)piperidine

From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 3-fluorophenol:

30 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 12.75min.

LCMS M+1 463 Nmr

Example 132.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(2-methanesulfonylphenoxy)piperidine

- 5 From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 2-methanesulphonylphenol:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 10.8min.

LCMS M+1 523 Nmr.

10

Example 133.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(2-methylmercaptophenoxy)piperidine

- 15 From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 2-methylmercaptophenol:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 12.7min

LCMS M+1 491 Nmr.

20 Example 134.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(2-fluorophenoxy)piperidine

From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and 2-fluorophenol:

- 25 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 15.8min.

LCMS M+1 463 Nmr.

Example 135.

- 30 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(phenoxy)piperidine

From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine and phenol:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
16.8min.

LCMS M+1 445

5 Example 136.

1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(3-pyridoxy)piperidine

From 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine
and 3-hydroxypyridine:

10 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
11.4min

LCMS M+1 446 Nmr

Example 137.

15 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(4-fluorophenoxy)piperidine

To a solution of triphenylphosphine (285mg, 1.09mmol) in dry THF (5ml) under argon at -15°C was added slowly (<-10°C) diethyl azodicarboxylate (DEAD) (208mg, 1.19mmol) and the
20 solution stirred at <-10°C for 5min. To this mixture was added a solution of 4-methoxybenzoyl-D-phenylglyciny1-4-hydroxypiperidine (400mg, 1.08mmol) and 4-fluorophenol (122mg, 1.09mmol) in dry THF (5ml) over 5min at <-10°C. The
25 reaction was warmed to room temperature and monitored by tlc (SiO₂ - ethyl acetate). The reaction mixture was poured into water (5ml) and extracted with dichloromethane (100ml). The organic solution was washed with sat. sodium bicarbonate (50ml) and 0.5% hydrochloric acid (50ml), dried (MgSO₄) and concentrated and the residue purified by flash
30 chromatography, (SiO₂ - 30% ethyl acetate in hexane to give 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(4-fluorophenoxy)piperidine, (107mg, 21%)

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,

16.0min

LCMS M+1 463. Nmr.

5 Examples 138 to 142

Preparation of Starting Materials

Benzyloxycarbonyl-D-phenylglycinyll-R,S-3-hydroxypyrrolidine

Benzyloxycarbonyl-D-phenylglycine (18.01g, 63.1mmol) and

10 R,S-3-hydroxypyrrolidinol (5.0g, 57.4mmol) were suspended in dimethylformamide (300ml). HOAt (8.61g, 63.1mmol) was added, the mixture stirred for 3min. and then EDCI (12.1g 63.1mmol) was added with stirring and the mixture left overnight. The orange solution was concentrated *in vacuo* and the residue
15 taken up in ethyl acetate (300ml). The organic solution was washed with sat. sodium bicarbonate (2 x 100ml), 0.5% aqueous hydrochloric acid (50ml) and brine (100ml). The organic solution was dried (MgSO₄) and evaporated *in vacuo* to give an orange solid. Flash chromatography (SiO₂ 1:1
20 dichloromethane: ethyl acetate gave benzyloxycarbonyl-D-phenylglycinyll-R,S-3-hydroxypyrrolidine, (11.4g, 56%).
Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
12.7min

LCMS M+1 355 Nmr.

25

Benzyloxycarbonyl-D-phenylglycinyll-4-hydroxypiperidine

By a similar method using benzyloxycarbonyl-D-phenylglycine and 4-hydroxypiperidine, benzyloxycarbonyl-D-phenylglycinyll-4-hydroxypiperidine was prepared.

30 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
11.9min

LCMS M+1 369 Nmr.

D-Phenylglycinyll-R,S-3-hydroxypyrrolidine

Benzyloxycarbonyl-D-phenylglycinyll-R,S-3-hydroxypyrrolidine,
(5.49g, 15.5mmol) was dissolved in ethanol (120ml) and Pd/C
5 (10%, 100mg) added. The mixture was hydrogenated at
atmospheric pressure until complete by tlc (SiO₂ ethyl
acetate - starting material Rf. 0.6, product 0.05). The
catalyst was filtered off through celite and concentrated in
vacuo to give D-phenylglycinyll-R,S-3-hydroxypyrrolidine as a
10 yellow oil, (3.54g, 16.1mmol).

D-Phenylglycinyll-4-hydroxypiperidine

By a similar method benzyloxycarbonyl-D-phenylglycinyll-4-
hydroxypiperidine was converted to D-phenylglycinyll-4-
15 hydroxypiperidine

Benzyloxycarbonyl-D-phenylglycinyll-4-(3-pyridoxy)piperidine

To a solution of benzyloxycarbonyl-D-phenylglycinyll-4-
hydroxypiperidine (500mg, 1.36mmol), 3-hydroxypyridine
20 (129mg, 1.36mmol) and triphenylphosphine (356mg, 1.36mmol)
in dry THF (20ml) at 0°C, was slowly added diethyl
azodicarboxylate (259mg, 1.19mmol) and the mixture stirred
for 1h at 0°C and then 16h at room temperature. Water (5ml)
was added and the mixture extracted with ethyl acetate (2 x
25 10ml). The organic solution was washed with water and brine,
dried (MgSO₄) and concentrated to an oil which was purified
by flash chromatography, (SiO₂ - hexane/ethyl acetate 1:1)
to give benzyloxycarbonyl-D-phenylglycinyll-4-(3-
pyridoxy)piperidine, (490mg 65% - contaminated with
30 triphenylphosphine)

**Benzyloxycarbonyl-D-phenylglycinyll-R,S-3-(3-pyridoxy) -
pyrrolidine**

A solution of benzyloxycarbonyl-D-phenylglycinyll-R,S-3-
hydroxypyrrolidine (2.0g, 8.64mmol), 2-triphenylphosphonium
5 4,4-dimethyl-tetrahydro-1,2,5-thiadiazolidine 1,1-dioxide
(Reference: J. Castro et al. J. Org. Chem. 1994, 59, 2289-
2291) (3.479g, 8.47mmol) and 3-hydroxypyridine (0.805g,
8.47mmol) in benzene (30ml) was stirred at room temperature
for 18h. The mixture was poured onto ether (50ml) and the
10 organic solution was washed with sat. sodium bicarbonate (2
x 50ml). The product was extracted into 5% hydrochloric acid
which was then basified (pH8) with 2M sodium hydroxide
solution and extracted with ether (3 x 100ml). The organic
solution was dried (MgSO₄) and evaporated to give
15 benzyloxycarbonyl-D-phenylglycinyll-R,S-3-(3-
pyridoxy)pyrrolidine

D-Phenylglycinyll-4-(3-pyridoxy)piperidine

Benzyloxycarbonyl-D-phenylglycinyll-4-(3-pyridoxy)piperidine
20 (1.18g 2.64mmol) was dissolved in ethanol (120ml) containing
Pd/C 10% (100mg) and acetic acid (0.3ml) and hydrogenated at
atmospheric pressure for 8h - (incomplete by tlc). The
catalyst was removed by filtration and the solution
evaporated to an oil. The oil was re-hydrogenated as before.
25 The catalyst was removed by filtration and the solvent
evaporated in vacuo to an oil which was taken up in dilute
hydrochloric acid. The aqueous solution was washed with
dichloromethane and then basified with solid sodium
bicarbonate. Extraction with chloroform, drying (MgSO₄) and
30 evaporation of the solvent in vacuo gave D-phenylglycinyll-4-
(3-pyridoxy)piperidine, (331mg 40%). Nmr

D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine

In a similar manner D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine was prepared from benzyloxycarbonyl-D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine by

5 hydrogenation over Pd/C in ethanol. Nmr.

Example 138.

1-(Indole-6-carbonyl-D-phenylglyciny1)-4-(3-pyridoxy)piperidine

10 A mixture of EDCI (169mg 0.88mmol), HOAt (120mg 0.88mmol) and indole-6-carboxylic acid (142mg 0.88mmol) in DMF (5ml) was stirred for 2min and then added to a solution of D-phenylglyciny1-4-(3-pyridoxy)piperidine (229mg 0.735mmol) and triethylamine (89mg 0.88mmol) in DMF (20ml). The mixture
15 was stirred at room temperature for 3h and excess solvent removed in vacuo. The residue was taken up in ethyl acetate (150ml) and washed with sat. sodium bicarbonate (50ml). The solution was dried (MgSO₄), evaporated and the residue purified by flash chromatography (SiO₂ ethyl acetate:
20 methanol 0% - 5%) to give 1-(indole-6-carbonyl-D-phenylglyciny1)-4-(3-pyridoxy)piperidine (122mg 41%)
Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 10.8min.

LCMS M+1 455 Nmr

25

The following were prepared in a similar manner:

Example 139.

1-(3-Chloroindole-6-carbonyl-D-phenylglyciny1)-4-(3-pyridoxy)piperidine

30

From D-phenylglyciny1-4-(3-pyridoxy)piperidine and 3-chloro-6-indolecarboxylic acid:

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt
11.95min

LMCS M+1 489 Nmr

5 Example 140.

1-(Indole-6-carbonyl-D-phenylglyciny1)-3-(R,S)-(3-pyridoxy)pyrrolidine

From D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine and 6-indolecarboxylic acid.

10 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
6.4min.

LCMS M+1 441 Nmr (mixture of diastereomers).

Example 141.

15 1-(3-Chloroindole-6-carbonyl-D-phenylglyciny1)-3-(R,S)-(3-pyridoxy)pyrrolidine

From D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine and 3-chloro-6-indolecarboxylic acid.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,

20 7.2min.

LCMS M+1 475 Nmr (mixture of diastereomers).

Example 142.

1-(3-Methylindole-6-carbonyl-D-phenylglyciny1)-3-(R,S)-(3-pyridoxy)pyrrolidine

25 From D-phenylglyciny1-R,S-3-(3-pyridoxy)pyrrolidine and 3-methyl-6-indolecarboxylic acid.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 6.84
and 7.0min.

30 LCMS M+1 455 Nmr (mixture of diastereomers).

Example 143.

(R)-2-(1'-(3-Chloroindole-6-carboxamido)benzyl)-4-methoxyphenyl-1,3-thiazole

5 (R)-2-(1'-benzyloxycarbonylamidobenzyl)-4-methoxyphenyl-1,3-thiazole

To a solution of benzyloxycarbonyl-D-phenylglycine thioamide (1g, 3.33mmol.) in acetone (25ml) was added α -bromo-4-methoxyacetophenone (0.76g, 3.32mmol) and the mixture
10 stirred at room temperature for 30min. Chloroform (25ml) and sat. aqueous sodium hydrogen carbonate (30ml) were added and the organic solution separated, dried (MgSO₄) and evaporated in vacuo. The residue was dissolved in
15 dichloromethane (30ml) and pyridine (0.5ml, 6.18mmol) and trifluoroacetic anhydride (0.5ml, 3.54mmol) were added. The mixture was stirred at room temperature until complete by
tlc (SiO₂ dichloromethane - 1h.), washed with 5% hydrochloric acid, dried (MgSO₄) and evaporated in vacuo. Flash chromatography of the residue (0.87g). (SiO₂ -
20 dichloromethane) gave (R)-2-(1'-benzyloxycarbonylamidobenzyl)-4-methoxyphenyl-1,3-thiazole (0.74g 1.72mmol. 52%)

Nmr: CDCl₃ 7.85(2H, d), 7.3-7.5 (11H, m), 6.95 (2H, d), 6.44 (0.5H, bd), 6.16 (0.5H, bd), 5.02-5.22 2H, m), 3.83 (3H, m).

25

(R)-2-(1'-aminobenzyl)-4-methoxyphenyl-1,3-thiazole

(R)-2-(1'-Benzyloxycarbonylamidobenzyl)-4-methoxyphenyl-1,3-thiazole (0.70g, 1.63mmol) was dissolved in acetic acid (50ml) and HBr in acetic acid (25ml) added. The mixture was
30 heated in a 50°C oil bath for 2h when no starting material remained by tlc (SiO₂ 30% ether in dichloromethane). The

mixture was evaporated in vacuo, basified with sat. aqueous sodium hydrogen carbonate and extracted with ethyl acetate (x3). The organic solution was dried (MgSO₄) and evaporated in vacuo. Flash chromatography (SiO₂ dichloromethane then 30% ether in dichloromethane) gave (R)-2-(1'-aminobenzyl)-4-methoxyphenyl-1,3-thiazole (172mg, 36%)

Nmr: CDCl₃ 7.7 (2H, d), 7.5 (2H, d), 7.17-7.4 (3H, m), 6.85 (2H, d), 3.76 (3H, s)

(R)-2-(1'-(3-Chloroindole-6-carboxamido)benzyl)-4-methoxyphenyl-1,3-thiazole

(R)-2-(1'-Aminobenzyl)-4-methoxyphenyl-1,3-thiazole (80mg, 0.27mmol) was coupled to 3-chloroindolecarboxylic acid using EDC/HOAt to give: (R)-2-(1'-(3-Chloroindole-6-carboxamido)benzyl)-4-methoxyphenyl-1,3-thiazole (49%)

Hplc (Luna C18 Gradient3) rt 17.2min.

LCMS M+1 474. Nmr.

Examples 144 to 147.

The compounds of Examples 144 to 147 were prepared by coupling to the appropriate carboxylic acid to D-phenylglyciny-4,4'-(1'-methylbispiperidine) using EDC and HOAt as described previously.

Example 144.

1-(4-Methylbenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

Hplc (Luna C18 Gradient3) rt 11.2min.

LCMS M+1 434. Nmr.

Example 145.

1-(4-Chlorobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine

Hplc (Luna C18 Gradient3) rt 11.5min.

5 LCMS M+1 454. Nmr.

Example 146.

1-(4-Methoxybenzoyl-D-phenylglyciny)-1'-methyl-4,4'-
bispiperidine

10 Hplc (Luna C18 Gradient3) rt 11.1min.

LCMS M+1 450. Nmr.

Example 147

1-(3,4-Methylenedioxybenzoyl-D-phenylglyciny)-1'-methyl--
4,4'-bispiperidine

15

Hplc (Luna C18 Gradient3) rt 10.65min.

LCMS M+1 464. Nmr.

Example 148.

20 1-(Indole-6-carbonyl-D-phenylglyciny)-1'-isopropyl-4,4'-
bispiperidine

Benzyloxycarbonyl-D-phenylglyciny-4,4'-(1'-bispiperidine)

25 Benzyloxycarbonyl-D-phenylglyciny- 1'-isopropyl-4,4'-
bispiperidine

D-phenylglyciny-1'-isopropyl-4,4'-bispiperidine

30 1-(Indole-6-carbonyl-D-phenylglyciny)- 1'-isopropyl-4,4'-
bispiperidine

Prepared by coupling the appropriate carboxylic acid to D-phenylglyciny-4,4'-(1'-(2''-propyl)bispiperidine).

Hplc (Luna C18 Gradient3) rt 11.46min.

LCMS M+1 487. Nmr.

5

Examples 149 to 154.

The compounds of Examples 149 to 154 were prepared by coupling Boc-D-4-carboxamidophenylglycine to the appropriate
10 amine with EDCI/HOAt, deprotection with TFA/DCM and coupling to 3-amino-4-chlorobenzoic acid with EDCI/HOAt as previously described.

Example 149.

15 2-(3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglyciny-1,2,3,4-tetrahydroisoquinoline

Hplc (Luna C18 Gradient3) rt 13.15min.

LCMS M+1 463. Nmr.

20 **Example 150.**

1-(3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglyciny)-4-benzylpiperazine

Hplc (Luna C18 Gradient3) rt 11.4min.

LCMS M+1 512. Nmr.

25

Example 151.

1-(3-Amino-4-chlorobenzoyl-D-4-carboxamidophenylglyciny)-4-(2-methylthiophenyl)piperazine

Hplc (Luna C18 Gradient3) rt 14.3min.

30 LCMS M+1 539. Nmr.

Example 152.

1- (3-Amino-4-chlorobenzoyl-D-4-carboxamidophenyl-
glyciny1)-4- (2-phenylethyl)piperazine

Hplc (Luna C18 Gradient3) rt 11.1min.

5 LCMS M+1 521. Nmr.

Example 153.

1- (3-Amino-4-chlorobenzoyl-D-4-carboxamidophenyl-
glyciny1)-4-benzoylpiperidine

10 Hplc (Luna C18 Gradient3) rt 12.8min.

LCMS M+1 520. Nmr.

Example 154.

1- (3-Amino-4-chlorobenzoyl-D-4-carboxamidophenyl-
15 glyciny1)-4- (2-ethylphenyl)piperazine

Hplc (Luna C18 Gradient3) rt 13.9min.

LCMS M+1 521. Nmr.

Example 155.

20 1- (3-Methoxyindole-6-carbonyl-D-phenylglyciny1)-1'-
methyl-4,4'-bispiperidine

Methyl 1-acetyl-3-formylindole-6-carboxylate

A suspension of methyl 3-formylindole-6-carboxylate (1g,
25 4.93 mmol) in acetic anhydride (10ml) was refluxed for 2 h.
The acetic anhydride was removed under reduced pressure to
afford a pinkish solid (1.2g, 100%) that was used without
further purification. ¹H NMR (CDCl₃) 2.7 (3H, s), 3.9 (3H,
s), 8.05 (1H, d), 8.15 (1H, s), 8.25 (1H, d), 9.0 (1H, s),
30 10.1 (1H, s); LCMS M+H 246.

Methyl 1-acetyl-2,3-dihydroindol-3-one-6-carboxylate

This was prepared from methyl 1-acetyl-3-formylindole-6-carboxylate (1.03g, 4.20 mmol) using the method of Merour et al. (*Synthesis*, 1994, 411) to yield the formate (680 mg).

5 The formate was dissolved in THF (50ml) and treated with sat. NaHCO₃ solution (10ml). After 15 min. the reaction mixture was extracted with ethyl acetate, washed with water, dried and concentrated to give the ketone (574mg). ¹H NMR (CDCl₃) 2.3 (3H, br.), 3.9 (3H, s), 4.3 (2H, s), 7.75 (1H, d), 7.85 (1H, d), 9.1 (1H, br.); LCMS M+H 234.

Methyl 1-acetyl-3-methoxyindole-6-carboxylate

Methyl 1-acetyl-2,3-dihydroindol-3-one-6-carboxylate (233mg, 1 mmol), trimethyl orthoformate (10ml) and *p*-toluene sulphonic acid (20 mg) were heated under reflux for 3 h. in methanol (10ml). The reaction mixture was concentrated under reduced pressure, poured into water and extracted with chloroform. After drying and evaporation, the product was purified by prep hplc; ¹H NMR (CD₃CN) 2.56 (3H, s), 3.93 (3H, s), 3.97 (3H, s), 7.25 (1H, s), 7.62 (1H, d), 7.90 (1H, d), 9.0 (1H, br.); LCMS M+H 248.

3-Methoxyindole-6-carboxylic acid

To a solution of methyl 1-acetyl-3-methoxyindole-6-carboxylate (74 mg, 0.3 mmol) in THF (10ml) and water (2ml) was added lithium hydroxide hydrate (63 mg, 1.5 mmol). The reaction mixture was warmed to 50°C and stirred for 3 h. The THF was removed under reduced pressure and the pH of the aqueous phase adjusted to 3. Extraction of the aqueous layer with ethyl acetate, drying and concentration gave the acid (50 mg, 87%); ¹H NMR (CD₃CN) 3.75 (3H, s), 3.97 (3H, s), 6.9

(1H, s), 7.45 (1H, d), 7.55 (1H, d), 8.2 (1H, s); LCMS M+H
192.

1- (3-Methoxyindole-6-carbonyl-D-phenylglyciny1)-4,4'-(1'-
5 methylbispiperidine)

Prepared by coupling to D-phenylglyciny1-4,4'-(1'-
methylbispiperidine) using EDC and HOAt as described
previously.

Hplc (Luna C18, Gradient3) rt 8.35min.

10 LCMS M+1 489 Nmr.

Example 156.

1- (3-Amino-4-chlorobenzoyl-D-cyclohexylglyciny1)-4-(4-
fluoro-2-methylsulfonylphenyl)-piperazine

15 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt
15.37min.

LCMS M+1 551

Example 157.

20 1- (3-Amino-4-chlorobenzoyl-D,L-1-naphthylglyciny1)-4-(4-
fluoro-2-methylsulfonylphenyl)-piperazine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt
15.69min.

LCMS M+1 595

25

Example 158.

1- (3-Chloroindole-6-carbonyl-D,L-(2-methylthiazol-4-
yl)glyciny1)-1'-methyl-4,4'-bispiperidine

Ethyl oximinoacetoacetate

This was prepared from ethyl acetoacetate (10.00g) using the method of Fischer (*Organic Synthesis Coll. Vol. 3*, 513-516) to yield the titled compound (12.45g); ¹H NMR (CDCl₃) 1.25

5 (3H, t), 2.35 (3H, s), 4.3 (2H, q), 8.8 (1H, br.).

Ethyl-γ-chloro-α-oximinoacetoacetate

This was prepared from ethyl oximinoacetoacetate (1.73g) using the method of Hatanaka et al. (*Journal of Medicinal*
10 *Chemistry*, 1973, 16(9), 978-984) to yield the titled compound (1.44g); ¹H NMR (CDCl₃) 1.25 (3H, t), 4.3 (2H, q), 4.55 (2H, s), 9.45 (1H, s), contains 20% starting material by NMR.

15 Ethyl-α-oximino-2-methylthiazole-4-acetate

This was prepared from ethyl-γ-chloro-α-oximinoacetoacetate (1.44g) using the method of Hatanaka et al. (*Journal of Medicinal Chemistry*, 1973, 16(9), 978-984) to yield the titled compound (0.64g); ¹H NMR (CDCl₃) 1.35 (3H, t), 2.7

20 (3H, s), 4.35 (2H, q), 8.2 (1H, s).

D,L-(2-methylthiazol-4-yl)glycine ethyl ester

This was prepared from ethyl-α-oximino-2-methylthiazole-4-acetate (0.62g) using the method of Hatanaka et al. (*Journal*
25 *of Medicinal Chemistry*, 1973, 16(9), 978-984) to yield the titled compound (0.40g); ¹H NMR (CDCl₃) 1.15 (3H, t), 1.95 (2H, br.), 2.6 (3H, s), 4.15 (2H, m), 4.65 (1H, s), 6.95 (1H, s).

N-Boc-D,L-(2-methylthiazol-4-yl)glycine ethyl ester

To a solution of D,L-(2-methylthiazol-4-yl)glycine ethyl ester (0.397g, 1.982 mmol) in tetrahydrofuran (20 cm³), was added di-tert-butylidicarbonate (0.475g, 2.180 mmol) and triethylamine (0.304 cm³, 2.180 mmol). This was allowed to stir for 1 hour and the solution concentrated in vacuo. The oil was taken up in ethyl acetate (c.a. 50 cm³) washed with 0.5% hydrochloric acid solution (c.a. 20 cm³), and saturated sodium bicarbonate solution (c.a. 20 cm³). This was then dried over magnesium sulphate and concentrated in vacuo to yield a yellow oil (0.654g, 2.177 mmol) [-100% yield]; ¹H NMR (CDCl₃) 1.1 (3H, s), 1.35 (9H, s), 2.6 (3H, s), 4.15 (3H, m), 5.3 (1H, d), 5.7 (1H, s), 7.0 (1H, s).

N-Boc-D,L-(2-methylthiazol-4-yl)glycine

To a solution of N-Boc-D,L-(2-methylthiazol-4-yl)glycine ethyl ester (0.595g, 1.982 mmol) in methanol (c.a. 15 cm³), was added 2M sodium hydroxide (1.98 cm³, 3.964 mmol), and allowed to stir for 30 minutes. The solution was concentrated in vacuo and taken up in water (c.a. 50 cm³). The aqueous solution was washed with ethyl acetate (c.a. 30 cm³), and then acidified to pH 2 with 5% hydrochloric acid solution (c.a. 50 cm³). The product was extracted with ethyl acetate (c.a. 3x60 cm³), dried over magnesium sulphate, and concentrated in vacuo to yield a pale yellow oil (0.645g, 2.368 mmol) [-100% yield]; ¹H NMR (CDCl₃) 1.35 (9H, s), 2.6 (3H, s), 5.4 (1H, d), 5.9 (1H, s), 7.1 (1H, s).

1-(N-Boc-D,L-(2-methylthiazol-4-yl)glycinyl) 1'-methyl-4,4'-

bispiperidine

Prepared by coupling N-Boc-D,L-(2-methylthiazol-4-yl)-glycine to 4,4'-(1'-methylbispiperidine) di-HCl salt using EDC and HOAt as described previously; ¹H NMR (CDCl₃) 0.5-1.3

5 (10H, br.), 1.35 (9H, s), 1.4-1.85 (6H, br.), 2.2 (3H, d), 2.6 (3H, s), 3.75-4.0 (1H, br.), 4.55 (1H, br.), 5.7 (1H, d), 6.1 (1H, d), 6.95 (1H, d)

1-(D,L-(2-Methylthiazol-4-yl)glyciny)- 1'-methyl-4,4'-

10 **bispiperidine**

Prepared from 1-(N-Boc-D,L-(2-methylthiazol-4-yl)glyciny)- 1'-methyl-4,4'- bispiperidine using DCM/TFA deprotection as described previously; ¹H NMR (CDCl₃) 0.9-1.8 (10H, br.), 2.1-2.3 (2H, br.), 2.45 (3H, br.), 2.6 (3H, s), 3.1-3.4 (3H,

15 br.), 4.6 (1H, br.), 4.95 (1H, s), 6.85 (1H, d).

1-(3-Chloroindole-6-carbonyl- D,L-(2-Methylthiazol-4-yl)glyciny)- 1'-methyl-4,4'-bispiperidine

Prepared by coupling 1-(D,L-(2-methylthiazol-4-yl)-

20 glyciny)- 1'-methyl-4,4'-bispiperidine to 3-chloroindole-6-carboxylic acid using EDC and HOAt as described previously;

¹H NMR (CDCl₃) 0.5-1.9 (12H, br.), 2.4 (2H, br.), 2.55 (3H, s), 2.65 (3H, s), 3.5 (2H, br.), 4.1 (1H, br.), 4.55 (1H, br.), 6.15 (1H, d), 7.15 (1H, d), 7.5 (2H, br.), 7.8-8.1

25 (2H, br.), 8.9-9.25 (1H, br.), 12.2-12.6 (1H, br. d); HPLC (Luna C18, Gradient3) rt 8.75min; LCMS M+1 514.

Example 159.

1-(3-Chloroindole-6-carbonyl-D,L-4-thiazolyglyciny)- 1'-

methyl-4,4'-bispiperidine**Ethyl- α -oximino-thiazole-4-acetate**

To a 2 necked r.b. flask (100 cm³) with ethanol thermometer,
5 concentrated sulphuric acid (25 cm³) was added and cooled to
0°C with stirring. To this solution, was added the ethyl- α -
oximino-2-aminothiazole-4-acetate (5.00g, 23.231 mmol).
Water (10 cm³) was then added and cooled to -10°C. A
solution of sodium nitrite (1.683g, 24.393 mmol) in water (5
10 cm³) was then added slowly over an hour keeping the
temperature below -5°C.

To a separate r.b. flask (500 cm³), water (180 cm³) was added
and cooled to 3°C. The reaction solution was poured on to
the cold water with stirring and then cooled to -5°C. To
15 this solution, 50% hypophosphoric acid (90 cm³) was added
dropwise over 10 minutes keeping the temperature at -5°C.
The solution was allowed to warm to room temperature and
stirred overnight. The product was extracted with diethyl
ether (c.a. 3x150 cm³) and washed with water. The ether
20 layer was concentrated in vacuo and treated to flash
chromatography (50% ethyl acetate/n-hexane) to yield a
orange oil upon concentration in vacuo (0.60g, 3.00 mmol)
[13% yield]; ¹H NMR (CDCl₃) 1.35 (3H, m), 4.35 (2H, m), 8.4
(1H, s), 8.9 (1H, s), 14.4 (1H, s).

25

D,L-4-thiazolyglycine ethyl ester

This was prepared from ethyl- α -oximino-thiazole-4-acetate
(0.60g) using the method of Hatanaka et al. (*Journal of*
Medicinal Chemistry, 1973, 16(9), 978-984) to yield the

titled compound (0.46g); ^1H NMR (CDCl_3) 1.25 (3H, t), 1.8-2.3 (2H, br.), 4.1 (2H, m), 4.75 (1H, s), 7.25 (1H, d), 8.7 (1H, d).

5 **N-Boc-D,L-4- thiazolyglycine ethyl ester**

To a solution of D,L-4-thiazolyglycine ethyl ester (0.460g, 2.470 mmol) in tetrahydrofuran (20 cm^3), was added di-tert-butylidicarbonate (0.530g, 2.470 mmol) and triethylamine (0.344 cm^3 , 2.470 mmol). This was allowed to stir for 10 hour and the solution concentrated in vacuo. The oil was taken up in ethyl acetate (c.a. 50 cm^3) washed with 0.5% hydrochloric acid solution (c.a. 20 cm^3), and saturated sodium bicarbonate solution (c.a. 20 cm^3). This was then dried over magnesium sulphate and concentrated in vacuo to yield an orange oil (0.709g, 2.477 mmol) [-100% yield]; ^1H NMR (CDCl_3) 1.15 (3H, t), 1.35 (9H, s), 4.1 (2H, m), 5.45 (1H, d), 5.75 (1H, d), 7.3 (1H, d), 8.7 (1H, d).

N-Boc-D,L-4- thiazolyglycine

20 To a solution of N-Boc-D,L-4- thiazolyglycine ethyl ester (0.700g, 2.470 mmol) in methanol (c.a. 15 cm^3), was added 2M sodium hydroxide (2.47 cm^3 , 4.940 mmol) and allowed to stir for 90 minutes. The solution was concentrated in vacuo and taken up in water (c.a. 20 cm^3). The aqueous solution was 25 washed with ethyl acetate (c.a. 20 cm^3), and then acidified to pH 2 with 5% hydrochloric acid solution (c.a. 50 cm^3). The product was extracted with ethyl acetate (c.a. 3x30 cm^3), dried over magnesium sulphate, and concentrated in vacuo to yield a pale yellow oil (0.582g, 2.254 mmol) [91% 30 yield]; ^1H NMR (CDCl_3) 1.35 (9H, s), 5.5 (1H, d), 5.8 (1H,

d), 7.35 (1H, d), 8.75 (1H, d), 9.8-10.2 (1H, br.).

1-(N-Boc-D,L-4-thiazolyglyciny)- 1'-methyl-4,4'-bispiperidine

- 5 Prepared by coupling N-Boc-D,L-4-thiazolyglycine to 4,4'-(1'-methylbispiperidine) di-HCl salt using EDC and HOAt as described previously; ¹H NMR (CDCl₃) 0.8-1.25 (10H, br.), 1.35 (9H, m), 1.7 (6H, br.), 2.0 (6H, m), 2.4 (3H, br.), 3.1 (2H, br.), 3.7 (1H, d), 4.6 (1H, d), 5.8 (1H, d), 10 6.0 (1H, br.), 7.25 (1H, 1H, br.), 8.65 (1H, m).

1-(D,L-4-Thiazolyglyciny)- 1'-methyl-4,4'-bispiperidine

- Prepared from 1-(N-Boc-D,L-4-thiazolyglyciny)- 1'-methyl-4,4'-bispiperidine using DCM/TFA deprotection as described 15 previously. The product was purified by prep HPLC; LCMS M+1 323.

1-(3-Chloroindole-6-carbonyl- D,L- thiazol-4-ylglyciny)- 1'-methyl-4,4'-bispiperidine

- 20 Prepared by coupling 1-(D,L-4-Thiazolyglyciny)- 1'-methyl-4,4'-bispiperidine to 3-chloroindole-6-carboxylic acid using EDC and HOAt as described previously; ¹H NMR (CD₃CN) 0.5-2.0 (10H, br.), 2.5 (2H, m), 2.8 (3H, br.), 3.1 (2H, m), 3.5 (2H, br.), 4.2 (1H, d), 4.6 (1H, d), 6.4 (1H, m), 7.5 25 (1H, br.), 7.8 (2H, br.), 8.15 (2H, br.), 9.05 (1H, br.), 9.9 (1H, br.); HPLC (Luna C18, Gradient3) rt 6.69min; LCMS M+1 500.

Preparation of starting materials:

Boc-R-4-(carboxymethyl)phenylglycine

5 **R-4-Hydroxyphenylglycine methyl ester hydrochloride.**

To a dry 250ml three necked round bottom flask, equipped with a low temperature thermometer, a septum for nitrogen coverage and another for introduction of thionyl chloride by syringe, was added R-4-hydroxyphenylglycine (12.5g) and dry
10 methanol (24ml). The mixture was stirred (magnetic stirrer) and cooled to an internal temperature of -20°C using cardice/acetone. Using a syringe, thionyl chloride was added dropwise to the cooled mixture over a period of 10min.

(Care: the reaction of thionyl chloride with methanol is
15 very exothermic and rate of addition should be such that the thionyl chloride is efficiently stirred into the mixture and that the temperature does not rise above -20°C. Once the addition was complete the mixture was allowed to warm to room temperature overnight (16-18hr). Dry ether (150ml) was
20 added and the white ppt. that formed was filtered off, washed with a little more ether and dried. Yield 15.5g 95%..
Nmr.

Boc-R-4-Hydroxyphenylglycine methyl ester hydrochloride

25 To a stirred mixture of R-4-hydroxyphenylglycine methyl ester hydrochloride 14g and sodium bicarbonate 11.7g in tetrahydrofuran (THF) 150ml and water 50ml, was added in one portion, di- t-butyl dicarbonate 15.9g. The mixture was stirred rapidly to allow thorough mixing for 4h. Hexane
30 (75ml) was added and the organic layer separated and washed

with sat. sodium bicarbonate solution, then brine and then dried with magnesium sulphate. The drying agents was filtered off and washed with a little THF and evaporated to dryness, finishing with a high vacuum pump to remove the
5 last traces of di- t-butyl dicarbonate. Yield 19.7g 96%.

— Nmr.

Boc-R-4-(trifluoromethanesulphonyloxy)phenylglycine methyl ester hydrochloride

10 To a stirred solution of Boc-R-4-hydroxyphenylglycine methyl ester 19g in dichloromethane 400ml was added 2,6-lutidine 9.44ml and 4-dimethylaminopyridine 1.65g and the mixture cooled in an ice bath. Trifluoromethane-sulphonic anhydride 13.74ml was added over a period of 5min and then
15 the reaction left to warm to room temperature over 4h. The organic solution was washed with water, 2 x 150ml, 1N HCl 2 x 150ml and the saturated sodium bicarbonate 150ml. The organics were dried with magnesium sulphate and then evaporated to and oil. The mixture was purified using flash
20 chromatography (SiO₂ 250g eluting with 1:1 hexane/dichloromethane and then neat dichloromethane). Pure product fractions were combined and evaporated, finishing with a high vacuum pump to remove all traces of solvent, to give a white solid, 19g 77%. Nmr.

25

Boc-R-4-(carboxymethyl)phenylglycine methyl ester.

Boc-R-4-trifluoromethanesulphonyloxyphenylglycine methyl ester (15g), methanol (32.6ml), bis-1,3-diphenylphosphinylpropane (448mg), palladium (II) acetate
30 (255mg), triethylamine (10.2ml) and dimethylformamide (72ml)

were placed in the glass liner of the Parr reactor and the reactor assembled. The vessel was pressurised to ~10psi with nitrogen and the gas released (repeated five times to remove all oxygen from the system). Carbon monoxide gas was then
5 carefully introduced (use extreme care -the gas cylinder is pressurised to far beyond the bursting disc pressure of the Parr, ideally use a pressure regulator to reduce the pressure to ~100psi) to ~20psi and released three times (into the back of a fume hood). Carbon monoxide was then
10 added to ~100psi and the stirrer started. The vessel was slowly heated to 65°C internal temperature and then stirred at 65°C overnight. (At the early stages more carbon monoxide was added to maintain ~100psi) A sample was removed after 18h and examined by tlc. When complete, the reaction was
15 cooled to ~30°C, the gas released and the vessel flushed five times with nitrogen as before. The reaction mixture was partitioned between ethyl acetate and water and the organic layer washed with 1M hydrochloric acid and then saturated sodium bicarbonate. The solution was dried with MgSO₄ and
20 evaporated. Flash chromatography of the resulting oil gave the product, pure by tlc, 10.6g 90%. Nmr

Boc-R-4-(carboxymethyl)phenylglycine.

To a solution of Boc-R-4-carboxymethylphenylglycine methyl
25 ester 692mg in THF 10ml was added a solution of lithium hydroxide hydrate 90mg in water 7ml. The mixture immediately became cloudy and over 15min cleared. After 30min, tlc showed the reaction to be complete. Ethyl acetate 20ml and water 20ml were added and the aqueous layer separated. The
30 aqueous solution was acidified with 2M hydrochloric acid and extracted with ethyl acetate (3 x 20ml). The organic

solution was then washed with water x 2 and brine x 2, dried with MgSO_4 and evaporated to give the mono-ester (650mg, 98%), pure by tlc. Nmr.

5 Boc-R-4-(carboxybenzyl)phenylglycine methyl ester

By the same method as described above, using 27.6g of Boc-R-4-trifluoromethanesulphonyloxyphenylglycine methyl ester and benzyl alcohol to give the Boc-D-4-

(carboxybenzyl)phenylglycine methyl ester 18.7g pure, 70% plus a further 6g of impure material (the major contaminant is benzyl alcohol). Nmr

Boc-R-4-(carboxamido)phenylglycine methyl ester

15 Boc-R-4-(carboxy)phenylglycine methyl ester

Boc-R-4-(carboxybenzyl)phenylglycine methyl ester (500mg) was dissolved in THF containing Pd/C 10% (100mg) and hydrogenated at 1atm for 2h. Removal of the catalyst by filtration and evaporation of solvent gave Boc-R-4-

20 (carboxy)phenylglycine methyl ester (330mg, 87%).

Nmr.

Boc-R-4-(carboxamido)phenylglycine methyl ester

To a solution of Boc-R-4-(carboxy)phenylglycine methyl ester (3.5g) in DMF 30ml was added EDCI (2.60g 1.36mmol) and HOBT (1.4g 10.4mmol) and the mixture stirred for 10min before cooling in a ice bath and bubbling in ammonia gas for 5min. The mixture was stirred for 2h at room temperature and then diluted with ethyl acetate and washed with water. The

aqueous solution was extracted with a little ethyl acetate and the combined organics washed with brine. The organic solution was evaporated to an oil which was purified by flash chromatography (SiO_2 - dichloromethane/ ethyl acetate 5 0 - 25%) to give Boc-R-4-(carboxamido)phenylglycine methyl ester (1.7g 48%). Nmr.

Boc-R-4-(methylcarboxamido)phenylglycine methyl ester

Was prepared by a similar method to that descibed above.

10 Nmr

Boc-R-4-Methoxyphenylglycine.

Boc-R-4-hydroxyphenylglycine methyl ester was converted to Boc-R-4-methoxyphenylglycine using the alkylation method 15 described by Basak et al. (Tetrahedron Lett. 1998, 39 (27), 4883-4886) followed by hydrolysis of the methyl ester with lithium hydroxide in aqueous THF. Nmr

Boc-D,L-2-chlorophenylglycine

20 2-Chlorobenzaldehyde (20mmol., 2.252ml) and 2,4 dimethoxybenzylamine (20mmol., 3.004ml) were added together and stirred for 2 hours. DCM (5ml) was added and any water separated and removed. tert-Butyl isonitrile (20mmol., 2.262ml) was added and stirred for 10mins followed by acetic 25 acid (20mmol., 1.145ml). Stirring was continued for 3 days. The reaction mixture was then treated with TFA (30ml) and triethylsilane (5ml). After 3 hours the mixture was evaporated to dryness, 6M HCl (100ml) added and the whole refluxed overnight at 130°C, stirring rapidly. The mixture

was allowed to cool and extracted with EtOAc (50ml x2) the aqueous fraction was evaporated to dryness and treated with 2M NaOH solution. The mixture was extracted with EtOAc (50ml x2) excess boc anhydride (5.2g) in dioxan (20ml) was added to the aqueous fraction and stirred overnight. The mixture was extracted with diethyl ether (100ml x2) acidified to pH 1 (CHCl₃) and extracted with EtOAc (50ml x2). The combined organic fractions were washed with water and evaporated to dryness under high vacuo. The product Boc-2-chlorophenylglycine (4.252g, 74.5%)

¹H nmr (CD₃CN/D₂O) 7.3 (4H, m); 5.5 (1H, s); 1.3 (9H, s). MS 286 (M+1)

By a similar method the following amino acids were obtained

15

Boc-D,L-3-fluorophenylglycine

¹H nmr (CD₃CN/D₂O) 7.3 (1H, m), 7.1 (3H, m); 5.2 (1H, s); 1.3 (9H, s). MS 270 (M+1)

20 **Boc-D,L-4-fluorophenylglycine**

¹H nmr (CD₃CN/D₂O) 7.3 (2H, m); 6.9 (2H, m), 5.0 (1H, s); 1.3 (9H, s). MS 270 (M+1)

Boc-D,L-2-methylphenylglycine

25 ¹H nmr (CD₃CN/D₂O) 7.3 (4H, m); 5.5 (1H, s); 2.5 (3H, s); 1.3 (9H, s). MS 266 (M+1)

Boc-D,L-3-thienylglycine

¹H nmr (CD₃CN/D₂O) 7.5 (2H, m); 7.1 (1H, d); 5.3 (1H, s);
1.3 (9H, s). MS 258 (M+1)

5 Boc-D,L-2-fluorophenylglycine

Was obtained by treating D,L-2-fluorophenylglycine (Aldrich) with Boc anhydride (1.1eq) and 2M NaOH (1eq) in Ethanol. Aqueous work up as described above yielded the protected amino acid.

10 Nmr.

These protected aminoacids were then coupled with first an amine and then, after removal of the Boc protecting group, with a carboxylic acid by method 2 to give the following

15 inhibitor examples:

Example 160.

1-(4 Methoxybenzoyl-D,L-3-thienylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

20 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.18
LCMS M+1 514. Nmr.

Example 161.

1-(Indol-6-carbonyl-D,L-3-thienylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

25

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.44
LCMS M+1 523. Nmr.

Example 162.

1-(4 Methoxybenzoyl-D,L-3-fluorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

5 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.61

LCMS M+1 526. Nmr.

Example 163.

10 1-(Indol-6-carbonyl-D,L-3-fluorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.88

LCMS M+1 535. Nmr.

Example 164.

15 1-(4 Methoxybenzoyl-D,L-4-fluorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.52

LCMS M+1 526. Nmr.

20 Example 165.

1-(Indol-6-carbonyl-D,L-4-fluorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.92

LCMS M+1 535. Nmr.

Example 166.

1-(4 Methoxybenzoyl-D,L-2-chlorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.82

5 LCMS M+1 542 Nmr.

Example 167.

1-(Indol-6-carbonyl-D,L-2-chlorophenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

10 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.63

LCMS M+1 551 Nmr.

Example 168.

1-(4 Methoxybenzoyl-D,L-2-methylphenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

15

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.69

LCMS M+1 522 Nmr.

Example 169.

1-(Indol-6-carbonyl-D,L-2-methylphenylglyciny1) 4-(2-methylsulfonylphenyl)-piperazine

20

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA) rt 10.76

LCMS M+1 531 Nmr.

Example 170.

1-(Indol-6-carbonyl-D-2-fluorophenylglyciny) 4-(4-fluoro -
2-methylsulfonylphenyl)-piperazine

Hplc (Luna 2 C18 3u water/acetonitrile/TFA, gradient = 5-
5 100%MeCN over 7 min)rt 10.92

LCMS M+1 553 Nmr.

Example 171.

1-(Indol-6-carbonyl-D-(4-carboxyphenylglyciny)-(4-(1-
10 methylpiperidin-4-yl)piperazine)

By coupling of Boc-D-4-carboxymethylphenylglycine with 1-(4-
(1-methylpiperidin-4-yl)piperazine) using HOAt and EDCI,
followed by deprotection (TFA), coupling to indol-6-
carboxylic acid using HOAt and EDCI followed by hydrolysis
15 of the methyl ester with lithium hydroxide.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
6.05min

LCMS M+1 504

Nmr.

20

Example 172.

1-(Indol-6-carbonyl-D-phenylglyciny)-4-(4-
hydroxyphenyl)piperazine

By coupling of Boc-D-phenylglycine with 1-(4-
25 hydroxyphenyl)piperazine using HOAt and EDCI, followed by
deprotection (TFA) and coupling to indol-6-carboxylic acid
using HOAt and EDCI.

Hplc (Symmetry C8, Gradient3, water/acetonitrile/TFA), rt,

6.0min

LCMS M+1 455

Nmr.

5 Example 173.

1-(3-Chloroindol-6-carbonyl-D-phenylglyciny)-4-(4-hydroxyphenyl)piperazine

By coupling of Boc-D-phenylglycine with 1-(4-hydroxyphenyl)piperazine using HOAt and EDCI, followed by
10 deprotection (TFA) and coupling to 3-chloroindol-6-carboxylic acid using HOAt and EDCI.

Hplc (Symmetry C8, Gradient3, water/acetonitrile/TFA), rt,
6.55min

LCMS M+1 489

15 Nmr.

Example 174.

1-(4-methoxybenzoyl-D-4-methoxyphenylglyciny)-4-(2-methylsulphonylphenyl)piperazine

20 By coupling of Boc-D-4-methoxyphenylglycine with-(2-methylsulphonylphenyl)piperazine using HOAt and EDCI, followed by deprotection (TFA) and coupling to 4-methoxybenzoic acid using HOAt and EDCI.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
25 10.4min

LCMS M+1 538

Nmr.

Example 175.

1-(5-Fluoroindole-6-carbonyl-D-phenylglyciny)-1-methyl-4,4'-bispiperidine.

5

N-(2,2-Dimethoxyethyl)-4-fluoro-3-methoxyaniline

To a solution of 4-fluoro-3-methoxyaniline (0.98g 6.9mmol) in ethanol (20ml) was added glyoxal 1,1-dimethyl acetal (0.89g 8.27mmol). Pd/C 5% (50mg) was added and the mixture
10 hydrogenated. Removal of the catalyst by filtration and evaporation of solvent in vacuo gave N-(2,2-dimethoxyethyl)-4-fluoro-3-methoxyaniline 1.6g

NMR LCMS M+1 (less MeO) 199

15 N-(2,2-Dimethoxyethyl)-N-methanesulphonyl-4-fluoro-3-methoxyaniline

N-(2,2-dimethoxyethyl)-4-fluoro-3-methoxyaniline (1.46g, 6.37mmol) in dichloromethane (20ml) was treated with pyridine (0.5g 6.37mmol) and methanesulphonyl chloride
20 (766mg, 6.69mmol) and the mixture stirred until the reaction was complete by tlc. Aqueous work up and removal of solvent in vacuo gave N-(2,2-dimethoxyethyl)-N-methanesulphonyl-4-fluoro-3-methoxyaniline 1.91g

NMR

25

5-Fluoro-1-methanesulphonyl-6-methoxyindole

To a solution of N-(2,2-dimethoxyethyl)-N-methanesulphonyl-4-fluoro-3-methoxyaniline (1.91g, 0.65mmol) in dry toluene

at 0°C under argon, was added slowly a solution of TiCl_4 (0.173g, 0.911mmol) in dry toluene (10ml). The solution was then heated to 70°C for 1h. cooled and poured onto ice/sat. sod. bicarbonate solution (20ml). The organic layer was separated, washed with sat. sod. bicarbonate solution, 0.5% hydrochloric acid (2 x 20ml) and water (2 x 20ml). The solution was dried (MgSO_4) and evaporated in vacuo to give 5-fluoro-1-methanesulphonyl-6-methoxyindole ((0.102g)

NMR

10

5-Fluoro-6-hydroxy-1-methanesulphonylindole

To a solution of 5-fluoro-1-methanesulphonyl-6-methoxyindole (0.10g 0.41mmol) in dry dichloromethane (3ml) at -10°C was added a solution of BBr_3 (1M in dichloromethane, 1.23ml) over one minute. The reacture was warmed to room temperature and stirred for 2h and then poured onto ice/1M hydrochloric acid (10ml). After stirring for 15min the mixture was extracted with ethyl acetate (1 x 50ml, 2x 20ml), dried (MgSO_4) and evaporated in vacuo to give 5-fluoro-6-hydroxy-1-methanesulphonylindole (70mg)

NMR

5-Fluoro-1-methanesulphonyl-6-trifluoromethanesulphonyloxy-indole

To a solution of 5-fluoro-6-hydroxy-1-methanesulphonylindole (0.57mg, 2.49mmol) in dry dichloromethane (20ml) at 0°C was added pyridine (0.24ml, 2.99mmol) and then trifluoromethanesulphonic anhydride (0.50ml, 2.99mmol) and the mixture stirred for 2h. The reaction mixture was washed with 0.5% hydrochloric acid (2 x 50ml), sodium bicarbonate

solution (2 x 50ml) and water (50ml), dried (MgSO_4) and filtered through a short pad of silica. Evaporation of solvent in vacuo gave 5-fluoro-1-methanesulphonyl-6-trifluoromethanesulphonyloxy-indole, (0.67g).

5 NMR

Methyl 5-fluoro-1-methanesulphonyl-indol-6-carboxylate,

To a solution of 5-fluoro-1-methanesulphonyl-6-trifluoromethanesulphonyloxy-indole, (0.70g 1.94mmol) was added, Pd (II) acetate (14mg), bis 1,3-diphenylphosphinylpropane (24mg), dimethylformamide (4ml) and methanol (2ml) and triethylamine (0.54ml) and the mixture stirred for 2 min. Carbon monoxide gas was bubbled in for 15min and then the mixture was heated to 75°C under an atmosphere of carbon monoxide and stirred overnight. After cooling to room temperature the mixture was poured into ethyl acetate (80ml) and washed with 1M hydrochloric acid (50ml), sat. sod. bicarbonate (50ml) and water (50ml). Drying (MgSO_4), evaporation of solvent gave crude product (0.53g). Purification of a portion (225mg) by flash chromatography (SiO_2 , 25% ethyl acetate in hexane) gave methyl 5-fluoro-1-methanesulphonyl-indol-6-carboxylate, (173mg)

NMR

25

5-fluoro-1-methanesulphonyl-indol-6-carboxylic acid

To a solution of methyl 5-fluoro-1-methanesulphonyl-indol-6-carboxylate (173mg) in THF (15ml) and water (2ml) was added 2M lithium hydroxide solution (3 equiv) and the mixture heated to 50°C for 2h. and then allowed to cool overnight.

The solution was concentrated in vacuo, diluted with 2M sodium hydroxide solution (10ml) and washed with ethyl acetate. The aqueous solution was acidified to pH3 with conc. hydrochloric acid and extracted with ethyl acetate (3 x 30ml). The organic solution was evaporated in vacuo to give 5-fluoro-1-methanesulphonyl-indol-6-carboxylic acid (164mg) - (circa 80% pure)

NMR

10 1-(5-fluoro-1-methanesulphonyl-indol-6-carbonyl-D-phenylglyciny-4,4'-(1'-methylbispiperidine)

5-fluoro-1-methanesulphonyl-indol-6-carboxylic acid (164mg) was coupled to D-phenylglyciny-4,4'-(1'-methylbispiperidine) using EDCI/HOAt as previously described to give 1-(5-fluoro-1-methanesulphonyl-indol-6-carbonyl-D-phenylglyciny-4,4'-(1'-methylbispiperidine) (111mg) - (~70% pure)

NMR

20 1-(5-fluoroindol-6-carbonyl-D-phenylglyciny-4,4'-(1'-methylbispiperidine)

1-(5-fluoro-1-methanesulphonyl-indol-6-carbonyl-D-phenylglyciny-4,4'-(1'-methylbispiperidine) (111mg--70% pure) was refluxed in ethanol (5ml) and sodium hydroxide solution (34mg in 0.34ml) for 2.25h. The mixture was evaporated to dryness, taken up in water (10ml) and extracted with chloroform (60ml). The organic solution was dried (MgSO₄) and evaporated in vacuo and the residue purified by Prep. Hplc. To give 1-(5-fluoroindol-6-carbonyl-D-phenylglyciny-4,4'-(1'-methylbispiperidine) (19mg)

Hplc (Luna C18 Gradient 3) rt 11.37min

LCMS M+1 477

NMR

5 Example 176.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide

1-t-Butoxycarbonyl-4-(2-pyridoxy)piperidine

- 10 1-t-Butoxycarbonyl-4-piperidinol (5.0g 24.88mmol) in dry dimethylformamide (60ml) was treated with sodium hydride (60% 2.99g 74.75mmol) at room temperature under argon and then with 2-chloropyridine hydrochloride (4.1g 27.33mmol). Then mixture was heated at 80°C overnight. After cooling the
- 15 reaction was carefully quenched with water (5ml) and then diluted with more water (20ml) and extracted with ethyl acetate (30ml). The organic solution was washed with sat. sodium bicarbonate, dried (MgSO₄) and evaporated to give 1-t-butoxycarbonyl-4-(2-pyridoxy)piperidine (4.96g 72%).

20

4-(2-pyridoxy)piperidine dihydrochloride.

- 1-t-Butoxycarbonyl-4-(2-pyridoxy)piperidine (6.5g) was treated with a solution of hydrogen chloride in ethyl acetate (110ml) for 7h and the mixture evaporated to give 4-
- 25 (2-pyridoxy)piperidine dihydrochloride, (7.4g 90%)

1-(Benzyoxycarbonyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide

Benzyloxycarbonyl-D-phenylglycine (3.75g 13.14mmol) was coupled to 4-(2-pyridoxy)piperidine dihydrochloride (3.0g 11.94mmol) using EDCI (2.52g 13.14g), HOAt (1.79g 13.13mmol) and triethylamine (3.63g 35.87mmol) to give, after work up with ethyl acetate and sodium bicarbonate solution, 1-(benzyloxycarbonyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide, (4.9g 92%)

1-D-phenylglyciny-4-(2-pyridoxy)piperidinamide

10 1-(Benzyloxycarbonyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide (400mg) was hydrogenated in ethanol with 5% Pd/C overnight. Removal of catalyst and evaporation of solvent gave 1-D-phenylglyciny-4-(2-pyridoxy)piperidinamide (162mg 58%)

15

Using a similar method and the appropriate starting materials the following intermediates were also prepared:

1-(D-phenylglyciny-4-(4-pyridoxy)piperidinamide

20 1-(D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide

1-(D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide

25 1-D-phenylglyciny-4-(2-pyridoxy)piperidinamide (162mg 0.52mmol) was treated with triethylamine (58mg 0.573mmol) and p-anisoyl chloride (93mg 0.545mmol) in dry dichloromethane for 1h. The reaction mixture was washed with sodium bicarbonate solution and brine, dried (MgSO₄)

and evaporated to an oil. Flash chromatography gave the product 1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide, (60mg 26%)

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,

5 8.94min

LCMS M+Na 468

Nmr

By a similar method the following compounds were prepared:

10 Example 177.

1-(Indol-6-carboxyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide

By the coupling of indol-6-carboxylic acid and 1-D-phenylglyciny-4-(2-pyridoxy)piperidinamide using EDCI and HOAt.

15 LCMS M+1 455

Nmr

20 Example 178.

1-(3-Chloroindol-6-carboxyl-D-phenylglyciny)-4-(2-pyridoxy)piperidinamide

By the coupling of 3-chloroindol-6-carboxylic acid and 1-D-phenylglyciny-4-(2-pyridoxy)piperidinamide using EDCI and HOAt.

25 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
10.29min

LCMS M+1 489

Nmr

Example 179.

- 5 1-(3-Chloroindol-6-carbonyl-D-phenylglyciny1)-4-(4-pyridoxy)piperidinamide

By the coupling of 3-chloroindol-6-carboxylic acid and 1-D-phenylglyciny1-4-(4-pyridoxy)piperidinamide using EDCI and HOAt.

10 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,

8.16min

LCMS M+1 489

Nmr

15 Example 180.

- 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-pyridoxy)piperidinamide

By the coupling of p-anisoyl chloride and 1-D-phenylglyciny1-4-(4-pyridoxy)piperidinamide in dichloromethane

- 20 with triethylamine

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
7.0min

LCMS M+1 446

Nmr

25

Exempl 181.

- 1-(Indol-6-carbonyl-D-phenylglyciny1)-4-(4-

pyridoxy)piperidinamide

By the coupling of indol-6-carboxylic acid and 1-D-phenylglyciny-4-(4-pyridoxy)piperidinamide with EDCI and HOAt.

5 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
7.08min

LCMS M+1 455

Nmr

10 **Example 182**

1-(4-Methoxybenzoyl-D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide

By the coupling of p-anisoyl chloride and 1-(D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide in
15 dichloromethane with triethylamine

LCMS M+1 432

Nmr

Example 183.

20 1-(Indol-6-carbonyl-D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide

By the coupling indol-6-carboxylic acid and 1-(D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide with EDCI and HOAt

25 LCMS M+1 441

Nmr

Example 184.

1-(3-Chloroindol-6-carbonyl-D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide

By the coupling 3-chloroindol-6-carboxylic acid and 1-(D-phenylglyciny)-3-R,S-(4-pyridoxy)pyrrolidinamide with EDCI and HOAt

LCMS M+1 475

Nmr

10 Example 185.

1-(4-Methoxybenzoyl-D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide

By the coupling of p-anisoyl chloride and 1-(D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide in dichloromethane with triethylamine

LCMS M+1 432

Nmr

Example 186.

20 1-(3-Chloroindol-6-carbonyl-D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide

By the coupling 3-chloroindol-6-carboxylic acid and 1-(D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide with EDCI and HOAt

25 LCMS M+1 475

Nmr

Example 187.

1-(Indol-6-carbonyl-D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide

By the coupling indol-6-carboxylic acid and 1-(D-phenylglyciny)-3-R,S-(2-pyridoxy)pyrrolidinamide with EDCI and HOAt

LCMS M+1 441

Nmr

10 Example 188.

1-(4-methoxybenzoyl-D-4-hydroxyphenylglyciny)-4-(2-methylsulphonylphenyl)piperazine

By coupling of Boc-D-4-hydroxyphenylglycine with-(2-methylsulphonylphenyl)piperazine using HOAt and EDCI, followed by deprotection (TFA) and coupling to 4-methoxybenzoic acid using HOAt and EDCI.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt, 9.1min

LCMS M+1 524

20 Nmr.

Example 189.

1-(Indol-6-carbonyl-D-4-hydroxyphenylglyciny)-4-(2-methylsulphonylphenyl)piperazine

25 By coupling of Boc-D-4-hydroxyphenylglycine with-(2-methylsulphonylphenyl)piperazine using HOAt and EDCI, followed by deprotection (TFA) and coupling to 6-indole carboxylic acid using HOAt and EDCI.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
9.0min

LCMS M+1 533

Nmr.

5

Example 190.

1-(Indol-6-carbonyl-D-4-hydroxyphenylglyciny1)-1'-methyl-
4,4'-bispiperidine

By coupling of Boc-D-4-hydroxyphenylglycine with 4,4'-(1'-
methylbispiperidine) di-HCl salt using HOAt and EDCI,
followed by deprotection (TFA) and coupling to 6-indole
carboxylic acid using HOAt and EDCI.

Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
6.8min

15 LCMS M+1 475

Nmr.

Example 191.

1-(3-Chloroindol-6-carbonyl-D-4-hydroxyphenylglyciny1)-1'-
methyl-4,4'-bispiperidine

By coupling of Boc-D-4-hydroxyphenylglycine with 4,4'-(1'-
methylbispiperidine) di-HCl salt using HOAt and EDCI,
followed by deprotection (TFA) and coupling to 3-
chloroindole-6-carboxylic acid using HOAt and EDCI.

25 Hplc (Luna C18, Gradient3, water/acetonitrile/TFA), rt,
7.3min

LCMS M+1 509

Nmr.

In the following examples the following additional abbreviations and meanings are included: CI-MS, chemical ionization mass spectrum; DMSO, dimethyl sulfoxide (perdeuterated if for NMR); EtOAc, ethyl acetate; EtOH, ethanol; IS-MS, ion spray mass spectrum; RPHPLC, reverse phase HPLC; SCX, strong cation exchange resin; THF, tetrahydrofuran; TLC, thin layer chromatography with R_f as relative mobility;

Reagents were obtained from a variety of commercial sources.

IR means an infrared spectrum was obtained. ^1NMR , $^1\text{H-NMR}$, or $^1\text{H NMR}$ means a proton magnetic resonance spectrum was obtained.

In general in this specification, "D-" or "R-" in the name of a product indicates the product was made beginning with a chiral starting material, for example D-phenylglycine; however, racemization may have occurred, and the enantiomeric purity may not have been determined.

Examples 201-210

25 Preparation of Starting Materials

4-[(Benzyloxycarbonyl-D-phenylglyciny] aminomethyl]-1-Boc-piperidine

Using Coupling Method C, benzyloxycarbonyl-D-phenylglycine (10.4 g, 36.5 mmol) and 4-aminomethyl-1-Boc-piperidine (7.3 g, 36.5 mmol) afforded, after purification by column

chromatography (SiO₂: 4:1 to 3:2 hexanes:EtOAc), 10.2 g (58%) of the title compound.

¹NMR

IS-MS, m/e 482 (M+1).

5

4-[(D-Phenylglyciny]aminomethyl]-1-Boc-piperidine

(Deprotection Method C) A solution of 4-[(benzyloxycarbonyl-D-phenylglyciny]aminomethyl]-1-Boc-piperidine (9.00 g, 18.7 mmol) and 10% palladium on carbon (2.34 g) in EtOAc (80 mL):EtOH (200 mL) was placed under an atmosphere of hydrogen gas (balloon). After 16 h, the mixture was filtered and concentrated affording 6.31 g (98%) of the title compound, which was used without further purification.

¹NMR

15 IS-MS, m/e 348 (M+1).

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-Boc-piperidine

(Acylation Method C) A solution of 4-[(D-phenylglyciny]-aminomethyl]-1-Boc-piperidine (2.38 g, 6.88 mmol) and pyridine (8 mL) in methylene chloride was treated with 4-methoxybenzoyl chloride (1.76 g, 10.3 mmol) in methylene chloride (prepared by treatment of 4-methoxy benzoic acid with excess oxalyl chloride in methylene chloride followed by concentration). After 2 days, the mixture was partitioned between water and methylene chloride. The organic extracts were washed with 1 N HCl, water, 1 N NaOH and brine, and concentrated. The residue was purified by column chromatography (SiO₂: 1:1 to 1:3 hexanes:EtOAc), affording 2.33 g (71%) of the title compound.

¹NMR

IS-MS, m/e 482 (M+1)

Analysis for $C_{27}H_{35}N_3O_5$:

Calcd: C, 67.3; H, 7.3; N, 8.7;

Found: C, 67.4; H, 7.4; N, 8.7.

5 4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-
piperidine

Using Deprotection Method D, 4-[(4-methoxybenzoyl-D-
phenylglyciny]aminomethyl]-1-Boc-piperidine (2.38 g)
afforded 1.56 g (82%) of 4-[(4-methoxybenzoyl-D-phenyl-
10 glyciny]aminomethyl]piperidine.

^1NMR

IS-MS, m/e 382 (M+1)

General Procedure: Unless otherwise indicated, the product
15 of Examples 201-210 was prepared from 4-[(4-methoxybenzoyl-
D-phenylglyciny]aminomethyl]piperidine and the indicated
aldehyde or ketone using Alkylation Method D.

Example 201.

20 4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-
isopropylpiperidine

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine
(0.10 g, 0.26 mmol) and acetone afforded 89 mg (81%) of the
title compound.

25 ^1NMR

IS-MS, m/e 424 (M+1)

Analysis for $C_{25}H_{33}N_3O_3$:

Calcd: C, 70.9; H, 7.9; N, 9.9;

Found: C, 70.8; H, 7.8; N, 9.9.

Example 202.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-(3-pentyl)piperidine

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine
5 (0.10 g, 0.26 mmol) and 3-pentanone afforded 57 mg (49%) of the title compound.

¹NMR

IS-MS, m/e 452 (M+1)

10 Example 203.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-(2-indanyl)piperidine

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine
15 (0.10 g, 0.26 mmol) and 2-indanone afforded 91 mg (78%) of the title compound.

¹NMR

IS-MS, m/e 498 (M+1)

Analysis for C₂₅H₃₃N₃O₃:

Calcd: C, 74.8; H, 7.1; N, 8.4;

20 Found: C, 74.5; H, 7.0; N, 7.9.

Example 204.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-cyclopentylpiperidine

25 4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine (0.10 g, 0.26 mmol) and cyclopentanone afforded 101 mg (86%) of the title compound.

¹NMR

IS-MS, m/e 450 (M+1)

Example 205.

4-[(4-Methoxybenzoyl-D-phenylglyciny] aminomethyl]-1-(cyclohexylmethyl)piperidine

- 5 4-[(4-Methoxybenzoyl-D-phenylglyciny] aminomethyl]piperidine
(0.10 g, 0.26 mmol) and cyclohexanecarboxaldehyde afforded
98 mg (79%) of the title compound.

¹NMR

IS-MS, m/e 478 (M+1)

10 Example 206.

4-[(4-Methoxybenzoyl-D-phenylglyciny] aminomethyl]-1-cyclohexylpiperidine

- 4-[(4-methoxybenzoyl-D-phenylglyciny] aminomethyl]piperidine
(0.10 g, 0.26 mmol) and cyclohexanone afforded 95 mg (79%)
15 of the title compound.

¹NMR

IS-MS, m/e 464 (M+1)

Example 207.

- 20 4-[(4-Methoxybenzoyl-D-phenylglyciny] aminomethyl]-1-(tetrahydropyran-4-yl)piperidine

4-[(4-methoxybenzoyl-D-phenylglyciny] aminomethyl]piperidine
(0.10 g, 0.26 mmol) and tetrahydro-4H-pyran-4-one afforded
78 mg (65%) of the title compound.

- 25 ¹NMR

IS-MS, m/e 466 (M+1)

Example 208.

- 30 4-[(4-Methoxybenzoyl-D-phenylglyciny] aminomethyl]-1-(tetrahydrothiopyran-4-yl)piperidine

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine (0.10 g, 0.26 mmol) and tetrahydro-4H-thiopyran-4-one afforded 63 mg (50%) of the title compound.

¹NMR

5 IS-MS, m/e 482 (M+1)

Example 209.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-methyl-piperidine

10 4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine (60 mg, 0.16 mmol) and paraformaldehyde afforded 59 mg (93%) of the title compound.

¹NMR

IS-MS, m/e 396 (M+1)

15

Example 210.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-ethyl-piperidine

20 4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]piperidine (60 mg, 0.16 mmol) and acetaldehyde afforded 23 mg (35%) of the title compound.

¹NMR

IS-MS, m/e 410 (M+1)

25 Examples 211-213

Preparation of Starting Materials

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-1-Boc-piperidine

30 Using Coupling Method C, 4-[(D-phenylglyciny]aminomethyl]-1-Boc-piperidine (2.5 g, 6.8 mmol) and indole-6-carboxylic acid (1.2 g, 7.6 mmol) afforded, after purification by

column chromatography (SiO₂: 2:3 hexanes:EtOAc to EtOAc),
2.57 g (83%) of the title compound.

¹NMR

IS-MS, m/e 491 (M+1)

5

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-
piperidine

Using Deprotection Method D, 4-[(indole-6-carbonyl-D-
phenylglyciny]aminomethyl]-1-Boc piperidine (1.6 g, 3.3
10 mmol) afforded 4-[(indole-6-carbonyl-D-phenylglyciny]-
aminomethyl]piperidine (1.27 g, 79%).

¹NMR

IS-MS, m/e 391 (M+1)

15 **General Procedure:** Unless otherwise indicated, the product
of Examples 211-213 was prepared from 4-[(indole-6-carbonyl-
D-phenylglyciny]aminomethyl]piperidine and the indicated
aldehyde or ketone using Alkylation Method D.

20 **Example 211.**

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-1-
isopropylpiperidine

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-
piperidine (0.10 g, 0.26 mmol) and acetone afforded 16 mg

25 (14%) of the title compound.

¹NMR

IS-MS, m/e 433 (M+1)

Example 212.

30 4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-1-
cyclopentylpiperidine

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-
piperidine (0.10 g, 0.26 mmol) and cyclohexanone afforded
19 mg (16%) of the title compound.

¹NMR

5 IS-MS, m/e 459 (M+1)

Example 213.

4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-1-
cyclohexylmethylpiperidine

10 4-[(Indole-6-carbonyl-D-phenylglyciny]aminomethyl]-
piperidine (0.10 g, 0.26 mmol) and cyclohexanecarboxaldehyde
afforded 14 mg (11%) of the title compound.

¹NMR

IS-MS, m/e 487 (M+1)

15

Examples 214-217

Preparation of Starting Materials

4-[(Benzyloxycarbonyl-D-phenylglyciny)]-1-Boc-piperidine

20 Using Coupling Method C, D-phenylglycine (6.10 g, 21.4 mmol)
and 4-amino-1-Boc-piperidine (4.27 g, 21.4 mmol) afforded,
after purification by column chromatography (SiO₂: 7:3
hexanes:EtOAc), 8.44 g (84%) of the title compound.

¹NMR

25 IS-MS, m/e 468 (M+1).

Analysis for C₂₆H₃₃N₃O₅:

Calcd: C, 66.3; H, 7.1; N, 9.0;

Found: C, 66.5; H, 7.1; N, 9.0.

30 4-[(D-Phenylglyciny]amino]-1-Boc-piperidine

Using Deprotection Method C, 4-[(benzyloxycarbonyl-D-
phenylglyciny]amino]-1-Boc-piperidine (8.0 g, 17 mmol)

afforded 6.1 g (90%) of the title compound, which was used without further purification.

¹NMR

IS-MS, m/e 334 (M+1).

5

4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]-1-Boc-piperidine

Using Acylation Method C, 4-[(D-phenylglyciny]amino]-1-Boc piperidine (2.23 g, 6.7 mmol) afforded, after purification
10 by column chromatography (SiO₂: 1:1 hexanes EtOAc), 2.44 g (78%) of the title compound.

¹NMR

IS-MS, m/e 468 (M+1).

15 4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]piperidine

Using Deprotection Method D, 4-[(4-methoxybenzoyl-D-phenylglyciny]amino]-1-Boc-piperidine (2.32 g) afforded
1.53 g (84%) of 4-[(4-methoxybenzoyl-D-phenylglyciny]-aminolpiperidine.

20 ¹NMR

IS-MS, m/e 368 (M+1).

General Procedure: Unless otherwise indicated, the product
of Examples 214-217 was prepared from 4-[(4-methoxybenzoyl-
25 D-phenylglyciny]amino]piperidine and the indicated aldehyde
or ketone using Alkylation Method D.

Example 214.

4-[(4-Methoxybenzoyl-D-phenylglyciny]aminomethyl]-1-
30 (3-pentyl)piperidine

4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]piperidine
(0.11 g, 0.3 mmol) and 3-pentanone afforded 81 mg (62%) of
the title compound.

¹NMR

5 IS-MS, m/e 438 (M+1).

Example 215.

4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]-1-(2-indanyl)-
piperidine

10 4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]piperidine
(0.11 g, 0.3 mmol) and 2-indanone afforded 121 mg (83%) of
the title compound.

¹NMR

IS-MS, m/e 484 (M+1).

15

Example 216.

4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]-1-cyclopentyl-
piperidine

20 4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]piperidine
(0.11 g, 0.3 mmol) and cyclopentanone afforded 103 mg (79%)
of the title compound.

¹NMR

IS-MS, m/e 436 (M+1).

25 Example 217.

4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]-1-cyclohexyl-
piperidine

30 4-[(4-Methoxybenzoyl-D-phenylglyciny]amino]piperidine
(0.11 g, 0.3 mmol) and 2-cyclohexanone afforded 112 mg (83%)
of the title compound.

¹NMR

IS-MS, m/e 450 (M+1).

Examples 218-220

Preparation of Starting Materials

5 4-[(Indole-6-carbonyl-D-phenylglyciny] amino]-1-Boc-piperidine

Using Acylation Method C, 4-[(D-phenylglyciny] amino]-1-Boc-piperidine (2.24 g, 6.15 mmol) and indole-6-carboxylic acid afforded 4-[(indole-6-carbonyl-D-phenylglyciny] amino]-1-Boc-piperidine (2.66 g, 82%).

¹NMR

IS-MS, m/e 477 (M+1).

15 4-[(Indole-6-carbonyl-D-phenylglyciny] amino]piperidine
Using Deprotection Method C, 4-[(indole-6-carbonyl-D-phenylglyciny] amino]-1-Boc-piperidine (1.2 g, 2.5 mmol) afforded 4-[(indole-6-carbonyl-D-phenylglyciny] amino]-piperidine (0.81 g, 83%).

¹NMR

20 IS-MS, m/e 377 (M+1).

General Procedure: Unless otherwise indicated, the product of Examples 218-220 was prepared from 4-[(indole-6-carbonyl-D-phenylglyciny] amino]piperidine and the indicated aldehyde
25 or ketone using Alkylation Method D.

Example 218.

4-[(Indole-6-carbonyl-D-phenylglyciny] amino]-1-isopropyl-piperidine

30 4-[(Indole-6-carbonyl-D-phenylglyciny] amino]piperidine (0.10 g, 0.27 mmol) and acetone afforded 21 mg (19%) of the title compound.

¹NMR

IS-MS, m/e 419 (M+1).

Example 219.

5 4-[(Indole-6-carbonyl-D-phenylglyciny]amino]-1-cyclopentylpiperidine

4-[(Indole-6-carbonyl-D-phenylglyciny]aminolpiperidine
(0.10 g, 0.27 mmol) and cyclopentanone afforded 28 mg
(24%) of the title compound.

10 ¹NMR

IS-MS, m/e 445 (M+1).

Example 220.

15 4-[(Indole-6-carbonyl-D-phenylglyciny]amino]-1-(cyclohexylmethyl)piperidine

4-[(Indole-6-carbonyl-D-phenylglyciny]aminolpiperidine
(0.10 g, 0.27 mmol) and cyclohexanecarboxaldehyde
afforded 17 mg (14%) of the title compound.

¹NMR

20 IS-MS, m/e 473 (M+1).

Examples 221-246

Preparation of Starting Materials

25 1-Methyl-4,4'-bispiperidine hydrobromide dihydrobromide

A solution of 4,4'-bipyridine (34.2 g, 100 mmol) in
acetone was treated with methyl p-toluenesulfonate.
After 3 days, the salt (28 g, 80%) was isolated by
filtration. The salt (44.0 g) was then treated with 10%
30 Pd/C in acetic acid (400 mL) and hydrogen gas (4.1 bar)
at 60 °C. After 16 h, the mixture was concentrated, the
residue was dissolved in acetone, and then treated with

hydrogen bromide in acetic acid. The resulting salt (36 g, 86%) was isolated by filtration as a dihydrobromide.

¹NMR

5 1-(D-Phenylglycinyl)-1'-methyl-4,4'-bispiperidine

Using Coupling Method A, benzyloxycarbonyl-D-phenylglycine (16 g, 56 mmol) and 1-methyl-4,4'-bispiperidine dihydrobromide (17.2 g, 50 mmol) afforded, after treatment of the crude acylation product with HBr (150 mL) and acetic acid (150 mL) at 60 °C for 6 h, 8.4 g (54%) of the title compound.

¹NMR

IS-MS, m/e 316 (M+1)

Analysis for C₁₉H₂₉N₃O:

15 Calcd: C, 72.3; H, 9.3; N, 13.3;
Found: C, 71.9; H, 9.2; N, 13.1.

General Procedure: Unless otherwise indicated, the product of Examples 221-246 (or a protected derivative thereof) was prepared from 1-(D-phenylglycinyl)-1'-methyl-4,4'-bispiperidine and the indicated acid using procedures similar to Acylation Method C.

25 Removal of Protecting Group: Where a protecting group was present in the acylation procedure, the procedure for its removal is described.

Example 221.

1-(4-Methoxy-3-methylbenzoyl-D-phenylglycinyl)-1'-methyl-4,4'-bispiperidine
30

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.64 mmol) and 4-methoxy-3-methylbenzoic acid (116 mg, 0.70 mmol) afforded 159 mg (54%) of the title compound.

¹NMR

5 IS-MS, m/e 464 (M+1)

Analysis for C₂₅H₃₃N₃O₃·0.35 H₂O:

Calcd: C, 71.6; H, 8.1; N, 8.9;

Found: C, 71.5; H, 7.8; N, 9.0.

10 Example 222.

1-[5-Methylthiophene-2-carbonyl-D-phenylglyciny1]-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.64 mmol) and 5-methylthiophene-2-carboxylic acid (120 mg, 0.70 mmol) afforded 190 mg (63%) of the title compound.

¹NMR

IS-MS, m/e 472 (M+1)

Example 223.

20 1-(3-Chloro-4-methoxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.64 mmol) and 3-chloro-4-methoxybenzoic acid (130 mg, 0.70 mmol) afforded 182 mg (59%) of the title compound.

25 ¹NMR

IS-MS, m/e 484 (M+1)

Example 224.

30 1-(5-Methoxybenzofuran-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.64 mmol) and 5-methoxybenzofuran-2-carboxylic acid (135 mg, 0.70 mmol) afforded 298 mg (96%) of the title compound.

¹NMR

5 IS-MS, m/e 490 (M+1)

Analysis for C₂₉H₃₅N₃O₄:

Calcd: C, 71.1; H, 7.2; N, 8.6;

Found: C, 71.5; H, 7.4; N, 8.8.

10 Example 225.

1-(5-Acetylthiophene-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.64 mmol) and 5-acetylthiophene-2-carboxylic acid (119 mg, 0.70 mmol) afforded 245 mg (83%) of the title compound.

¹NMR

IS-MS, m/e 468 (M+1)

Analysis for C₂₆H₃₃N₃O₃S:

Calcd: C, 66.8; H, 7.1; N, 9.0;

20 Found: C, 66.5; H, 7.1; N, 9.0.

Example 226.

1-(4-Chloro-3-methylbenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

25 1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 4-chloro-3-methylbenzoic acid (171 mg, 1.00 mmol) afforded 240 mg (51%) of the title compound.

¹NMR

IS-MS, m/e 468 (M+1)

30 Analysis for C₂₆H₃₃N₃O₃S:

Calcd: C, 69.3; H, 7.3; N, 9.0;

Found: C, 68.9; H, 7.2; N, 8.9.

Example 227.

1-(5-Methylindole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 5 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 5-methylindole-2-carboxylic acid (263 mg, 1.50 mmol) afforded 240 mg (51%) of the title compound.

¹NMR

IS-MS, m/e 473 (M+1).

10

Example 228.

1-(5-Methoxyindole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 15 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 5-methoxyindole-2-carboxylic acid (1.50 mmol) afforded 77 mg (16%) of the title compound.

¹NMR

IS-MS, m/e 489 (M+1)

Analysis for C₂₆H₃₃N₃O₃S:

- 20 Calcd: C, 69.3; H, 7.3; N, 9.0;
Found: C, 68.9; H, 7.2; N, 8.9.

Example 229.

1-(Benzothiazole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 25 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and benzothiazole-2-carboxylic acid (200 mg, 1.12 mmol) afforded 180 mg (16%) of the title compound.

¹NMR

- 30 IS-MS, m/e 477 (M-1)

Example 230.

1-(5-Fluoroindole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 5 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 5-fluoroindole-2-carboxylic acid (280 mg, 1.50 mmol) afforded 80 mg (17%) of the title compound.

¹NMR

IS-MS, m/e 477 (M+1)

Analysis for C₂₈H₃₃FN₄O₂·H₂O:

- 10 Calcd: C, 68.0; H, 7.1; N, 11.3;
Found: C, 68.0; H, 6.7; N, 11.1.

Example 231.

- 15 1-(Naphthalene-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and naphthalene-2-carboxylic acid (220 mg, 1.28 mmol) afforded 160 mg (38%) of the title compound.

¹NMR

- 20 IS-MS, m/e 470 (M+1)

Analysis for C₃₀H₃₅N₃O₂·0.5 H₂O:

Calcd: C, 75.3; H, 7.6; N, 8.8;
Found: C, 75.6; H, 7.4; N, 8.9.

25 Example 232.

1-(6-Methoxyindole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- Using Coupling Method C, 1-(D-phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 6-methoxyindole-2-carboxylic acid (191 mg, 1.00 mmol) afforded 200 mg (41%) of the title compound.

¹NMR

IS-MS, m/e 489 (M+1)

Analysis for $C_{29}H_{36}N_4O_3 \cdot 0.5 H_2O$:

Calcd: C, 70.0; H, 7.5; N, 11.3;

Found: C, 69.3; H, 7.5; N, 11.1.

5

Example 233.

1-(5-Chloroindole-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

Using Coupling Method A, 1-(D-phenylglyciny)-1'-methyl-

10 4,4'-bispiperidine (315 mg, 1.00 mmol) and 5-chloroindole-2-carboxylic acid (230 mg, 1.15 mmol) afforded 220 mg (45%) of the title compound.

1H NMR

IS-MS, m/e 493 (M+1)

15 Analysis for $C_{28}H_{33}ClN_4O_2 \cdot 0.75 H_2O$:

Calcd: C, 66.4; H, 6.9; N, 11.1;

Found: C, 66.8; H, 6.6; N, 10.9.

Example 234.

20 1-(3-Hydroxybenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 3-benzyloxybenzoic acid (158 mg, 0.698 mmol) afforded 100 mg (30%) of 1-(3-benzyloxybenzoyl-D-phenyl-

25 glyciny)-1'-methyl-4,4'-bispiperidine. A solution of this material and 10% Pd/C in 3 mL of EtOH was treated with hydrogen gas (1 atm). After 16 h, the mixture was filtered, concentrated, and the residue triturated with EtOAc, affording 27 mg (32%) of the title compound.

30 1H NMR

IS-MS, m/e 436 (M+1).

Example 235.

1-(3-Hydroxy-4-methylbenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 3-acetoxy-4-methylbenzoic acid (135 mg, 0.698 mmol) afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by column chromatography (4% to 6% 2 N NH₃ in methanol:-methylene chloride), 132 mg (46%) of the title compound.

¹NMR

IS-MS, m/e 450 (M+1).

Analysis for C₂₇H₃₅N₃O₃·0.5 H₂O:

Calcd: C, 71.4; H, 7.9; N, 9.3;

Found: C, 71.4; H, 7.9; N, 9.2.

15

The protected starting acid for the above procedure was prepared as follows:

3-Acetoxy-4-methylbenzoic acid

20 A solution of 3-hydroxy-4-methylbenzoic acid (3.0 g, 19.7 mmol) in acetic anhydride (5.6 mL) was treated with sulfuric acid (0.03 mL), heated to 70 °C, cooled and diluted with water. The resulting solid was collected by filtration yielding 1.14 g (30%) of the title compound, which was used
25 without further purification.

¹NMR

Example 236.

1-(2-Hydroxybenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

30

1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 2-acetoxybenzoic acid (125 mg, 0.698 mmol;

prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by column chromatography, 100 mg (36%) of the title compound.

¹NMR

IS-MS, m/e 436 (M+1).

Example 237.

10 1-(4-Chloro-3-hydroxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 4-chloro-3-acetoxybenzoic acid (150 mg, 0.698 mmol; prepared using methods substantially equivalent

15 to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by column chromatography, 110 mg (37%) of the title compound.

¹NMR

20 IS-MS, m/e 470 (M+1).

Example 238.

1-(4-Chloro-2-hydroxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

25 1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 4-chloro-2-acetoxybenzoic acid (150 mg, 0.698 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by radial chromatography, 60 mg (20%) of the title compound.

¹NMR

IS-MS, m/e 470 (M+1).

Example 239.

1- (4-Chloro-3-methoxybenzoyl-D-phenylglyciny1) -1'-methyl-
4,4'-bispiperidine

1- (D-Phenylglyciny1) -1'-methyl-4,4'-bispiperidine (200 mg,
0.635 mmol) and 4-chloro-2-methoxybenzoic acid (130 mg,
0.698 mmol) afforded, after purification by column
chromatography, 120 mg (39%) of the title compound.

¹NMR

IS-MS, m/e 484 (M+1)

Analysis for C₂₇H₃₄ClN₃O₃:

Calcd: C, 67.0; H, 7.1; N, 8.7;

Found: C, 66.8; H, 7.1; N, 8.8.

Example 240.

1- (3-Hydroxy-4-methoxybenzoyl-D-phenylglyciny1) -1'-methyl-
4,4'-bispiperidine

1- (D-Phenylglyciny1) -1'-methyl-4,4'-bispiperidine (200 mg,
0.635 mmol) and 3-acetoxy-4-methoxybenzoic acid (146 mg,
0.698 mmol; prepared using methods substantially equivalent
to those described for 3-acetoxy-4-methylbenzoic acid)
afforded, after treatment of the crude acylation mixture
with methanolic potassium carbonate and purification by
column chromatography, 52 mg (18%) of the title compound.

¹NMR

IS-MS, m/e 466 (M+1).

Example 241.

1- (2,4-Dihydroxybenzoyl-D-phenylglyciny1) -1'-methyl-4,4'-
bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 2,4-diacetoxybenzoic acid (167 mg, 0.698 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid)

- 5 afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by column chromatography, 145 mg (50%) of the title compound.

¹NMR

IS-MS, m/e 452 (M+1).

- 10 Analysis for C₂₆H₃₃N₃O₄·0.75 H₂O:

Calcd: C, 67.2; H, 7.5; N, 9.0;

Found: C, 67.3; H, 7.2; N, 9.3.

Example 242.

- 15 1-(2-Hydroxy-4-methoxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 2-acetoxy-4-methoxybenzoic acid (146 mg, 0.698 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid)

20 afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by ion exchange chromatography (Varian, SCX), 118 mg (40%) of the title compound.

- 25 ¹NMR

IS-MS, m/e 466 (M+1).

Analysis for C₂₇H₃₅N₃O₄·0.50 H₂O:

Calcd: C, 68.3; H, 7.7; N, 8.9;

Found: C, 68.2; H, 7.4; N, 9.1.

Example 243.

1-(5-Chloro-2-hydroxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (200 mg, 0.635 mmol) and 2-acetoxy-5-chlorobenzoic acid (150 mg, 0.698 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the crude acylation mixture with methanolic potassium carbonate and purification by ion exchange chromatography (Varian, SCX), 100 mg (33%) of the title compound.

¹NMR

IS-MS, m/e 471 (M+1).

Analysis for C₂₆H₃₂ClN₃O₃·0.25 H₂O:

Calcd: C, 65.8; H, 6.9; N, 8.9;
Found: C, 65.9; H, 7.0; N, 9.2.

Example 244.

1-(3-Chloro-4-hydroxybenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

1-(D-Phenylglyciny1)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 4-acetoxy-3-chlorobenzoic acid (321 mg, 1.50 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the acylation mixture with methanolic potassium carbonate, 50 mg (27%) of the title compound.

¹NMR

IS-MS, m/e 470 (M+1).

Analysis for C₂₆H₃₂ClN₃O₃·1.0 H₂O:

Calcd: C, 64.0; H, 7.0; N, 8.6;
Found: C, 63.7; H, 7.0; N, 8.7.

Example 245.

1-(3-Hydroxynaphthalene-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 5 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 3-acetoxynaphthalene-2-carboxylic acid (300 mg, 1.30 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the acylation product
10 with methanolic potassium carbonate, 128 mg (38%) of the title compound.

¹NMR

IS-MS, m/e 486 (M+1).

15 Example 246.

1-(6-Hydroxynaphthalene-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 1-(D-Phenylglyciny)-1'-methyl-4,4'-bispiperidine (315 mg, 1.00 mmol) and 6-acetoxynaphthalene-2-carboxylic acid
20 (300 mg, 1.30 mmol; prepared using methods substantially equivalent to those described for 3-acetoxy-4-methylbenzoic acid) afforded, after treatment of the acylation product with methanolic potassium carbonate, 210 mg (43%) of the title compound.

25 ¹NMR

IS-MS, m/e 486 (M+1).

Analysis for C₃₀H₃₅N₃O₃·1.0 H₂O:

Calcd: C, 71.6; H, 7.4; N, 8.3;

Found: C, 71.5; H, 7.3; N, 8.3.

30

Examples 247-251.

Preparation of Starting Materials

1-(Benzyloxycarbonyl-D-phenylglyciny)l)piperidine-4-methanol

Using Coupling Method C, benzyloxycarbonyl-D-phenylglycine (8.41 g, 29.5 mmol) and 4-piperidinemethanol (3.85 g, 37.4 mmol) afforded 10.2 g (93%) of the title compound.

¹NMR

1-(D-Phenylglyciny)l)piperidine-4-methanol

Using Deprotection Method C, 1-(benzyloxycarbonyl-D-phenylglyciny)l)piperidine-4-methanol (3.93 g, 29.5 mmol) and 10% palladium on carbon (1.30 g) afforded 2.31 g (88%) of the title compound.

¹NMR

IS-MS, m/e 249 (M+1).

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-methanol

Using methods substantially equivalent Acylation Method C described prior to Example 201, 1-(D-phenylglyciny)l)-piperidine-4-methanol (1.23 g, 4.96 mmol) and p-anisoyl chloride (0.888 g, 5.21 mmol) afforded, after purification by column chromatography (SiO₂: 1:1 to 1:9 hexanes:EtOAc), 1.26 g (66%) of the title compound.

¹NMR

IS-MS, m/e 383 (M+1).

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-carboxaldehyde

A solution of 1-(4-methoxybenzoyl-D-phenylglyciny)l)-piperidine-4-methanol (0.800 g, 2.08 mmol) and N-methyl-morpholine oxide (0.366 g, 3.13 mmol) in methylene chloride (15 mL) was treated with tetrapropylammonium perruthenate (TPAP, 2 mg). After 14 h, the mixture was treated with

additional TPAP (5 mg). After 20 h, the mixture was treated with additional TPAP (5 mg). After 32 h, the mixture was loaded directly onto a column and purified by column chromatography (SiO₂: 1:1 to 1:4 hexanes:EtOAc) affording
5 0.286 g (36%) of the title compound.

¹NMR

IS-MS, m/e 381 (M+1).

General Procedure: Unless otherwise indicated, the product
10 of Examples 247-251 was obtained from the indicated amine and 1-(4-methoxybenzoyl-D-phenylglyciny)l)piperidine-4-carboxaldehyde using Alkylation Method D.

Example 247.

15 1-[(4-Methoxybenzoyl-D-phenylglyciny)l]-4-[(isopropylamino)-methyl]piperidine hydrochloride

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and isopropylamine afforded, after treatment of the isolated product with
20 excess hydrochloric acid in methanol and concentration, 37 mg of the title compound as a hydrochloride salt.

¹NMR

IS-MS, m/e 424 (M+1)

25 Example 248.

1-(4-Methoxybenzoyl-D-phenylglyciny)l)-4-[(dimethylamino)-methyl]piperidine

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and dimethylamine
30 afforded 25 mg (47%) of the title compound.

¹NMR

IS-MS, m/e 410 (M+1)

Example 249.

1-[(4-Methoxybenzoyl-D-phenylglyciny)]-4-[(N,N-diethyl-amino)methyl]piperidine hydrochloride

- 5 1-(4-Methoxybenzoyl-D-phenylglyciny)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and diethylamine afforded, after treatment of isolated product with excess hydrochloric acid in methanol and concentration, 42 mg of the title compound as a hydrochloride salt.

10 ¹NMR

IS-MS, m/e 438 (M+1)

Example 250.

1-[(4-Methoxybenzoyl-D-phenylglyciny)]-4-[(1-pyrrolidinyl)-methyl]piperidine

15

1-(4-Methoxybenzoyl-D-phenylglyciny)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and pyrrolidine afforded 27 mg (47%) of the title compound.

¹NMR

20 IS-MS, m/e 436 (M+1)

Example 251.

1-[(4-Methoxybenzoyl-D-phenylglyciny)]-4-[(3-pyrrolin-1-yl)methyl]piperidine hydrochloride

- 25 1-(4-Methoxybenzoyl-D-phenylglyciny)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and 3-pyrroline afforded, after treatment of isolated product with excess hydrochloric acid in methanol and concentration, 43 mg of the title compound as a hydrochloride salt.

30 ¹NMR

IS-MS, m/e 434 (M+1)

Examples 252 to 253

Preparation of Starting Materials

4-[(Benzyloxycarbonyl-D-phenylglyciny]aminomethyl]-
5 piperidine

Using Deprotection Method D, 4-[(benzyloxycarbonyl-D-phenylglyciny]aminomethyl]-1-Boc piperidine (2.70 g, 5.61 mmol) afforded 1.56 g (73%) of the title compound.

¹NMR

10 IS-MS, m/e 382 (M+1)

4-[(Benzyloxycarbonyl-D-phenylglyciny]aminomethyl]-1-cyclopentylpiperidine

Using Alkylation Method D, 4-[(benzyloxycarbonyl-D-phenylglyciny]aminomethyl]piperidine (1.50 g, 3.93 mmol) and cyclopentanone afforded 3.48 g (91%) of the title compound.

¹NMR

IS-MS, m/e 450 (M+1)

20

4-[(D-Phenylglyciny]aminomethyl]-1-cyclopentylpiperidine

Using a deprotection procedure similar to that described above for preparation of 1-(D-phenylglyciny]-1'-methyl-

4,4'-bispiperidine, 4-[(benzyloxycarbonyl-D-phenylglyciny]-aminomethyl]-1-cyclopentylpiperidine (1.70 g, 3.78 mmol) afforded 0.75 g (63%) of the title compound.

¹NMR

IS-MS, m/e 316 (M+1)

30 General Procedure: Using Coupling Method A, 4-[(D-phenylglyciny]aminomethyl]-1-cyclopentylpiperidine was coupled with the indicated acid.

Example 252.

4-[(5-Chloroindole-2-carbonyl-D-phenylglyciny]aminomethyl]-1-cyclopentylpiperidine

- 5 4-[(D-Phenylglyciny]aminomethyl]-1-cyclopentylpiperidine
(0.100 g, 0.317 mmol) and 5-chloroindole-2-carboxylic
acid (0.075 g, 0.38 mmol) afforded 156 mg (98%) of the
title compound.

¹NMR

- 10 IS-MS, m/e 493 (M+1)

Example 253.

4-[(3-Methylindole-6-carbonyl-D-phenylglyciny]aminomethyl]-1-cyclopentylpiperidine

- 15 4-[(D-Phenylglyciny]aminomethyl]-1-cyclopentylpiperidine
(0.100 g, 0.317 mmol) and 3-methylindole-6-carboxylic
acid (0.067 g, 0.38 mmol) afforded 137 mg (91%) of the
title compound.

¹NMR

- 20 IS-MS, m/e 473 (M+1)

Particular Analytical Methods for Examples 254-276:

- HPLC Analysis (Method A): Dynamax (trademark) C18, 60Å
25 column. The elution system consisted of a linear gradient
from 90:10 (95% H₂O, CH₃CN)/(95% CH₃CN, H₂O) to (95% CH₃CN,
H₂O) over 20 min, followed by (95% CH₃CN, H₂O) isocratic
elution over 15 min. The flow rate was 1 mL/min. UV
Detection was performed at 254 nm unless otherwise noted.

30

HPLC Analysis (Method B): Microsorb-MV (trademark) C8 (4.6 x
250 mm) column. The elution system consisted of a linear

gradient from 95:5 (2.5% TFA in H₂O):(2.5% TFA in acetonitrile) to 0:100 (2.5% TFA in H₂O):(2.5% TFA in acetonitrile) over 25 min at 30 °C and a flow rate of 1 mL/min. UV Detection was performed at 254 nm unless
5 otherwise noted.

HPLC Analysis (Method C): Dynamax (trademark), C18, 60Å column. The elution system consisted of a linear gradient from 95:5 (0.2% TFA in H₂O)/ (0.2% TFA in CH₃CN) to 5:95
10 (0.2% TFA in H₂O)/ (0.2% TFA in CH₃CN) over 20 min, followed by (0.2% TFA in CH₃CN) isocratic elution over 15 min. The flow rate was 1 mL/min. UV Detection was performed at 254 nm unless otherwise noted.

15 HPLC Analysis (Method D): Waters Symmetry (trademark), C18 (4.6 x 250 mm) column. The elution system consisted of a linear gradient from 95:5 (0.2% TFA in H₂O)/(0.2% TFA in CH₃CN) to 5:95 (0.2% TFA in H₂O)/(0.2% TFA in CH₃CN) over
20 min, followed by (0.2% TFA in CH₃CN) isocratic over 15 min. The flow rate was 1 mL/min. UV Detection was performed at 254 nm unless otherwise noted.

HPLC Analysis (Method E): Microsorb-MV C18 (4.6 x 250 mm) column. The elution system consisted of a linear gradient
25 from 90:10 (2.5% TFA in H₂O):(2.5% TFA in acetonitrile) to 10:90 (2.5% TFA in H₂O):(2.5% TFA in acetonitrile) over 25 min at 30 °C and a flow rate of 1 mL/min. UV Detection was performed at 254 nm unless otherwise noted.

30 API-MS (atmospheric pressure chemical ionization mass spectra) were obtained on a PESCiex (trademark) API 150EX

with a heated nebulizer and nitrogen as the reagent gas in positive ion mode.

Examples 254 to 257

5 Preparation of Starting Materials

(R)-(-)-Boc-phenylglycinol: Di-tert-butyl dicarbonate (232.4 g, 1.06 mol) was added to a well stirred, ice bath cooled mixture of (R)-(-)-2-phenylglycinol (121.7 g, 0.887 mol), potassium carbonate (171.7 g, 1.24 mol), 1,4-dioxane (1 L), and water (1 L). The temperature rose from 5 °C - 11 °C during the addition. The reaction was allowed to stir overnight. The reaction was diluted with water (1 L), and cooled in ice-water. The resultant precipitate was collected by vacuum filtration, washed with water, air dried, and vacuum dried at 40 °C overnight to afford 201.7 g (95%) as a white solid.

¹H-NMR(CDCl₃)

TLC R_f = 0.45 (83% CH₂Cl₂, EtOAc)

20

(R)-(-)-[2-[(Methylsulphonyl)oxy]-1-phenylethyl]carbamic acid 1,1-dimethylethyl ester

The sulphonate was prepared from the above alcohol according to *J. Med. Chem.* 1994, 37, 1819.

25 ¹H-NMR(CDCl₃)

TLC R_f = 0.45 (95% CH₂Cl₂, EtOAc)

(R)-2-[(Butoxycarbonyl)amino]-2-phenylethyl azide

30 The azide was prepared from the above sulphonate according to *J. Med. Chem.* 1994, 37, 1819.

¹H-NMR(CDCl₃)

TLC R_f = 0.85 (95% CH₂Cl₂, EtOAc)

(R)-2-(4-Methoxybenzoylamino)-2-phenylethyl azide

(R)-2-[(Butoxycarbonyl)amino]-2-phenylethyl azide (47.8 g, 0.182 mole) was added to trifluoroacetic acid (500 mL) with

5 stirring and ice-water bath cooling. The cooling bath was removed, the reaction was allowed to stir 1 h, and the solvent was removed in vacuo at 35 °C water bath

temperature. The residue was co-evaporated with toluene to give a weight of 75.0 g. The residue was dissolved in

10 1,4-dioxane (500 mL) and water (500 mL), with ice-water bath cooling, and then potassium carbonate (113.5 g, 0.82 mol), and anisoyl chloride (37.3 g, 0.219 mol) were added.

Another portion of 1,4-dioxane (300 mL) was added to

15 facilitate stirring. After stirring over the weekend, water (1 L) was added. The mixture was cooled to -15 °C, and vacuum filtered to collect a white solid. The solid was washed with water, air dried, and then dried under vacuum at 50 °C for 4 h to afford 46.3 g (86%).

¹H-NMR(CDCl₃)

20 TLC R_f = 0.85 (83% CH₂Cl₂, EtOAc)

(R)-2-(4-Methoxybenzoylamino)-2-phenylethylamine

(R)-2-(4-methoxybenzoylamino)-2-phenylethyl azide (46.3 g) was combined with 10% palladium on carbon in THF (400 mL),

25 methanol (100 mL) and was stirred under a hydrogen atmosphere. Analysis by TLC (70% methylene chloride, ethyl acetate) indicated absence of starting material after stirring overnight. The solution was filtered through diatomaceous earth, rinsed with THF, and evaporated. The

30 resulting solid was recrystallized with ethyl acetate, and dried under vacuum at 60 °C for 1 h to afford 35.4 g (84%) of a white crystalline solid.

$^1\text{H-NMR}$ (CDCl_3)

TLC R_f = 0.17 (90% CH_2Cl_2 , 9% Methanol, 1% NH_4OH)

Examples 254-257 were prepared from (R)-2-(4-methoxybenzoyl-
5 amino)-2-phenylethylamine and the indicated acid chloride
using the acylation method described in Example 254
(Acylation Method A).

Example 254.

10 (R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-4-methyl-
benzamide

(Acylation Method A) p-Toluoyl chloride (0.22 mL, 1.6 mmol)
was added via syringe to a 15 °C stirring mixture of (R)-2-
(4-methoxybenzoylamino)-2-phenylethylamine (0.40 g, 1.48
15 mmol), potassium carbonate (0.27 g, 1.9 mmol), 1,4-dioxane
(8 mL), and water (4 mL). TLC analysis (80% methylene
chloride, 18% methanol, 2% ammonium hydroxide) indicated
reaction completion within 1 h. The solution was diluted
with water, and the precipitated solid was collected by
20 vacuum filtration. The precipitate was recrystallized from
methanol and dried under vacuum at 50 °C overnight to afford
the title compound (0.42 g, 72%) as a white solid.

$^1\text{H-NMR}$ (DMSO)

IS-MS, m/e = 389 ($M+1$)

25 Analysis for $\text{C}_{24}\text{H}_{24}\text{N}_2\text{O}_3$:

Calcd: C, 74.21; H, 6.23; N, 7.21;

Found: C, 73.82; H, 6.32; N, 7.04.

HPLC Analysis (Method A): 99.3%, RT: 21.35 min.

Melting Point: 230-238 °C

Example 255.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-4-ethylbenzamide

Prepared from 4-ethylbenzoyl chloride (84%).

5 ¹H-NMR (DMSO)

IS-MS, m/e = 403 (M+1)

Analysis for C₂₅H₂₆N₂O₃:

Calcd: C, 74.60; H, 6.51; N, 6.96;

Found: C, 74.25; H, 6.63; N, 6.83.

10 HPLC Analysis (Method A): 95.4%, RT=22.62 min.

Melting Point: 222-229 °C

Example 256.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-4-isopropylbenzamide

15

Prepared from 4-isopropylbenzoyl chloride (40%).

¹H-NMR (DMSO)

IS-MS, m/e = 417 (M+1)

Analysis for C₂₆H₂₈N₂O₃:

20 Calcd: C, 74.97; H, 6.78; N, 6.73;

Found: C, 74.61; H, 6.78; N, 6.61.

HPLC Analysis (Method A): 98.4%, RT=23.77 min.

Melting Point: 239-244 °C

25 Example 257.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-4-tert-butylbenzamide

Prepared from 4-tert-butylbenzoyl chloride (89%).

¹H-NMR (DMSO)

30 IS-MS, m/e = 431 (M+1)

Analysis for C₂₇H₃₀N₂O₃·0.25H₂O:

Calcd: C, 74.54; H, 7.07; N, 6.44;

Found: C, 74.39; H, 7.13; N, 6.34.

HPLC Analysis (Method A): 96.4%, RT=25.04 min.

Melting Point = 171-175 °C

5 Examples 258 to 266

Preparation of Starting Materials

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-tert-butoxycarbonylpiperidine-4-carboxamide.

- 10 N-Boc-iso-nipecotic acid (2.13 g, 9.5 mmol) followed by (R)-2-(4-methoxybenzoylamino)-2-phenylethylamine (2.34 g, 8.7 mmol) were added at 2 °C to a stirring mixture of EDCI (2.5 g, 13.0 mmol), and HOBT (1.64 g, 12.1 mmol) in DMF (50 mL). Triethylamine (1.8 mL, 13.0 mmol) was added
- 15 dropwise. The reaction was allowed to warm to room temperature, with stirring overnight. Water (100 mL) was added, and the aqueous mixture was extracted with ethyl acetate (2 X 200 mL). The extracts were combined, and THF (200 mL) was added. Next, the organic layers were washed
- 20 with water (5 X 70 mL), aqueous NaHCO₃ (70 mL), and brine (100 mL). The organic layer was dried over Na₂SO₄, filtered, and evaporated. The crude residue (4.2 g, 100%), was recrystallized from ethyl acetate and hexanes to afford 2.9 g (71%) of a white solid.

25 ¹H-NMR (DMSO)

IS-MS, m/e = 482 (M+1)

Analysis for C₂₇H₃₀N₂O₃:

Calcd: C, 67.34; H, 7.33; N, 8.73;

Found: C, 67.34; H, 7.46; N, 8.66.

- 30 HPLC Analysis (Method A): 98.8%, RT=20.72 min.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]piperidine-4-carboxamide trifluoroacetate

(Deprotection Method A) Trifluoroacetic acid was added to a stirring suspension of (R)-N-[2-(4-methoxybenzoylamino)-2-phenylethyl]-1-tert-butoxycarbonylpiperidine-4-carboxamide (2.0 g, 4.2 mmol), methylene chloride (20 mL), and anisole (0.5 g, 4.6 mmol) at room temperature. A solution was obtained and bubbling was observed. After 1 h, the reaction mixture was evaporated at 40 °C. The residue was taken up in warm methanol, and to this stirring solution was added ether to precipitate the product. The precipitate was collected by vacuum filtration, washed with ethyl acetate, then dried under vacuum at 60 °C overnight to afford 1.9 g (92%) of a white solid.

¹H-NMR(DMSO)

IS-MS, m/e = 382 (M+1)

Analysis for C₂₄H₂₈F₃N₃O₅:

Calcd: C, 58.18; H, 5.70; N, 8.48;

Found: C, 58.19; H, 5.78; N, 8.27.

HPLC Analysis (Method C): >99%, RT=20.40 min.

Except as otherwise noted, Examples 258-266 were prepared from (R)-N-[2-(4-methoxybenzoylamino)-2-phenylethyl]-piperidine-4-carboxamide trifluoroacetate and the indicated aldehyde or ketone using the reductive alkylation method described in Example 258 (Alkylation Method A).

Example 258.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-isopropylpiperidine-4-carboxamide

(Alkylation Method A) (R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]piperidine-4-carboxamide trifluoroacetate

(0.50 g, 1.0 mmol), acetone (4.5 mL, 61 mmol), acetic acid (0.28 mL, 4.9 mmol), and sodium cyanoborohydride (0.32 g, 5.1 mmol) were combined in methanol, and stirred. After 4 h, TLC (79% CH₂Cl₂, 19% methanol, 1% NH₄OH) indicated
5 reaction completion. The solution was diluted with methanol (100 mL), and passed through H⁺ form ion exchange resin (Varian SCX cartridge, Catalog #1225-6035) washed with methanol, and then with 2 M NH₃ in methanol to collect the product. The product was recrystallized from methanol and
10 ether to afford 0.30 g (70%) of a white crystalline solid:

¹H-NMR (DMSO)

IS-MS, m/e = 424 (M+1)

Analysis for C₂₅H₃₃N₃O₃·0.75H₂O:

Calcd: C, 68.70; H, 7.96; N, 9.61;

15 Found: C, 68.73; H, 7.68; N, 9.29.

HPLC Analysis (Method C): >99% RT=18.19 min.

Examples 259-262 were purified by passing a solution through a silica gel column, eluting with 200:10:1 methylene
20 chloride, methanol, and concentrated ammonium hydroxide.

Example 259.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-cyclopentylpiperidine-4-carboxamide

25 Prepared from cyclopentanone (44%).

¹H-NMR (DMSO)

IS-MS, m/e = 450 (M+1)

Analysis for C₂₇H₃₅N₃O₃·0.25H₂O:

Calcd: C, 71.42; H, 7.88; N, 9.25;

30 Found: C, 71.21; H, 7.93; N, 9.18.

HPLC Analysis (Method C): >99%, RT=18.84 min.

Melting Point = 253-257 °C

Example 260.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-cyclohexylpiperidine-4-carboxamide

5 Prepared from cyclohexanone (65%).

¹H-NMR (DMSO)

IS-MS, m/e = 464 (M+1)

Analysis for C₂₈H₃₇N₃O₃·1.0H₂O:

Calcd: C, 69.83; H, 8.16; N, 8.72;

10 Found: C, 69.64; H, 7.84; N, 8.90.

HPLC Analysis (Method C): >99%, RT=19.13 min.

Melting Point = 239-243 °C.

Example 261.

15 (R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-ethyl-piperidine-4-carboxamide

Prepared from acetaldehyde (36%).

¹H-NMR (DMSO)

IS-MS, m/e 410 (M+1)

20 Analysis for C₂₄H₃₁N₃O₃:

Calcd: C, 70.39; H, 7.63; N, 10.26;

Found: C, 70.06; H, 7.67; N, 10.00.

HPLC Analysis (Method D): 96.9%, RT=16.04 min.

Melting Point = 245-251 °C.

25

Example 262.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-(1-methyl-piperidin-4-yl)piperidine-4-carboxamide

Prepared from 1-methylpiperid-4-one (27%).

30 ¹H-NMR (DMSO)

IS-MS, m/e 479 (M+1)

Analysis for C₂₈H₃₈N₄O₃·0.25H₂O:

Calcd: C, 69.61; H, 8.03; N, 11.60;

Found: C, 69.72; H, 8.11; N, 11.48.

HPLC Analysis (Method D): 97.0%, RT=15.42 min.

Melting Point = 252-259 °C.

5

(No example for Examples 263-264.)

Examples 265-266 were purified by passing a solution through a silica gel column, eluting with 200:10:1 methylene chloride, methanol, and concentrated ammonium hydroxide.

Example 265.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-(3-pyridinylmethyl)piperidine-4-carboxamide

15 Prepared from pyridine-3-carboxaldehyde (68%).

¹H-NMR (DMSO)

CI-MS, m/e = 473 (M+1)

HPLC Analysis (Method D): 92.7%, RT=15.39 min.

20 Example 266.

(R)-N-[2-(4-Methoxybenzoylamino)-2-phenylethyl]-1-(4-pyridinylmethyl)piperidine-4-carboxamide

Prepared from pyridine-4-carboxaldehyde (63%).

¹H-NMR (DMSO)

25 CI-MS, m/e = 473 (M+1)

HPLC Analysis (Method D): 89.2%, RT=15.33 min.

Example 267.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(4-piperidinyl-30 methyl)piperazine trifluoroacetate

1-[D-(+)-Benzyloxycarbonylphenylglycinyll]-4-(tert-butoxycarbonyl)piperazine.

(Coupling Method A) D-(+)-Benzyloxycarbonylphenylglycine (58.0 g, 203 mmol) and 1-Boc-piperazine (41.7 g, 224 mmol)

5 were dissolved in DMF (1 L) and cooled to approximately -15 °C in an ice-methanol bath. Diethyl cyanophosphonate (37.0 mL, 244 mmol) was slowly added to the mixture. Triethylamine (59.4 mL, 426 mmol) was added dropwise to the solution. The mixture was stirred at -15 °C for 2 h and was
10 allowed to gradually warm to room temperature overnight.

The mixture was diluted with ethyl acetate and water. The layers were separated, and the water layer extracted with ethyl acetate. The organic layers were combined, washed with 10% citric acid (2 x 500 mL) and brine, dried (Na₂SO₄),
15 filtered and concentrated under vacuum. The crude product was filtered through a plug of silica gel (1.2 kg) using 1:1 hexanes:ethyl acetate as eluent to provide 1-[D-(+)-benzyloxycarbonylphenylglycinyll]-4-(tert-butoxycarbonyl)piperazine (69.9 g, 76%) as a colorless oil.

20 ¹H-NMR(CDCl₃)

API-MS, m/e = 454 (M+1)

1-[D-(+)-Phenylglycinyll]-4-(tert-butoxycarbonyl)piperazine

1-[D-(+)-Benzyloxycarbonylphenylglycinyll]-4-(tert-butoxycarbonyl)piperazine (69.5 g, 153 mmol) was dissolved in
25 ethanol (500 mL). The mixture was degassed with nitrogen and Pd/C (6.8 g) was added. Hydrogen was bubbled through the mixture for 1 h, and it was maintained under a hydrogen atmosphere for 16 h. The Pd/C was removed by filtration
30 through cellulose powder. The filter cake was rinsed with ethanol and ethyl acetate. The filtrate was concentrated under vacuum to give 1-[D-(+)-phenylglycinyll]-4-(tert-

butoxycarbonyl)piperazine (45.3 g, 93%) as a light yellow solid.

¹H-NMR(CDCl₃)

API-MS, m/e = 320 (M+1)

5

1-(4-Methoxybenzoyl-D-phenylglycinyll)-4-(tert-butoxycarbonyl)piperazine

(Acylation Method B) 1-[D-(+)-phenylglycinyll]-4-(tert-butoxycarbonyl)piperazine (42.0 g, 131.5 mmol) was dissolved
10 in 1,4-dioxane (420 mL) and water (210 mL) and was cooled to 10 °C. Potassium carbonate (36.4 g, 263 mmol) was added, followed by p-anisoyl chloride (24.7 g, 144 mmol). The mixture was stirred at room temperature overnight. The mixture was diluted with water and ethyl acetate. The
15 layers were separated, and the water layer extracted with ethyl acetate. The organic layers were combined, washed with brine, dried, filtered and concentrated to provide 1-(4-methoxybenzoyl-D-phenylglycinyll)-(4-tert-butoxycarbonyl)piperazine (58.7 g, 98%) as an off-white solid.

20 ¹H-NMR(CDCl₃)

API-MS, m/e = 454 (M+1)

1-(4-Methoxybenzoyl-D-phenylglycinyll)piperazine trifluoroacetate

25 1-(4-Methoxybenzoyl-D-phenylglycinyll)-(4-tert-butoxycarbonyl)piperazine (20.0 g, 44.1 mmol) was dissolved in dichloromethane (50 mL) and anisole (20 mL). To this vigorously stirred mixture was added trifluoroacetic acid (50 mL). The mixture was stirred for 25 min at room
30 temperature. The solvents were removed under vacuum. The residue was triturated in ether and sonicated for 60 min. The solid was collected by filtration and dried in a vacuum

pistol overnight to provide 1-(4-methoxybenzoyl-D-phenylglyciny)l)piperazine trifluoroacetate (18.2 g, 88%) as a light yellow solid.

¹H-NMR(CD₃OD)

5 API-MS, m/e = 354 (M+1)

1-Boc-isonipecotic acid

Isonipecotic acid (15.0 g, 116 mmol) was dissolved in THF (300 mL), water (150 mL) and 6 N NaOH (40 mL). Di-tert-butyl dicarbonate (26.6 g, 122 mmol) was added and the
10 mixture stirred overnight. The mixture was diluted with water and ethyl acetate, and the layers separated. The water layers were extracted with ethyl acetate, and the organic layers discarded. The water layer was diluted with
15 KHSO₄ (2 N, pH-4) and extracted with ethyl acetate. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated to provide 1-Boc-isonipecotic acid (23.9 g, 90%) as a white solid.

¹H-NMR(CDCl₃)

20 API-MS, m/e = 230 (M+1)

1-Boc-piperidine-4-methanol

1-Boc-isonipecotic acid (10.0 g, 214 mmol) was dissolved in THF (400 mL) and cooled to 0 °C. A solution of BH₃·THF
25 (180 mL, 1 N in THF, 180 mmol) was added slowly. The mixture stirred for 1 h at 0 °C and was allowed to warm to room temperature for 12 h. The mixture was carefully quenched with water and diluted with ethyl acetate. The water layer was extracted with ethyl acetate. The organic
30 layers were combined, washed with brine, dried (Na₂SO₄), filtered and concentrated to provide 1-Boc-piperidine-4-methanol (7.98 g, 85%) as a white solid.

¹H-NMR (CDCl₃)

API-MS, m/e = 220 (M+1)

1-Boc-piperidine-4-carboxaldehyde

- 5 Dimethyl sulfoxide (3.5 mL, 48.7 mmol) was dissolved in dichloromethane (100 mL) and was cooled to -78 °C. Oxalyl chloride (3.65 mL, 41.8 mmol) was added. The mixture stirred for 30 min. To this solution was added a solution of 1-Boc-piperidine-4-methanol (7.5 g, 34.8 mmol) in 10 dichloromethane (15 mL), and the mixture stirred for 1 h. Triethylamine (9.7 mL, 69.6 mmol) was added slowly and the mixture stirred at -78 °C for 30 min and warmed to room temperature over the course of 1 h. The mixture was diluted with water and the layers separated. The water layer was 15 extracted with dichloromethane and the organic layers combined, dried (Na₂SO₄), filtered and concentrated to provide 1-Boc-piperidine-4-carboxaldehyde (6.75 g, 91%) as a yellow oil.

¹H-NMR (CDCl₃)

- 20 API-MS, m/e = 214 (M+1)

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(1-Boc-piperidin-4-ylmethyl)piperazine

- (Alkylation Method B) Using Alkylation Method A, except 25 using sodium triacetoxymethylborohydride in 1,2-dichloroethane, 1-(4-methoxybenzoyl-D-phenylglyciny)-4-(1-Boc-piperidin-4-ylmethyl)piperazine was prepared from 1-(4-methoxybenzoyl-D-phenylglyciny)piperazine trifluoroacetate and 1-Boc-piperidine-4-carboxaldehyde (85%).

- 30 ¹H-NMR (CDCl₃)

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-piperidinyl-methyl)piperazine trifluoroacetate.

Using Deprotection Method A, the title compound was prepared from 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(1-Boc-

5 piperidin-4-ylmethyl)piperazine (90%).

Melting Point = 70-72 °C with decomposition

IR(KBr)

¹H-NMR(CD₃OD)

API-MS, m/e = 451 (M+1)

10 Analysis for C₂₆H₃₄N₄O₃·2.5TFA·0.4H₂O:

Calcd: C, 50.12; H, 5.06; N, 7.54;

Found: C, 49.81; H, 5.33; N, 7.39.

HPLC Analysis (Method B): 97.1% RT=14.3 min.

15 Examples 268 to 272

Unless otherwise indicated, using Alkylation Method A or B, the title compounds were prepared from 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(4-piperidinylmethyl)piperazine trifluoroacetate and the indicated aldehyde or ketone.

20

Example 268.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(1-methylpiperidin-4-ylmethyl)piperazine hydrochloride

Prepared from paraformaldehyde using Method A (56%).

25 IR (KBr)

¹H-NMR(CD₃OD)

CI-MS, m/e = 465 (M+1)

Example 269.

30 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(1-isopropyl-piperidin-4-ylmethyl)piperazine hydrochloride

Prepared from acetone using Method A (72%).

Melting Point = 172-180 °C with decomposition

IR (KBr)

¹H-NMR (CD₃OD)

CI-MS, m/e = 493 (M+1)

5 Analysis for C₂₉H₄₀N₄O₃·3HCl:

Calcd: C, 55.85; H, 7.34; N, 8.98;

Found: C, 55.63; H, 7.32; N, 8.66.

HPLC Analysis (Method B): 98.2% RT=14.4 min.

10 Example 270.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-[3-(3-pyridinyl)-propyl]piperazine hydrochloride

Prepared from pyridine-3-propionaldehyde (prepared as described below) using Method B (72%).

15 ¹H-NMR (CD₃OD)

CI-MS, m/e = 473 (M+1)

Pyridine-3-propionaldehyde

(Oxidation Method A) 1,1,1-Triacetoxy-1,1-dihydro-1,2-

- 20 benziodoxol-3(1H)-one (5.4 g, 12.7 mmol) was suspended in dichloromethane (45 mL). 3-Pyridinepropanol (1.59 g, 11.6 mmol) as a solution in dichloromethane (35 mL) was added slowly. The mixture stirred for 3 h at room temperature. The mixture was diluted with saturated aqueous NaHCO₃ and
- 25 ether. The mixture was stirred for 10 min and was diluted with sodium thiosulfate (2 N) and stirred until the solids dissolved. The layers were separated, and the water layer was extracted with ether. The organic layers were combined, washed with water and brine, dried (Na₂SO₄), filtered and
- 30 concentrated to provide pyridine-3-propionaldehyde (1.03 g, 66%) as a yellow oil.

¹H-NMR (CDCl₃)

Example 271.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[3-(4-pyridiny1)-propy1]piperazine hydrochloride.

- 5 Prepared from pyridine-4-propionaldehyde (prepared as described below) using Method A; the hydrochloride salt was prepared using HCl (2 M) in diethyl ether (76%).

¹H-NMR (CD₃OD)

CI-MS, m/e = 473 (M+1)

10

Pyridine-4-propionaldehyde

Prepared from 4-pyridinepropanol using Oxidation Method A (80%).

¹H-NMR (CDCl₃)

15

Example 272.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(2-cyclopentylethyl)piperazine hydrochloride hydrate

- 20 The free base was prepared from cyclopentylacetaldehyde (prepared as described below) using Method B (58%).

¹H NMR (CDCl₃)

- 25 To a stirred solution of 1-(4-methoxybenzyl-D-phenylglyciny1)-4-(2-cyclopentylethyl)piperazine (260 mg, 0.58 mmol) in ether (10 mL) and methylene chloride (1 mL) was added hydrogen chloride as a 2 N solution in ether (about 2 mL), and the resulting precipitate was filtered to give 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(2-cyclopentylethyl)piperazine hydrochloride as a pale yellow solid.

¹H NMR (CD₃OD)

- 30 IS-MS, m/e = 450 (M+1)

Analysis for C₂₇H₃₅N₃O₃·HCl·0.5H₂O:

Calcd: C, 65.51; H, 7.53; N, 8.49;

Found: C, 65.67; H, 7.58; N, 8.13.

HPLC Analysis (Method E): >99%, RT=15.84

Melting Point = 190-192 °C

5 **Cyclopentylacetaldehyde**

Using Oxidation Method A, the title compound was prepared from 2-cyclopentylethanol and used with trace amounts of ether and methylene chloride present due to volatility of product.

10 ¹H NMR (CDCl₃)

Example 273.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(3-pyrrolidiny)-piperazine trifluoroacetate.

15

(R)-(+)-1-Boc-3-pyrrolidinol

To a stirred solution of (R)-(+)-3-pyrrolidinol (2 g, 22.96 mmol) in tetrahydrofuran (60 mL) and water (30 mL) was added di-tert-butyl dicarbonate (5.27 g, 24.15 mmol) and 3 N

20 sodium hydroxide (16 mL), and the resulting solution was stirred for 6 h. Another portion of di-tert-butyl

dicarbonate (0.74 g, 0.34 mmol) was added and the solution was stirred overnight. The reaction was diluted with water (40 mL) and extracted with ethyl acetate (2 x 150 mL). The

25 combined organic extracts were washed with 2 N potassium hydrogen sulfate (200 mL), saturated sodium bicarbonate (2 x 150 mL), brine (150 mL) and dried over magnesium sulfate.

Removal of solvent in vacuo gave (R)-(+)-1-Boc-3-pyrrolidinol (4.21 g, 98%) as a yellow oil.

30 ¹H-NMR (CDCl₃)

1-Boc-3-pyrrolidinone

Using Oxidation Method A, the title compound was prepared from (R)-(+)-1-Boc-3-pyrrolidinol (85%).

¹H NMR (CDCl₃)

5 1-(4-Methoxybenzyl-D-phenylglyciny1)-4-(1-Boc-3-pyrrolidinyl)piperazine

Using Alkylation Method B, the title compound was prepared (69%) from 1-(4-methoxybenzyl-D-phenylglyciny1)piperazine trifluoroacetate and 1-Boc-3-pyrrolidinone.

10 ¹H NMR (CDCl₃)

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(3-pyrrolidinyl)-piperazine trifluoroacetate.

Using Deprotection Method A, the title compound was prepared from 1-(4-methoxybenzyl-D-phenylglyciny1)-4-(1-Boc-3-pyrrolidinyl)piperazine.

¹H NMR (CD₃OD)

Example 274.

20 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[2-(4-pyridinyl)-ethyl]piperazine

1-Boc-4-[2-(4-pyridinyl)ethyl]piperazine

1-Boc-piperazine (4.0 g, 21.5 mmol), 4-vinylpyridine (2.94 g, 27.9 mmol), and acetic acid (1.29 g, 21.5 mmol) were mixed in ethanol and heated to reflux for 48 h. The mixture was cooled to room temperature and concentrated under vacuum to provide 1-Boc-4-[2-(4-pyridinyl)ethyl]-piperazine (2.9 g, 45%) as an off white solid. The product was used without further purification.

¹H-NMR(CDCl₃)

CI-MS, m/e = 292 (M+1)

1-[2-(4-Pyridinyl)ethyl]piperazine hydrochloride

(Deprotection Method B) 1-Boc-4-[2-(4-pyridinyl)ethyl]-piperazine (1.0 g, 3.43 mmol) was dissolved in ethyl ether.

- 5 Ethyl acetate (15 mL) saturated with HCl was added, and the mixture stirred for 30 min at room temperature. The mixture was concentrated under vacuum and provided 1-[2-(4-pyridinyl)ethyl]piperazine hydrochloride (900 mg, 87%) as a tan solid.

10 ¹H-NMR(CD₃OD)

CI-MS, m/e = 192 (M+1)

1-(D-Boc-phenylglyciny)-4-[2-(4-pyridinyl)ethyl]piperazine

Using Coupling Method A, the title compound was prepared

- 15 from 1-[2-(4-pyridinyl)ethyl]piperazine and Boc-D-phenylglycine (95%).

¹H-NMR(CDCl₃)

CI-MS, m/e = 425 (M+1)

- 20 **1-(D-Phenylglyciny)-4-[2-(4-pyridinyl)ethyl]piperazine hydrochloride**

Using Deprotection Method B, the title compound was prepared from 1-(D-Boc-phenylglyciny)-4-[2-(4-pyridinyl)ethyl]-piperazine (89%).

- 25 ¹H-NMR(CD₃OD)

CI-MS, m/e = 325 (M+1)

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-[2-(4-pyridinyl)-ethyl]piperazine

- 30 Using Acylation Method B, the title compound was prepared from 1-(D-phenylglyciny)-4-[2-(4-pyridinyl)ethyl]piperazine hydrochloride and *p*-anisoyl chloride (70%).

¹H-NMR(CDCl₃)

CI-MS, m/e = 459 (M+1)

HPLC Analysis (Method E): 99.7%, RT=10.98 min.

5 **Examples 275 to 276**

Using Alkylation Method B, the title compounds were prepared from 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-(3-pyrrolidinyl)piperazine trifluoroacetate and the indicated aldehyde or ketone.

10

Example 275.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(1-methylpyrrolidin-3-yl)piperazine

Prepared from paraformaldehyde (20%).

15 ¹H-NMR(CDCl₃)

Example 276.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(1-isopropylpyrrolidin-3-yl)piperazine.

20 Prepared from acetone (59%).

¹H-NMR(CDCl₃)

The following analytical methods apply to Examples 277-336.

25 Analytical RPHPLC Method 1 = Vydac C18; linear gradient of 90/10 - 50/50 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min.

30 Analytical RPHPLC Method 2 = Vydac C18, linear gradient of 85/20 - 40/60 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min.

Examples 277 to 290

Unless otherwise indicated, the products of Examples 277 through 290 were obtained from the indicated acid and 1-D-phenylglyciny-1'-methyl-4,4'-bispiperidine using the procedure described in Example 277 (Coupling Method B).

Example 277.

1-(2-Chloropyridine-5-carbonyl-D-phenylglyciny-1'-methyl-4,4'-bispiperidine

(Coupling Method B) To a stirring solution of 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.20 g, 1.0 mmol) and 1-hydroxybenzotriazole hydrate (0.15 g, 1.1 mmol) in DMF (3 mL) was added 2-chloropyridine-5-carboxylic acid (0.14 g, 0.89 mmol) followed by a solution of 1-D-phenylglyciny-1'-methyl-4,4'-bispiperidine (0.25 g, 0.80 mmol) in DMF (2 mL). After stirring for 18 h, the solvent was removed in vacuo and the residue was partitioned between dichloromethane and 1 N sodium hydroxide. The aqueous phase was separated, extracted twice with dichloromethane, and the combined organic phases were dried with MgSO_4 , filtered and concentrated in vacuo. The resulting solid was dissolved in a minimum amount of dichloromethane and chromatographed over silica gel, eluting with 10% methanol (containing 2 N ammonia) in dichloromethane through 15% methanol (containing 2 N ammonia) in dichloromethane. The product containing fractions were combined and concentrated in vacuo to give 0.258 g (71%) of a white solid.

¹H-NMR

IS-MS, m/e 455.0 (M+1)

Analysis for $\text{C}_{25}\text{H}_{31}\text{N}_4\text{O}_2\text{Cl}\cdot 0.4\text{H}_2\text{O}$:

Calcd: C, 64.96; H, 6.93; N, 12.13;

Found: C, 64.68; H, 6.72; N, 12.02.

Analytical RPHPLC, Method 1, RT = 21.28 min (98%)

Example 278.

- 5 1-(5-Chloropyridine-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

Prepared from 2-chloropyridine-5-carboxylic acid (61%).

¹H-NMR

IS-MS, m/e 454.9 (M+1)

- 10 Analysis for C₂₅H₃₁N₄O₂Cl·0.4H₂O:

Calcd: C, 64.96; H, 6.93; N, 12.12;

Found: C, 64.75; H, 6.64; N, 12.00.

Analytical RPHPLC, Method 1, RT = 27.23 min (100%)

- 15 Example 279.

1-(3-Cyano-4-fluorobenzoyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

Prepared from 3-cyano-4-fluorobenzoic acid (66%).

¹H-NMR

- 20 IS-MS, m/e 463.0 (M+1)

Analysis for C₂₇H₃₁N₄O₂F·0.3H₂O:

Calcd: C, 69.30; H, 6.81; N, 11.97;

Found: C, 68.91; H, 6.58; N, 11.77.

Analytical RPHPLC [Vydac C18, linear gradient of 85/15 -

- 25 45/55 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min] RT = 21.54 (99%).

Example 280.

1-(5-Chlorobenzo[b]thiophene-2-carbonyl-D-phenylglyciny1)-

- 30 1'-methyl-4,4'-bispiperidine

Prepared from 5-chlorobenzo[b]thiophene-2-carboxylic acid (38%).

¹H-NMR

IS-MS, m/e 509.9 (M+1)

Analysis for C₂₈H₃₂N₃O₂SCl·0.3H₂O:

Calcd: C, 65.24; H, 6.37; N, 8.15;

5 Found: C, 65.01; H, 6.12; N, 8.07.

Analytical RPHPLC, Method 1, RT = 36.08 min (99%)

Example 281.

10 1-(2-Benzo[b]thiophenecarbonyl-D-phenylglyciny)-1'-methyl-
4,4'-bispiperidine

Prepared from 2-benzo[b]thiophenecarboxylic acid (82%).

¹H-NMR,

IS-MS, m/e 475.9 (M+1)

Analysis for C₂₈H₃₃N₃O₂S·0.4H₂O:

15 Calcd: C, 69.65; H, 7.06; N, 8.70;

Found: C, 69.45; H, 6.90; N, 8.58.

Analytical RPHPLC, Method 2, RT = 22.30 min (100%)

Example 282.

20 1-(6-Chlorobenzo[b]thiophene-2-carbonyl-D-phenylglyciny)-
1'-methyl-4,4'-bispiperidine

Prepared from 6-chlorobenzo[b]thiophene-2-carboxylic
acid (77%).

¹H-NMR

25 IS-MS, m/e 509.9 (M+1)

Analysis for C₂₈H₃₂N₃O₂SCl·0.3H₂O:

Calcd: C, 65.24; H, 6.37; N, 8.15;

Found: C, 64.97; H, 6.23; N, 8.07.

Analytical RPHPLC, Method 2, RT = 27.62 min (100%)

Example 283.

1- (Indole-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-
bispiperidine

Prepared from 2-indolecarboxylic acid (57%).

5 1H-NMR

IS-MS, m/e 459.0 (M+1)

Analysis for $C_{28}H_{34}N_4O_2 \cdot 0.4H_2O$:

Calcd: C, 71.10; H, 7.59; N, 11.85;

Found: C, 70.82; H, 7.25; N, 11.74.

10 Analytical RPHPLC, Method 1, RT = 29.60 min (99%)

Example 284.

1- (1-Methylindole-2-carbonyl-D-phenylglyciny1)-1'-methyl-
4,4'-bispiperidine

15 Prepared from 1-methylindole-2-carboxylic acid (43%).

1H-NMR

IS-MS, m/e 473.0 (M+1)

Analytical RPHPLC, Method 2, RT = 22.20 min (98%).

20 **Example 285.**

1- (Benzofuran-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-
bispiperidine

Prepared from 2-benzofurancarboxylic acid (50%).

1H-NMR

25 IS-MS, m/e 460.0 (M+1)

Analytical RPHPLC, Method 1, RT = 27.59 min (100%)

Example 286.

1- (3-Methylbenzofuran-2-carbonyl-D-phenylglyciny1)-1'-
30 methyl-4,4'-bispiperidine

Prepared from 3-methylbenzofuran-2-carboxylic acid (47%).

1H-NMR

IS-MS, m/e 474.1 (M+1)

Analytical RPHPLC, Method 1, RT = 31.31 min (95%)

Example 287.

- 5 1-(5-Methylbenzofuran-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

Prepared from 5-methylbenzofuran-2-carboxylic acid (45%).

¹H-NMR

IS-MS, m/e 474.3 (M+1)

- 10 Analytical RPHPLC, Method 1, RT = 30.91 min (100%)

Example 288.

1-(6-Methoxybenzofuran-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

- 15 Prepared from 6-methoxybenzofuran-2-carboxylic acid (50%).

¹H-NMR

IS-MS, m/e 490.0 (M+1)

Analytical RPHPLC, Method 1, RT = 29.26 min (100%)

- 20 Example 289.

1-(5-Chlorobenzofuran-2-carbonyl-D-phenylglyciny1)-1'-methyl-4,4'-bispiperidine

Prepared from 5-chlorobenzofuran-2-carboxylic acid (59%).

¹H-NMR

- 25 IS-MS, m/e 493.9 (M+1)

Analysis for C₂₈H₃₂N₃O₃Cl·0.5H₂O:

Calcd: C, 66.85; H, 6.61; N, 8.35;

Found: C, 66.46; H, 6.28; N, 8.25.

Analytical RPHPLC, Method 1, RT = 34.86 min (100%)

Example 290.

1-(2-Aminobenzimidazole-5-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

Prepared from 2-amino-5-carboxybenzimidazole hydrochloride
5 (32%).

¹H-NMR

IS-MS, m/e 475.2 (M+1)

Analytical RPHPLC [Vydac C18, linear gradient of 98/2 -
58/42 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40
10 min, 1 mL/min] RT = 24.56 (90%).

Example 291. 1-(3-Aminobenzisoxazole-5-carbonyl-D-phenylglycine)-1'-methyl-4,4'-bispiperidine

To a stirring solution of acetoxime (98 mg, 7.1 mmol) in DMF
15 (5 mL) was added a 1 M solution of potassium tert-butoxide
(1.3 mL, 1.3 mmol) in THF. After 2 min, 1-(3-cyano-4-fluorobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine
(303 mg, 0.65 mmol) was added; and, after another hour, the
solvent was partially removed and the residue was
20 partitioned between brine and dichloromethane. The layers
were separated and the aqueous phase was extracted another
two times with dichloromethane. The combined organics were
dried (MgSO₄), filtered and concentrated in vacuo.

IS-MS, m/e 516.0 (M+1)

25

The residue was then dissolved in ethanol (3.6 mL) and 1 N
HCl was added. The stirring solution was heated to reflux.

After 5 h, the heating mantle was removed and after
cooling, the solution was diluted ethyl acetate and water.

30 The pH of the aqueous phase was adjusted to 11 with 2 N
sodium hydroxide and extracted twice with dichloromethane.
The combined dichloromethane extracts were dried (MgSO₄),

filtered and concentrated in vacuo. The resulting solid was dissolved in a minimum amount of dichloromethane and chromatographed over silica gel, eluting with 2% methanol (containing 2 N ammonia) in dichloromethane through 10% methanol (containing 2 N ammonia) in dichloromethane. The product containing fractions were combined and concentrated in vacuo to give 89 mg (29%) of an off-white solid.

¹H-NMR

IS-MS, m/e 476.3 (M+1)

10 Analytical RPHPLC, Method 1, RT = 19.55 min (99%)

Examples 292 to 303

Preparation of Starting Materials

15 1-(Boc-D-phenylglyciny1)-4-hydroxypiperidine

(Coupling Method C) To a stirring solution of 1-hydroxy-7-azabenzotriazole (10.24 g, 75.2 mmol) and EDCI (14.42 g, 75.2 mmol) in DMF (160 mL) was added a solution of Boc-D-phenylglycine (18.9 g, 75.2 mmol) in DMF (80 mL). After 10 min, 4-hydroxypiperidine (6.85 g, 67.7 mmol) was added. After stirring over night, the solvent was evaporated in vacuo and the residue was partitioned between ethyl acetate and water. The organic phase separated and washed with saturated aqueous NaHCO₃, followed by brine, dried over MgSO₄, flitered and concentrated in vacuo. Two-thirds of this material was dissolved in a minimum amount of dichloromethane and chromatographed over silica gel, eluting with a gradient of dichloromethane through 1:1 dichloromethane/ethyl acetate. The product containing fractions were combined and concentrated in vacuo to give 15.71 g (94%) of a white foam.

¹H-NMR

IS-MS, m/e 335.1 (M+1)

Analysis for $C_{18}H_{26}N_2O_4$:

Calcd: C, 64.65; H, 7.84; N, 8.37;

Found: C, 64.40; H, 7.77; N, 8.12.

5

1-(D-phenylglyciny1)-4-hydroxypiperidine

(Deprotection Method D) To a stirring solution of 1-(Boc-D-phenylglyciny1)-4-hydroxypiperidine (5 g, 15 mmol) in dichloromethane (290 mL) was added anisole, (8 mL) followed
10 by trifluoroacetic acid (29 mL). After stirring for 4 h, the solvent was concentrated in vacuo and the residue was suspended with stirring in diethyl ether. After 1 h, the mixture was filtered and the solid was partitioned between ethyl acetate and saturated aqueous $NaHCO_3$. The organic
15 phase was washed with brine, dried with $MgSO_4$, filtered and concentrated to give 0.41 g of white solid. The combined aqueous phase was back extracted with 3:1 chloroform/-isopropanol and this organic phase was separated, dried with $MgSO_4$, filtered and concentrated in vacuo to give 1.6 g of
20 white solid. The two crops of solid were combined to give 2.02 g (90%) of the title compound.

1H -NMR

IS-MS, m/e 235.1 (M+1)

25 **1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-hydroxypiperidine**

To a stirring solution of 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.4 g, 7.4 mmol), 1-hydroxybenzotriazole hydrate (1.0 g, 7.4 mmol) and N,N-diisopropylethylamine (1.4 mL) in DMF (20 mL) was added
30 a solution of 1-(D-phenylglyciny1)-4-hydroxypiperidine (2.0 g, 7.38 mmol) in DMF (10 mL) followed by a solution of 4-methoxybenzoic acid (1.0 g, 6.7 mmol) in DMF (10 mL).

After stirring overnight at room temperature, the solvent was removed in vacuo and the residue was partitioned between ethyl acetate and water. The organic phase was washed again with water followed by saturated aqueous NaHCO_3 (2X) and
5 brine, then dried with MgSO_4 , filtered and concentrated in vacuo to give 2.4 g of off-white solid. A portion of this material (2.0 g) was dissolved in a minimal amount of dichloromethane and chromatographed over silica gel, eluting
10 acetate/dichloromethane. The product-containing fractions were combined and concentrated in vacuo to give 1.3 g (60%) of a white foam.

$^1\text{H-NMR}$

IS-MS, m/e 369.2 (M+1)

15 Analysis for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$:

Calcd: C, 68.46; H, 6.57; N, 7.60;

Found: C, 67.88; H, 6.73; N, 7.33.

Analytical RPHPLC, Method 1, RT = 24.24 min (100%)

20 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-oxopiperidine
(Oxidation Method B) To a stirring solution of oxalyl chloride (0.26 mL, 3 mmol) in dichloromethane (6.5 mL) at -50 °C, was added a solution of DMSO (0.43 mL, 6 mmol) in dichloromethane (1.3 mL). After 3 min, a solution of
25 1-(4-methoxybenzoyl-D-phenylglyciny1)-4-hydroxypiperidine (1.0 g, 2.7 mmol) in dichloromethane (4 mL) was added and the solution was allowed to warm to -20 °C over 45 min. Triethylamine (2 mL) was then added and the solution was allowed to warm to room temperature. The solution was then
30 diluted with dichloromethane and water and the layers were separated. The organic phase was washed with brine, dried over MgSO_4 , filtered and concentrated in vacuo. The residue

was dissolved in a minimum amount of dichloromethane and chromatographed over silica gel, eluting with a gradient of dichloromethane through 50% ethyl acetate/dichloromethane. The product containing fractions were combined and

5 concentrated in vacuo to give 0.77 g (78%) of a white foam.

¹H-NMR

IS-MS, m/e 367.2 (M+1)

Analysis for C₂₁H₂₂N₂O₄:

Calcd: C, 68.84; H, 6.05; N, 7.65;

10 Found: C, 68.33; H, 6.01; N, 7.27.

Analytical RPHPLC, Method 1, RT = 25.52 min (100%)

General Procedure: Unless otherwise indicated, the product of Examples 292-303 was obtained from 1-(4-methoxybenzoyl-D-phenylglyciny)-4-oxopiperidine and the indicated amine using the alkylation procedure described for Example 292 (Alkylation Method C).

Example 292.

20 1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(1-pyrrolidinyl)-piperidine

(Alkylation Method C) To a stirring solution of

1-(4-methoxybenzoyl-D-phenylglyciny)-4-oxopiperidine (50 mg, 0.14 mmol) and pyrrolidine (0.011 mL, 0.13 mmol) in

25 1,2-dichloroethane (1 mL) was added sodium triacetoxymethylborohydride (45 mg, 0.21 mmol). After stirring overnight, the mixture was loaded onto an SCX column (pretreated with a 5% glacial acetic acid in methanol solution), rinsed with methanol (2 column volumes) and eluted with a 30% 2 N
30 ammonia/methanol in dichloromethane solution. The solution was concentrated in vacuo. The product containing fractions

were combined and concentrated in vacuo to give 48 mg (87%) of the title compound.

¹H-NMR

IS-MS, m/e 422.0 (M+1)

5 Analytical RPHPLC, Method 1, RT = 21.02 min (100%)

Example 293.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(1-piperidinyl)-
piperidine

10 Prepared from piperidine (49%).

¹H-NMR

IS-MS, m/e 436.0 (M+1)

Analytical RPHPLC, Method 1, RT = 22.14 min (100%)

15 Example 294.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(4-methylpiperidin-
1-yl)piperidine

Prepared from 4-methylpiperidine (78%).

¹H-NMR

20 IS-MS, m/e 450.0 (M+1)

Analytical RPHPLC, Method 1, RT = 24.06 min (100%)

Example 295.

1-(4-Methoxybenzoyl-D-phenylglyciny)-4-(4-methylpiperazin-
25 1-yl)piperidine

Prepared from 1-methylpiperazine (98%).

¹H-NMR

IS-MS, m/e 451.0 (M+1)

Analytical RPHPLC, Method 1, RT = 18.66 min (99%)

Example 296.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-ethylpiperazin-1-yl)piperidine

Prepared from 1-ethylpiperazine (76%).

5 ¹H-NMR

IS-MS, m/e 465.0 (M+1)

Analytical RPHPLC, Method 1, RT = 19.11 min (100%)

Example 297.

10 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-isopropylpiperazin-1-yl)piperidine

Prepared from 1-isopropylpiperazine (83%).

¹H-NMR

IS-MS, m/e 479.2 (M+1)

15 Analysis for C₂₈H₃₈N₄O₃·0.3H₂O:

Calcd: C, 69.48; H, 8.04; N, 11.58;

Found: C, 69.22; H, 7.91; N, 11.34.

Analytical RPHPLC, Method 1, RT = 19.56 min (99%)

20 Example 298.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(hexahydro-1,4-diazapin-1-yl)piperidine hydrochloride

¹H-NMR

IS-MS, m/e 451.0 (M+1)

25 Analytical RPHPLC, Method 1, RT = 16.86 min (100%)

Example 299.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[4-methyl-(hexahydro-1,4-diazapin-1-yl)]piperidine

30 Prepared from 4-methyl-hexahydro-1,4-diazapine (63%).

¹H-NMR

IS-MS, m/e 465.0 (M+1)

Analytical RPHPLC, Method 1, RT = 18.86 min (98%)

Example 300.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(3-pyridylamino)-
5 piperidine

Prepared from 3-aminopyridine (25%).

¹H-NMR

IS-MS, m/e 445.0 (M+1)

Analytical RPHPLC, Method 1, RT = 23.87 min (100%)

10

Example 301.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[(N-methyl-N-
benzyl)amino]piperidine

Prepared from N-methylbenzylamine (89%).

15 ¹H-NMR

IS-MS, m/e 472.0 (M+1)

Analysis for C₂₉H₃₃N₃O₃·0.1H₂O:

Calcd: C, 73.58; H, 7.07; N, 8.88;

Found: C, 73.39; H, 7.19; N, 9.06.

20 Analytical RPHPLC, Method 1, RT = 26.27 min (98%)

Example 302.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[(3-pyridylmethyl)-
amino]piperidine

25 Prepared from 3-aminomethylpyridine (72%).

¹H-NMR

IS-MS, m/e 459.0 (M+1)

Analysis for C₂₇H₃₀N₄O₃·0.2H₂O:

Calcd: C, 70.17; H, 6.63; N, 12.12;

30 Found: C, 70.00; H, 6.53; N, 12.13.

Analytical RPHPLC, Method 1, RT = 16.38 min (100%)

Example 303.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[(4-pyridylmethyl)-aminol]piperidine

prepared from 4-aminomethylpyridine (46%).

5 ¹H-NMR

IS-MS, m/e 459.0 (M+1)

Analysis for C₂₇H₃₀N₄O₃·0.9H₂O:

Calcd: C, 68.30; H, 6.75; N, 11.80;

Found: C, 67.99; H, 6.42; N, 11.59.

10 Analytical RPHPLC, Method 1, RT = 18.36 min (100%)

Examples 304 to 314

General Procedure: Unless otherwise indicated, the product of Examples 304-314 was obtained from 1-(4-methoxybenzoyl-D-phenylglyciny1)piperazine and the indicated aldehyde or ketone using the alkylation procedure described for Example 304 (Alkylation Method D).

Example 304.

20 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(2-pyridylmethyl)-piperazine

(Alkylation Method D) To a stirring solution of

1-(4-methoxybenzoyl-D-phenylglyciny1)piperazine (50 mg, 0.14 mmol) and 2-pyridinecarboxaldehyde (0.020 mL, 23 mg, 0.21

25 mmol) in 5% acetic acid/methanol (1 mL) was added NaBH₃CN (20 mg, 0.32 mmol). After 4 h, the solution was loaded onto an SCX column (pretreated with a 5% glacial acetic acid in methanol solution), rinsed with methanol (2 column volumes) and eluted with a 30% 2N ammonia/methanol in dichloromethane solution. The solution was concentrated in vacuo and the residue was dissolved in a minimum amount of dichloromethane and chromatographed over silica gel, eluting with

30

dichloromethane, followed by 50% ethyl acetate/dichloromethane, and finally with a gradient of 2%-10% (2 N NH₃ in MeOH) in dichloromethane. The product containing fractions were combined and concentrated in vacuo to give 30 mg (48%) of the title compound.

¹H-NMR

IS-MS, m/e 444.9 (M+1)

Analytical RPHPLC, Method 1, RT = 21.70 min (100%)

10 Example 305.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(3-pyridylmethyl)-piperazine

Prepared from 3-pyridine carboxaldehyde (42%).

¹H-NMR

15 IS-MS, m/e 444.9 (M+1)

Analytical RPHPLC, Method 1, RT = 17.84 min (99%).

Example 306.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(4-pyridylmethyl)-piperazine

Prepared from 4-pyridine carboxaldehyde (45%).

¹H-NMR

IS-MS, m/e 444.9 (M+1)

Analytical RPHPLC, Method 1, RT = 18.36 min (99%)

25

Example 307.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-phenethylpiperazine

Prepared from phenylacetaldehyde (34%).

¹H-NMR

30 IS-MS, m/e 458.0 (M+1)

Analytical RPHPLC, Method 1, RT = 27.44 min (100%)

Example 308.

1- (4-Methoxybenzoyl-D-phenylglyciny1)-4- (3-pentyl)piperazine

Prepared from 3-pentanone (88%).

¹H-NMR

5 IS-MS, m/e 424.0 (M+1)

Analytical RPHPLC, Method 1, RT = 23.62 min (100%)

Example 309.

1- (4-Methoxybenzoyl-D-phenylglyciny1)-4-cyclopentyl-

10 piperazine

Prepared from cyclopentanone (95%).

¹H-NMR

IS-MS, m/e 422.0 (M+1)

Analytical RPHPLC, Method 1, RT = 20.76 min (100%)

15

Example 310.

1- (4-Methoxybenzoyl-D-phenylglyciny1)-4- (4-methyl-
cyclohexyl)piperazine

Prepared from 4-methylcyclohexanone (46%).

20 ¹H-NMR

IS-MS, m/e 450.0 (M+1)

Analytical RPHPLC, Method 1, RT = 27.07 min (isomer 1),
27.74 min (isomer 2).

25 Example 311.

1- (4-Methoxybenzoyl-D-phenylglyciny1)-4- (tetrahydro-
thiopyran-4-yl)piperazine

Prepared from tetrahydro-4H-thiopyran-4-one (86%).

¹H-NMR

30 IS-MS, m/e 453.9 (M+1)

Analytical RPHPLC, Method 1, RT = 22.96 min (100%)

Example 312.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(2-indany1)-
piperazine

Prepared from 2-indanone (92%).

5 ¹H-NMR

IS-MS, m/e 469.9 (M+1)

Analytical RPHPLC, Method 1, RT = 26.32 min (100%)

Example 313.

10 1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-benzylpiperazine

Prepared from benzaldehyde (87%).

¹H-NMR

IS-MS, m/e 444.0 (M+1)

Analytical RPHPLC, Method 1, RT = 25.78 min (96%)

15

Example 314.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(cyclohexyl-
methyl)piperazine

Prepared from cyclohexanecarboxaldehyde (86%).

20 ¹H-NMR

IS-MS, m/e 450.2 (M+1)

Analytical RPHPLC, Method 1, RT = 28.07 min (94%)

Examples 315 to 316

25 Preparation of Starting Materials

1-(Boc-D-Phenylglyciny1)-4-oxopiperidine

Using Oxidation Method B, the title compound was prepared
from 1-(Boc-D-phenylglyciny1)-4-hydroxypiperidine (44%).

30 ¹H-NMR

IS-MS, m/e 333.0 (M+1)

1-(Boc-D-Phenylglyciny1)-4-(4-methylpiperazin-1-yl)-
piperidine

Using Alkylation Method C, the title compound was prepared
from 1-(Boc-D-phenylglyciny1)-4-oxopiperidine and

5 methylpiperazine (65%).

¹H-NMR

IS-MS, m/e 417.3 (M+1)

Analysis for C₂₃H₃₆N₄O₃:

Calcd: C, 66.32; H, 8.71; N, 13.45;

10 Found: C, 66.25; H, 8.58; N, 13.42.

~~1-D-Phenylglyciny1-4-(4-methylpiperazin-1-yl)piperidine~~

HCl gas was bubbled through a stirring solution of 1-(Boc-D-
phenylglyciny1)-4-(4-methylpiperazin-1-yl)piperidine (1.36

15 g, 3.26 mmol) in ethyl acetate (150 mL). A white

precipitate was formed immediately, but then went back into
solution. After about 5 min, a white precipitate again fell
out of solution. After 10 min, the addition of HCl was

discontinued and after stirring for a total of 1 h, the
20 mixture was filtered to give 1.38 g (quantitative) of white
solid.

¹H-NMR

IS-MS, m/e 317.3 (M+1)

Analysis for C₁₈H₂₈N₄O·2.9HCl·2.5H₂O:

25 Calcd: C, 46.27; H, 7.74; N, 11.99; Cl, 22.01;

Found: C, 46.06; H, 7.51; N, 11.63; Cl, 21.78.

General Procedure: The product of Examples 315-316 was
prepared from 1-(D-phenylglyciny1)-4-(4-methylpiperazin-
30 1-yl)piperidine and the indicated acid using Coupling
Method B.

Example 315.

1-(Indole-6-carbonyl-D-phenylglyciny1)-4-(4-methylpiperazin-1-yl)piperidine

Prepared from indole-6-carboxylic acid (66%).

5 ¹H-NMR

IS-MS, m/e 460.2 (M+1)

Analytical RPHPLC, Method 1, RT = 17.83 min (99%)

Example 316.

10 1-(3-Chloroindole-6-carbonyl-D-phenylglyciny1)-4-(4-methylpiperazinyl)piperidine

Prepared from 3-chloroindole-6-carboxylic acid (69%)..

¹H-NMR

IS-MS, m/e 494.3 (M+1)

15 Analytical RPHPLC, Method 1, RT = 22.99 min (99%)

Examples 317 to 320

Preparation of Starting Materials

20 (Cbz-D-phenylglyciny1)piperazine.

Using Deprotection Method D, the title compound was prepared from 1-(Cbz-D-phenylglyciny1)-4-Boc-piperazine (85%)

¹H-NMR

IS-MS, m/e 354.2 (M+1)

25 Analysis for C₂₀H₂₃N₃O₃·0.2H₂O:

Calcd: C, 67.28; H, 6.61; N, 11.77;

Found: C, 67.10; H, 6.46; N, 11.63.

1-(Cbz-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)-

30 piperazine

Using Alkylation Method C, the title compound was prepared from (Cbz-D-phenylglyciny1)piperazine and 1-methylpiperidin-

4-one (49%). The product was purified using silica gel chromatography, eluting with a gradient of dichloromethane through 10% (2 N ammonia in methanol) / dichloromethane.

¹H-NMR

5 IS-MS, m/e 451.3 (M+1)

Analysis for C₂₆H₃₄N₄O₃:

Calcd: C, 69.31; H, 7.61; N, 12.43;

Found: C, 69.36; H, 7.71; N, 13.14.

10 1-D-Phenylglyciny1-4-(1-methylpiperidin-4-yl)piperazine dihydrochloride.

To a stirring suspension of 5% Pd/C (0.6 g) in ethanol (25 mL) under nitrogen was added a solution of 1-(Cbz-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine (2.6 g, 15 5.77 mmol) and acetic acid (1.6 mL) in ethanol (50 mL). The flask was placed under vacuum and the atmosphere was replaced with hydrogen (balloon). After 4 h, diatomaceous earth was added and the mixture was filtered through a pad of diatomaceous earth and concentrated in vacuo. The 20 residue was dissolved in ethyl acetate and HCl gas was bubbled through the stirring solution to precipitate the dihydrochloride salt. The mixture was filtered and the solid was dried in vacuo to give 2.6 g (quantitative) of the title compound.

25 ¹H-NMR

IS-MS, m/e 317.3 (M+1)

General Procedure: The product of Examples 317-320 was prepared from 1-(D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine dihydrochloride and the indicated acid using 30 Coupling Method B.

Example 317.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine

Prepared from 4-methoxybenzoic acid (19%).

5 ¹H-NMR

IS-MS, m/e 451.0 (M+1)

Analytical RPHPLC, Method 1, RT = 16.76 min (100%)

Example 318.

10 1-(Indole-6-carbonyl-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine

Prepared from indole-6-carboxylic acid (65%).

¹H-NMR

IS-MS, m/e 460.2 (M+1)

15 Analytical RPHPLC, Method 1, RT = 16.68 min (100%)

Example 319.

1-(3-Methylindole-6-carbonyl-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine

20 Prepared from 3-methylindole-6-carboxylic acid (50%).

¹H-NMR

IS-MS, m/e 474.3 (M+1)

Analytical RPHPLC, Method 1, RT = 22.20 min (98%)

25 Example 320.

1-(3-Chloroindole-6-carbonyl-D-phenylglyciny1)-4-(1-methylpiperidin-4-yl)piperazine

Prepared from 3-chloroindole-6-carboxylic acid (76%).

¹H-NMR

30 IS-MS, m/e 493.9 (M+1)

Analytical RPHPLC, Method 1, RT = 22.66 min (100%)

Examples 321 to 324

Preparation of Starting Materials

Ethyl hydroxyimino-pyridine-2-acetate

5 To a stirring solution of ethyl pyridine-2-acetate (12.6 g, 76.3 mmol) in acetic acid (19 mL) at 5 °C was added a solution of sodium nitrite (6.05 g, 87.7 mmol) in water (12 mL) at a rate sufficient to maintain the internal temperature below 15 °C. After complete addition and an
10 additional 30 min, an additional 30 mL of water was added. The resulting white precipitate was filtered, washed with water, saturated aqueous NaHCO₃, and again with water. The solid was then dried under vacuum to give 14.1 g (95%) of the title compound.

15 ¹H-NMR

IS-MS, m/e 194.9 (M+1)

Analysis for C₉H₁₀N₂O₃:

Calcd: C, 55.67; H, 5.19; N, 14.43;

Found: C, 55.79; H, 5.14; N, 14.13.

20

Boc-D,L-(2-Pyridinyl)glycine ethyl ester

To a solution of ethyl hydroxyimino-pyridine-2-acetate (7.8 g, 40.15 g) in ethanol (175 mL) and glacial acetic acid (20 mL) was added 5% Pd/C, and the mixture was shaken in a
25 hydrogenation apparatus under an atmosphere of hydrogen at 3.1 bar for 4 h. The mixture was filtered through diatomaceous earth and concentrated in vacuo. The residue was dissolved in THF/H₂O (1:1, 240 mL) and treated with di-tert-butyl dicarbonate (14.23 g, 65.2 mmol) and sodium
30 bicarbonate (27.4 g, 326 mmol). After stirring at room temperature for 2 h, the solution was concentrated in vacuo and the residue was partitioned between EtOAc and water.

The organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated in vacuo. The crude material was purified via chromatography over silica gel, eluting with a stepwise gradient of 10-20% ethyl acetate in dichloromethane, to give 8.11 g (72%) of a yellow oil.

¹H-NMR

IS-MS, m/e 281.1 (M+1)

10 1- [Boc-D,L-(2-Pyridinyl)glyciny]l-1'-methyl-4,4'-
bispiperidine

To a stirring solution of Boc-D,L-(2-pyridinyl)glycine ethyl ester (3.89 g, 13.88 mmol) in 1, 4-dioxane (20 mL) was added a solution of lithium hydroxide hydrate (0.64 g, 15.27 mmol) in water (20 mL). After stirring for 2 h, the solution was concentrated in vacuo. The residue was dried under vacuum for 15 h then dissolved in DMF (50 mL). The solution was cooled to 0 °C, purged with nitrogen, and diethyl cyanophosphonate (2.5 g, 16.66 mmol) was slowly added.

20 After 2 min, the solution was treated with a solution of 1-methyl-4,4'-bispiperidine dihydrochloride (3.9 g, 15.27 mmol) and triethylamine (6.8 mL, 48.58 mmol) in DMF (50 mL).

After 2 h, the cold bath was removed and the solution was allowed to stir overnight. The next morning, the solvent was evaporated in vacuo and the resulting oil was partitioned between 3:1 chloroform:isopropyl alcohol and saturated aqueous sodium bicarbonate. The organic phase was dried over magnesium sulfate, filtered and concentrated in vacuo. The crude material was purified via chromatography over silica gel, eluting with a stepwise gradient of 5-9% (2 N ammonia in methanol) in dichloromethane to give 2.6 g (45%) of a clear oil.

¹H-NMR

IS-MS, m/e 417.2 (M+1)

1-[D,L-(2-Pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine

5 (Deprotection Method E) To a stirring solution of 1-[Boc-D,L-(2-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine (1.8 g, 4.32 mmol) in dichloromethane (90 mL) was added anisole (2.3 mL, 21.6 mmol), followed by trifluoroacetic acid (8.3 mL, 108 mmol). After 4 h, the solvents were evaporated in
10 vacuo, the crude product was dissolved in methanol and loaded onto an SCX column (pretreated with a 5% glacial acetic acid in methanol solution), rinsed with methanol (2 column volumes) and eluted with a 30% 2 N ammonia/methanol in dichloromethane solution. The product containing
15 fractions were combined and concentrated in vacuo to give 1.08 g (77%) of a yellow oil.

¹H-NMR

IS-MS, m/e 317.2 (M+1)

Analysis for C₁₈H₂₈N₄O·0.55H₂O:

20 Calcd: C, 66.25; H, 8.99; N, 17.17;

Found: C, 66.07; H, 8.49; N, 16.66.

General Procedure: The product of Examples 321-324 was prepared from 1-[D,L-(2-pyridinyl)glycinyll]-1'-methyl-4,4'-
25 bispiperidine and the indicated acid using the procedure described for Example 321 (Coupling Method D).

Example 321.

1-[Indole-6-carbonyl-D,L-(2-pyridinyl)glycinyll]-1'-methyl-
30 4,4'-bispiperidine

(Coupling Method D) To a stirring solution of 1-[D,L-(2-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine (0.3 g,

0.95 mmol) in N, N-dimethylformamide (3 mL) was added
indole-6-carboxylic acid (0.15 g, 0.95 mmol) and 1-hydroxy-
benzotriazole hydrate (0.13 g, 0.95 mmol), followed by
1,3-dicyclohexylcarbodiimide (0.19 g, 0.95 mmol). After
5 stirring overnight, the mixture was filtered and the
filtrate was loaded onto an SCX column (pretreated with a 5%
glacial acetic acid in methanol solution), rinsed with
methanol (2 column volumes) and eluted with a 30% (2 N
ammonia in methanol) in dichloromethane solution. The
10 product containing fractions were concentrated in vacuo and
the residue was chromatographed over silica gel, eluting
with a stepwise gradient of 5-9% (2 N ammonia in methanol)
in dichloromethane to give 255 mg (58%) of a tan foam.

¹H-NMR

15 IS-MS, m/e 460.3 (M+1)

Analytical RPHPLC, Method 1, RT = 14.90 min (100%)

Example 322.

1- [4-Methoxybenzoyl-D,L-(2-pyridinyl)glyciny]l-1'-methyl-
20 4,4'-bispiperidine

Prepared from 4-methoxybenzoic acid (53%).

¹H-NMR

IS-MS, m/e 451.2 (M+1)

Analytical RPHPLC, Method 1, RT = 14.79 min (98%)

25

Example 323.

1- [3-Methylindol-6-carbonyl-D,L-(2-pyridinyl)glyciny]l-1'-
methyl-4,4'-bispiperidine

Prepared from 3-methyl-6-carboxyindole (40%).

30 ¹H-NMR

IS-MS, m/e 474.3 (M+1)

Analytical RPHPLC, Method 1, RT = 18.28 min (97%)

Example 324.

1-[3-Chloroindole-6-carbonyl-D,L-(2-pyridinyl)glyciny]-1'-methyl-4,4'-bispiperidine

5 Prepared from 3-chloro-6-carboxyindole (71%).

¹H-NMR

IS-MS, m/e 494.0 (M+1)

Analysis for C₂₇H₃₂N₅O₂Cl·0.2H₂O:

Calcd: C, 65.17; H, 6.56; N, 14.07;

10 Found: C, 65.57; H, 6.56; N, 13.23.

Analytical RPHPLC, Method 1, RT = 20.96 min (99%)

Examples 325 to 328**Preparation of Starting Materials**

15

Ethyl hydroxyimino-pyridine-3-acetate

Using the procedure of Tikk et al. [Acta. Chimica Hungarica, 114(3-4), 355], a mixture of ethyl hydroxyimino-pyridine-3-acetate and n-butyl hydroxyimino-pyridine-3-acetate was

20 prepared from ethyl pyridine-3-acetate and n-butyl nitrite.

¹H-NMR

IS-MS, m/e 195 (M+1), 223.1 (M+1)

Boc-D,L-(3-Pyridinyl)glycine ethyl ester

25 Using methods substantially equivalent to those described above in preparation of Boc-D,L-(2-pyridinyl)glycine ethyl ester, the title compound was prepared from the above ethyl hydroxyimino-pyridine-3-acetate (57%).

¹H-NMR

30 IS-MS, m/e 281.1 (M+1)

1- [Boc-D,L- (3-Pyridinyl)glycinyll]-1'-methyl-4,4'-
bispiperidine

Using methods substantially equivalent to those described in
preparation of 1- [Boc-D,L- (2-pyridinyl)glycinyll]-1'-methyl-
5 4,4'-bispiperidine, the title compound was prepared from
Boc-D,L- (3-pyridinyl)glycine ethyl ester (20%).

¹H-NMR

IS-MS, m/e 417.2 (M+1)

10 1- [D,L- (3-Pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine

Using methods substantially equivalent to those described in
preparation of 1- [D,L- (2-pyridinyl)glycinyll]-1'-methyl-4,4'-
bispiperidine, the title compound was prepared from
1- [Boc-D,L- (3-pyridinyl)glycinyll]-1'-methyl-4,4'-

15 bispiperidine (75%).

¹H-NMR

IS-MS, m/e 317.2 (M+1)

General Procedure: The product of Examples 325-328 was
20 prepared from 1- [D,L- (3-pyridinyl)glycinyll]-1'-methyl-4,4'-
bispiperidine and the indicated acid using the procedure
described for Example 325 (Coupling Method D).

Example 325.

25 1- [4-Methoxybenzoyl-D,L- (3-pyridinyl)glycinyll]-1'-methyl-
4,4'-bispiperidine

Prepared from 4-methoxybenzoic acid (45%).

¹H-NMR

IS-MS, m/e 451.2 (M+1)

30 Analysis for C₂₆H₃₄N₄O₃·1.2H₂O:

Calcd: C, 66.13; H, 7.77; N, 11.87;

Found: C, 66.61; H, 7.27; N, 11.87.

Analytical RPHPLC, Method 1, RT = 12.98 min (98%)

Example 326.

1-[Indole-6-carbonyl-D,L-(3-pyridinyl)glycinyll]-1'-methyl-
5 4,4'-bispiperidine

Prepared from indole-6-carboxylic acid (36%).

¹H-NMR

IS-MS, m/e 460.3 (M+1)

Analysis for C₂₇H₃₃N₅O₂·1.5H₂O:

10 Calcd: C, 66.64; H, 7.46; N, 14.39;
Found: C, 66.71; H, 6.87; N, 13.89.

Analytical RPHPLC, Method 1, RT = 14.39 min (100%)

Example 327.

15 1-[3-Methylindole-6-carbonyl-D,L-(3-pyridinyl)glycinyll]-1'-
methyl-4,4'-bispiperidine

Prepared from 3-methylindole-6-carboxylic acid (40%).

¹H-NMR

IS-MS, m/e 474.3 (M+1)

20 Analysis for C₂₈H₃₅N₅O₂·1.6H₂O:

Calcd: C, 66.93; H, 7.66; N, 13.94;
Found: C, 66.63; H, 6.99; N, 13.52.

Analytical RPHPLC, Method 1, RT = 16.98 min (98%)

25 Example 328.

1-[3-Chloroindole-6-carbonyl-D,L-(3-pyridinyl)glycinyll]-1'-
methyl-4,4'-bispiperidine

Prepared from 3-chloroindole-6-carboxylic acid (46%).

¹H-NMR

30 IS-MS, m/e 494.2 (M+1)

Analysis for C₂₇H₃₂ClN₅O₂·1.1H₂O:

Calcd: C, 63.11; H, 6.71; N, 13.63;

Found: C, 62.84; H, 6.32; N, 13.26.

Analytical RPHPLC, Method 1, RT = 19.63 min (100%)

Examples 329 to 330

5 Preparation of Starting Materials

Boc-D-[3-(ethanesulfonylamino)phenyl]glycine

To a stirring solution of D-3-(ethanesulfonylamino)-phenylglycine (20 g, 77.43 mmol) and sodium carbonate (8.2 g, 77.43 mmol) in 3:1 THF/water (200 mL) at 0 °C, was added di-tert-butyl dicarbonate (18.5 g, 85.17 mmol). After stirring for 30 min, the cold bath was removed; and after an additional 30 min at room temperature, the solvent was removed and the residue was partitioned between ethyl acetate and water. The aqueous layer was acidified to pH 2 with KHSO₄ and extracted twice with ethyl acetate. The combined ethyl acetate extracts were washed with water, dried with Na₂SO₄, filtered and concentrated in vacuo to give 17.51 g (63%) of a white solid.

20 ¹H-NMR

IS-MS, m/e 357.0 (M-1)

1-[Boc-D-[3-(ethanesulfonylamino)phenyl]glycinyll]-1'-methyl-4,4'-bispiperidine

25 To a stirring solution of Boc-D-[3-(ethanesulfonylamino)-phenyl]glycine (5 g, 13.95 mmol) in dichloromethane at 0 °C, diethyl cyanophosphonate (2.12 mL, 13.95 mmol) and diisopropylethylamine (4.86 mL, 27.91 mmol) and then N-methylbispiperidine dihydrobromide (4.32 g, 12.56 mmol) were added; and the mixture was stirred at 0 °C for 3 h. The reaction mixture was then stirred at room temperature overnight, filtered, washed with saturated aqueous sodium

bicarbonate and water, dried over sodium sulfate, filtered and concentrated in vacuo to give 5 g (76%) of a tan foam.

¹H-NMR

IS-MS, m/e (M+1)

5

1-[D-[3-(ethanesulfonylamino)phenyl]glycinyll]-1'-methyl-4,4'-bispiperidine

Using Deprotection Method E, the title compound was prepared from 1-[Boc-D-[3-(ethanesulfonylamino)phenyl]glycinyll]-1'-

10 methyl-4,4'-bispiperidine (74%).

¹H-NMR

IS-MS, m/e 423.1 (M+1)

Analysis for C₂₁H₃₄N₄O₃S·1.3H₂O:

Calcd: C, 56.55; H, 8.27; N, 12.56;

15 Found: C, 56.68; H, 7.87; N, 11.97.

General Procedure: The product of Examples 329-330 was prepared from 1-[D-[3-(ethanesulfonylamino)phenyl]glycinyll]-1'-methyl-4,4'-bispiperidine and the indicated acid using
20 the procedure described for Example 321 (Coupling Method D).

Example 329.

1-[4-Methoxybenzoyl-D-[3-(ethanesulfonylamino)-phenyl]glycinyll]-1'-methyl-4,4'-bispiperidine

25 Prepared from 4-methoxybenzoic acid (43%).

¹H-NMR

IS-MS, m/e 557.3 (M+1)

Analysis for C₂₉H₄₀N₄O₅S·0.9H₂O:

Calcd: C, 60.79; H, 7.35; N, 9.78;

30 Found: C, 60.49; H, 7.08; N, 9.62.

Analytical RPHPLC, Method 1, RT = 22.68 min (98%)

Example 330.

1-[Indole-6-carbonyl-D-[3-(ethanesulfonylamino)-phenyl]glycinyll]-1'-methyl-4,4'-bispiperidine

Prepared from indole-6-carboxylic acid (58%).

5 1H-NMR

IS-MS, m/e (M+1)

Analysis for $C_{30}H_{39}N_5O_4S \cdot 2H_2O$:

Calcd: C, 59.88; H, 7.20; N, 11.64;

Found: C, 59.97; H, 6.65; N, 11.43.

10 Analytical RPHPLC, Method 1, RT = 29.02 min (98%)

Example 331.

1-(3-Aminoindazole-5-carbonyl-D-phenylglycinyll)-1'-methyl-4,4'-bispiperidine

15 To a stirring solution of 1-(3-cyano-4-fluorobenzoyl-D-phenylglycinyll)-1'-methyl-4,4'-bispiperidine (120 mg, 0.259 mmol) in p-dioxane (6 mL) was added hydrazine hydrate (26 mg, 0.518 mmol), and the solution was heated to reflux.

After 2 h, the heat was removed and the solvent was
20 evaporated in vacuo. The residue was dissolved in ethanol and heated to reflux. After 12 h, the solution was cooled and concentrated in vacuo. The residue was chromatographed over silica gel, eluting with 10% (2 N ammonia in methanol) in dichloromethane. The product containing fractions were
25 combined and concentrated in vacuo to give 75 mg (62%) of an off white solid.

1H-NMR

IS-MS, m/e 475.3 (M+1)

Analytical RPHPLC, Method 1, RT = 14.72 min (100%)

Example 332.

1-(1-Methyl-3-aminoindazole-5-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

Using methods substantially equivalent to those described in

- 5 Example 331, the title compound was prepared from methylhydrazine and 1-(3-cyano-4-fluorobenzoyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine (31%).

¹H-NMR

IS-MS, m/e 489.2 (M+1)

- 10 Analytical RPHPLC [Vydac C18, linear gradient of 98/2 - 80/20 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min] RT = 38.99 min (100%).

Example 333.

- 15 1-(Imidazo[1,2-a]pyrimidine-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

Imidazo[1,2-a]pyrimidine-2-carboxylic acid

- To a stirring solution of ethyl 1-(imidazo[1,2-a]pyrimidine-20 2-carboxylate (1 g, 5.2 mmol) [Abignente, et al. Eur. J. Med. Chem. (1994) 29, 279] in ethanol (30 mL) was added 2 N aqueous KOH (10 mL, 20 mmol). The solution was heated to reflux; and after 2 h, the heating mantle was removed, the solution was allowed to cool and the solvent was removed by
25 rotary evaporation. The residue was dissolved in water (20 mL) and acidified to pH 3 with 5 N HCl. The resulting precipitate was filtered, washed with water and dried in vacuo to give 700 mg (83%) of a tan solid.

¹H-NMR

- 30 FD-MS, m/e 163.2 (M+1)

Analysis for C₇H₅N₃O₂:

Calcd: C, 51.54; H, 3.09; N, 25.76;

Found: C, 51.12; H, 3.25; N, 25.25.

1-(Imidazo[1,2-a]pyrimidine-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- 5 Using Coupling Method B, the title compound was prepared from imidazo[1,2-a]pyrimidine-2-carboxylic acid and 1-D-phenylglyciny-1'-methyl-4,4'-bispiperidine (56%).

¹H-NMR

IS-MS, m/e 461.2 (M+1)

- 10 Analytical RPHPLC [Vydac C18, linear gradient of 98/2 - 80/20 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min] RT = 32.72 min (96%).

Example 334.

- 15 1-(5,6,7,8-Tetrahydro-imidazo[1,2-a]pyrimidine-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine

- To a stirring solution of 1-(imidazo[1,2-a]pyrimidine-2-carbonyl-D-phenylglyciny)-1'-methyl-4,4'-bispiperidine (250 mg, 0.542 mmol) in ethanol (5 mL) was added sodium
20 borohydride (103 mg, 2.71 mmol). After 24 h, the mixture was diluted with water and extracted 3 times with dichloromethane. The organic phase was dried (MgSO₄), filtered and concentrated in vacuo. The residue was dissolved in dichloromethane and chromatographed over silica
25 gel, eluting with 5% through 10% (2 N NH₃ in MeOH) in dichloromethane. The product containing fractions were combined and concentrated in vacuo to give 55 mg (20%) of the title compound.

¹H-NMR

- 30 IS-MS, m/e 465.2 (M+1)

Analytical RPHPLC [Vydac C18, linear gradient of 98/2 - 80/20 (0.1% TFA in water / 0.1% TFA in acetonitrile) over 40 min, 1 mL/min] RT = 28.44 min (97%).

5 Examples 335 to 338

Preparation of Starting Materials

Ethyl hydroxyimino-pyridine-4-acetate

10 The oxime was prepared in 82% yield from ethyl pyridine-4-acetate using a procedure similar to that described above under Examples 321-324 for the preparation of ethyl hydroxyimino-pyridine-2-acetate.

¹H-NMR (DMSO)

IS-MS, m/e 194.9 (M+1)

15

Boc-D,L-(4-Pyridinyl)glycine ethyl ester

The protected amino ester is prepared from ethyl hydroxyimino-pyridine-4-acetate using a procedure similar to that described above under Examples 321-324 for the preparation of Boc-D,L-(2-pyridinyl)glycine ethyl ester.

20

1-[Boc-D,L-(4-Pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine

25 The protected amide is prepared from Boc-D,L-(4-pyridinyl)-glycine ethyl ester and 1-methyl-4,4'-bispiperidine dihydrochloride using a procedure similar to that described above under Examples 321-324 for the preparation of 1-[Boc-D,L-(2-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine.

30

1-[D,L-(4-Pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine

The amine is prepared from 1-[Boc-D,L-(4-pyridinyl)-glycinyll]-1'-methyl-4,4'-bispiperidine using a procedure

similar to that described above under Examples 321-324 for the preparation of 1-[D,L-(2-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine.

- 5 **General Procedure:** The product of Examples 335-338 is prepared from 1-[D,L-(4-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine and the indicated acid using Coupling Method D.

10 **Example 335.**

1-[4-Methoxybenzoyl-D,L-(4-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine
From 4-methoxybenzoic acid.

15 **Example 336.**

1-(Indole-6-carbonyl-D,L-(4-pyridinyl)glycinyll)-1'-methyl-4,4'-bispiperidine
From indole-6-carboxylic acid.

20 **Example 337.**

1-[3-Methylindole-6-carbonyl-D,L-(4-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine
From 3-methylindole-6-carboxylic acid.

25 **Example 338.**

1-[3-Chloroindole-6-carbonyl-D,L-(4-pyridinyl)glycinyll]-1'-methyl-4,4'-bispiperidine
From 3-chloroindole-6-carboxylic acid.

30 **Assay protocols**

Enzyme Inhibition assays:

The ability of a test compound to inhibit factor Xa may be evaluated in one or more of the following Enzyme Inhibition assays, or in other standard assays known to those skilled
5 in the art.

Enzyme Inhibition Assay 1

Enzyme assays were carried out at room temperature in 0.1M
10 phosphate buffer, pH7.4 according to the method of Tapparelli et al (J. Biol. Chem. 1993,268,4734-4741).

Purified human factor Xa, trypsin, thrombin and plasmin were purchased from Alexis Corporation, Nottingham, UK. Urokinase was purchased from Calbiochem, Nottingham, UK. Chromogenic

15 substrates for these enzymes; pefachrome-FXA, pefachrome-TRY, pefachrome-TH, pefachrome-PL and pefachrome-UK were purchased from Pentapharm AG, Basel, Switzerland. Product

(p-nitroaniline) was quantified by adsorption at 405nm in 96 well microplates using a Dynatech MR5000 reader (Dynex Ltd,

20 Billingshurst, UK). Km and Ki were calculated using SAS PROC NLIN (SAS Institute, Cary, NC, USA, Release 6.11) Km values were determined as 100.9µM for factor Xa/pefachrome-FXA and

81.6µM for trypsin/pefachrome-TRY. Inhibitor stock solutions were prepared at 40mM in Me2SO and tested at 500µM, 50µM and

25 5µM. Accuracy of Ki measurements was confirmed by

comparison with Ki values of known inhibitors of factor Xa and trypsin.

In agreement with published data, benzamidine inhibited

30 factor Xa, trypsin, thrombin, plasmin and urokinase with Ki values of 155µM, 21µM, 330nM, 200nM and 100nM respectively.

NAPAP inhibited thrombin with a Ki value of 3nM. Compounds

of the invention were found to have activity in these assays.

Enzyme Inhibition Assay 2

5

Human factor Xa and human thrombin were purchased from Enzyme Research Laboratories (South Bend, Indiana, USA). Other proteases were from other commercial sources. Chromogenic para-nitroanilide peptide protease substrates
10 were purchased from Midwest Biotech (Fishers, Indiana, USA).

The binding affinities for human factor Xa were measured as apparent association constants (K_{ass}) derived from protease
15 inhibition kinetics as described previously.^{a,b,c,d} The apparent K_{ass} values were obtained using automated (BioMek-1000) dilutions of inhibitors (K_{ass} determinations are performed in triplicate at each of four-eight inhibitor concentrations) into 96-well plates and chromogenic
20 substrate hydrolysis rates determined at 405 nm using a Thermomax plate reader from Molecular Devices (San Francisco). For factor Xa inhibition, the assay protocol was: 50 μ l buffer (0.06 M tris, 0.3 M NaCl, pH 7.4); 25 μ l inhibitor test solution (in MeOH); 25 μ l human factor Xa (32
25 nM in 0.03 M tris, 0.15 M NaCl, 1 mg/ml HSA); finally, 150 μ l BzIleGluGlyArgpNA (0.3 mM in water) added within 2 min to start hydrolysis. Final factor Xa was 3.2 nM. Free [Xa] and bound [Xa] were determined from linear standard curves on the same plate by use of SoftmaxPro software for each
30 inhibitor concentration and apparent K_{ass} calculated for each inhibitor concentration which produced hydrolysis inhibition between 20% and 80% of the control (3.2 nM factor

Xa): apparent $K_{ass} = [E:I]/[E_f][I_f] = [E_b]/[E_f][I^0 - I_b]$.

The apparent K_{ass} values so obtained are approximately the inverse of the K_i for the respective inhibitors [$i/appK_{ass} = app K_i$]. The variability of mean apparent K_{ass} values

5 determined at the single substrate concentration was +/- 15%. The assay system K_m was measured as 0.347 +/- 0.031 mM [$n=4$]; and V_{max} was 13.11 +/- 0.76 $\mu M/min$.

K_{ass} values were determined with thrombin and other
10 proteases using the same protocol with the following enzyme and substrate concentrations: thrombin 5.9 nM with 0.2 mM BzPheValArgpNA; XIa 1.2 nM with 0.4 mM pyroGluProArgpNA; XIIa 10 nM with 0.2 mM HDProPheArgpNA; plasmin 3.4 nM with 0.5 mM HDValLeuLyspNA; nt-PA 1.2 nM with 0.8 mM
15 HDIleProArgpNA; and urokinase 0.4 nM with 0.4 mM pyroGluGlyArgpNA; aPC 3 nM with 0.174 mM pyroGluProArgpNA; plasma kallikrein 1.9 nM with D-PropPheArgpNA; bovine trypsin 1.4 nM with 0.18 mM BzPheValArgpNA.

20 Citations

(a) Sall DJ, JA Bastian, SL Briggs, JA Buben, NY Chirgadze, DK Clawson, ML Denny, DD Giera, DS Gifford-Moore, RW Harper, KL Hauser, VJ Klimkowski, TJ Kohn, H-
25 S Lin, JR McCowan, AD Palkowitz, GF Smith, ME Richett, K Takeuchi, KJ Thrasher, JM Tinsley, BG Utterback, S-CB Yan, M Zhang. Dibasic Benzo[b]thiophenes Derivatives as a Novel Class of Active Site Directed Thrombin Inhibitors. 1. Determination of the Serine Protease
30 Selectivity, Structure-Activity Relationships and Binding Orientation. J Med Chem 40 3489-3493 (1997).

(b) Smith GF, TJ Craft, DS Gifford-Moore, WJ Coffman, KD Kurz, E Roberts, RT Shuman, GE Sandusky, ND Jones, N Chirgadze, and CV Jackson. A Family of Arginal Thrombin Inhibitors Related to Efegatran. Sem. Thrombos. Hemost. 22,
5 173-183 (1996).

(c) Smith GF, DS Gifford-Moore, TJ Craft, N Chirgadze, KJ Ruterbories, TD Lindstrom, JH Satterwhite. Efegatran: A New Cardiovascular Anticoagulant. In New Anticoagulants for the
10 Cardiovascular Patient. Ed. R Pifarre. Hanley & Belfus, Inc., Philadelphia (1997) pp 265-300.

(d) Sall DJ, JA Bastian, NY Chirgadze, ML Denny, MJ Fisher, DS Gifford-Moore, RW Harper, VJ Klimkowski, TJ
15 Kohn, HS Lin, JR McCowan, ME Richett, GF Smith, K Takeuchi, JE Toth, M Zhang. Diamino Benzo[b]thiophene Derivatives as a Novel Class of Active Site Directed Thrombin Inhibitors: 5. Potency, Efficacy and Pharmacokinetic Properties of Modified C-3 Side Chain
20 Derivatives. In press, J Med Chem (1999).

In general, the compounds of formula (I) exemplified herein have been found to exhibit a K_i of 10 μM or less in Assay 1 and/or a K_{ass} of at least 0.1×10^6 L/mole in Assay 2.

25

The ability of a test compound to elongate Partial Thromboplastin Time (Prothrombin Time) may be evaluated in the following test protocols.

30 Partial Thromboplastin Time (Prothrombin) Test Protocol

Venous blood was collected into 3.2% (0.109M) trisodium citrate vacutainer tubes at 1 volume of anticoagulant to nine volumes of blood. The blood cells were separated by centrifugation at 700g for ten minutes to yield plasma, which was frozen at 70°C until required.

To perform the test, 100µl of plasma was pipetted into a glass test tube, 1µl of test compound in DMSO was added, and allowed to warm to 37°C over two minutes. 100µl of warm (37°C) Manchester (tissue thromboplasin) reagent (Helena

Biosciences, UK) was added, allowed to equilibrate for two minutes. 100µl of warm (37°C) 25mM calcium chloride solution was added to initiate clotting. The test tube was tilted three times through a 90° angle every five seconds to mix the reagents and the time to clot formation recorded. Data from a series of observations and test compound concentrations are analysed by a SAS statistical analysis program and a CT2 (Concentration required to double clotting time) for each compound is generated.

Compounds of the invention were found to significantly elongate the partial thromboplastin time (Prothrombin time).

Example No.	Conc. necessary to double the prothrombin time (μM) ^a
8	26
27	6.7
30	7.8
32	11
35	8.8
38	9.0
39	12
40	12
62	8.6
63	2.1
64	4.4
65	6.1

66	2.1 (average of 3 tests)
68	3.6
69	5.8
70	4.0

a The concentration quoted is that of the solution which, when added to the other reagents in the assay, doubles prothrombin time. The final concentration in the assay mixture is one third of this value.

By way of comparison with the result for the compound of Example 66, the compound of Example 75 of WO99/11657 was found to double prothrombin time at a concentration of 11.4 μ M (average of 3 tests).

10 By way of comparison with the result for the compound of Example 35, 1-aminoisoquinolin-7-oyl-D-phenylglycine-4-(4-fluoro-2-methanesulfonylphenyl)-piperazinamide ditrifluoroacetate salt (a compound within the scope of WO99/11657) was found to double prothrombin time at a concentration of 45 μ M (average of 3 tests).

Alternative Prothrombin Time and APTT Protocols

20 Coagulation Determinations. Prothrombin Times and APTT values were determined in HUMAN PLASMA with a STA instrument (Stago). BioPT is a special non-plasma clotting assay triggered with human tissue factor (Innovin). Possible binding to albumen or to lipid was assessed by comparing the BioPT effects in the presence/absence of 30 mg/ml human

albumen (HSA) and 1 mg/ml phosphatidyl choline (PC).
Inhibitors were delivered in 50% MeOH vehicle.

APTT ASSAY

- 5 75 μ l plasma Citrol *Baxter-Dade* Citrated Normal
Human Plasma
25 μ l test sol'n
75 μ l Actin *Baxter-Dade* Activated Cephaloplastin incubate 2
min min. @ 37°
- 10 75 μ l CaCl_2 (0.02 M)


PT ASSAY

- 75 μ l plasma
25 μ l test sol'n
- 15 75 μ l saline incubate 1 min. @ 37° C
75 μ l *Innovin Baxter-Dade* Recombinant Human Tissue Factor

Compounds of the invention were found to be potent
inhibitors of factor Xa.

CLAIMS

1. A serine protease inhibitor compound of formula (I)



(I)

where R₂ represents a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring atom, optionally being substituted in the 3 and/or 4 position by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO₂- or R₁, or the substituents at the 3 and 4 positions taken together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring optionally substituted by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}, and optionally substituted in the position alpha to the X-X group (i.e. 6 position for a six membered aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio with the proviso that R₂ cannot be aminoisoquinolyl;

each X independently is a C, N, O or S atom or a CO, CR_{1a}, C(R_{1a})₂ or NR_{1a} group, at least one X being C, CO, CR_{1a} or C(R_{1a})₂;

each R_{1a} independently represents hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl, alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

R₁ is as defined for R_{1a}, provided that R₁ is not unsubstituted aminoalkyl;

L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or
5 cyclic group;

Y is a nitrogen atom or a CR_{1b} group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group optionally substituted by groups R_{3a} or phenyl optionally substituted by R_{3a};

10 each R_{3a} independently is R_{1c}, amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkyl imidazolyl, thiazolyl, alkyl thiazolyl, alkyl oxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl,
15 aminosulphonyl, haloalkoxy and haloalkyl;

Lp is a lipophilic organic group;

D is a hydrogen bond donor group; and n is 0, 1 or 2;

and

R_{1b}, R_{1c} and R_{1j} are as defined for R_{1a},
20 or a physiologically tolerable salt thereof.

2. A compound as claimed in Claim 1, where

R₂ represents a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, oxygen or sulphur ring
25 atom, optionally being substituted in the 3 and/or 4 position by halo, nitro, haloalkoxy, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO₂- or R₁, or the substituents at the 3 and 4 positions taken together form a
30 fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring optionally substituted by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido,

alkylthio, alkenyl, alkynyl or R_{1j} , and optionally substituted in the position alpha to the X-X.. group (i.e. 6 position for a six membered aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, 5 alkoxy or alkylthio with the proviso that R_2 cannot be aminoisoquinolyl; and

each R_{1a} independently represents hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl alkoxyalkyl, alkoxy carbonyl, acyloxymethoxycarbonyl or 10 alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl.

3. A compound as claimed in Claim 1 or Claim 2, in which n is 0.

15

4. A compound as claimed in any one of Claims 1 to 3, in which X-X is selected from -CH=CH-, -CONH-, -CONR $_{1a}$ -, -NH-CO-, -NH-CH $_2$ -, -CH $_2$ -NH-, -CH $_2$ O-, -OCH $_2$ -, -COO-, -OC=O- and -CH $_2$ CH $_2$ - is CONH.

20

5. A compound as claimed in Claim 4, in which X-X is CONH.

25

6. A compound as claimed in any one of Claims 1 to 5, in which Y is a CR $_{1b}$ group and has the conformation that would result from construction from a D- α -aminoacid NH $_2$ -CR $_{1b}$ (Cy)-COOH where the NH $_2$ represents part of X-X.

7. A compound as claimed in any one of Claims 1 to 6, in which Y is CH.

30

8. A compound as claimed in any one of Claims 1 to 7, in which Cy represents an optionally R $_{3a}$ substituted phenyl,

pyridyl, thienyl, thiazolyl, naphthyl, piperidiny1 or cycloalkyl group.

9. A compound as claimed in Claim 8, in which R_{3a}
 5 is selected from hydrogen, hydroxyl, methoxy, ethoxy, methyl, ethyl, methylaminomethyl, dimethylaminomethyl, hydroxymethyl, carboxy, methoxymethyl, methoxycarbonyl, ethoxycarbonyl, methylaminocarbonyl, dimethylamino-carbonyl, aminomethyl, $CONH_2$, CH_2CONH_2 , acetyl amino,
 10 methoxycarbonylamino, ethoxycarbonylamino, t-butoxycarbonylamino, amino, fluoro, chloro, cyano, nitro, thiol, methylthio, methylsulphonyl, ethylsulphonyl, methylsulphenyl, methylsulphonylamido, ethylsulphonylamido, methylaminosulphonyl, ethylaminosulphonyl, aminosulphonyl,
 15 trifluoromethoxy and trifluoromethyl.

10. A compound as claimed in any one of Claims 1 to 9, in which Cy is phenyl, 4-aminophenyl, 4-amidophenyl, 4-(N-methyl)amidophenyl, 4-(N,N-dimethyl)amidophenyl, 2-
 20 chlorophenyl, 2-methylphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-methoxyphenyl, 4-methoxyphenyl, 4-carboxyphenyl, 3-ethylsulphonylaminophenyl, thien-2-yl, thien-3-yl, thiazol-4-yl, thiazol-5-yl, 2-methylthiazol-4-yl, pyrid-2-yl, pyrid-
 25 3-yl, pyrid-4-yl, piperidin-4-yl, 1-methylpiperidin-4-yl, cyclohexyl or naphth-1-yl.

11. A compound as claimed in any one of Claims 1 to 10, in which L represents CO , CH_2NH , $CONR_{1d}(CH_2)_m$,
 30 $(CH_2)_mN(R_{1d})CO(CH_2)_m$, $(CH_2)_{m+2}$, $CO(CH_2)_m$, $(CH_2)_mCO$, $(CH_2)_mOC=O$, $(CH_2)_mO$, $CH=CH(CH_2)_m$, SO_2 , SO_2NR_{1d} , $SO_2(CH_2)_m$,


$(\text{CH}_2)_m\text{SO}_2$ or $(\text{CH}_2)_m\text{SO}_2\text{NR}_{1d}$ (where each m is independently 0 or 1 and R_{1d} is as defined for R_{1a}).

12. A compound as claimed in Claim 11, in which L is CO,
5 CONH, CH_2NHCO and CONHCH_2 .

13. A compound as claimed in any one of Claims 1 to 12,
in which L_p is an alkyl, alkenyl, carbocyclic or
heterocyclic group, or a combination of two or more such
10 groups linked by a spiro linkage or a single or double bond
or by C=O, O, S, SO, SO_2 , CONR_{1e} , $\text{NR}_{1e}\text{-CO-}$ or NR_{1e} linkage
(where R_{1e} is as defined for R_{1a}), optionally substituted by
one or more oxo or R_3 groups in which R_3 is as defined for
 R_{3a} .

15

14. A compound as claimed in Claim 13, in which L_p is a
group of formula:



in which:

20 r is 1 or 2;

one of X_a and X_b is N and the other is CH or N provided that
when r is 1, X_a and X_b are not both N;


s , t and u are each 0 or 1;

L_a and L_b are each independently selected from a single
25 bond, C=O, O and NR_{1e} , in which R_{1e} is hydrogen or (1-
6C)alkyl;

G is (1-6C)alkanediyl; and

R_{10} is (1-6C)alkyl, (3-6C)cycloalkyl which is unsubstituted
or substituted by (1-6C)alkyl, indanyl, pyridyl,


tetrahydropyranyl, tetrahydrothiopyranyl, phenyl which is unsubstituted or substituted by one or two R_3 groups, pyrrolinyl, or a group of formula



- 5 in which v is 1, 2 or 3; one of X_c and X_d is N and the other is CH or N, provided that when v is 1, X_c and X_d are not both N; and R_{11} is hydrogen, (1-6C)alkyl or when X_d is CH, hydroxy(1-6C)alkyl; provided that when t is 0, the sum of s and u is 1; when X_b is N, L_a is a bond or $C=O$; when X_c is N, L_b is a bond or $C=O$; when X_b and X_c are both N, t is 1; and when $(L_a)_s-(G)_t-(L_b)$ represents an alkyl group and X_b and X_c both represent N, the alkyl group contains at least two chain carbon atoms.

- 15 15. A compound as claimed in Claim 14, in which either X_a is N and L is CO or CH_2CO , or X_a is CH and L is $CONH$, $CONHCH_2$ or CH_2NHCO .


16. A compound as claimed in Claim 13, in which in which
20 $-L-Lp(D)_n$ is:
(i)



in which q is 1 or 2;

- (a) Q is a direct bond; and R_q is piperidin-4-yl which
25 may bear a C_{1-3} alkyl substituent at the 1-position; or R_q is


tetrahydropyranyl, tetrahydrothiopyranyl, phenyl which is unsubstituted or substituted by one or two R_3 groups, pyrrolinyl, or a group of formula



- 5 in which v is 1, 2 or 3; one of X_c and X_d is N and the other is CH or N, provided that when v is 1, X_c and X_d are not both N; and R_{11} is hydrogen, (1-6C)alkyl or when X_d is CH, hydroxy(1-6C)alkyl; provided that when t is 0, the sum of s and u is 1; when X_b is N, L_a is a bond or C=O; when X_c is N, L_b is a bond or C=O; when X_b and X_c are both N, t is 1; and when $(L_a)_s-(G)_t-(L_b)$ represents an alkyl group and X_b and X_c both represent N, the alkyl group contains at least two chain carbon atoms.

- 15 15. A compound as claimed in Claim 14, in which either X_a is N and L is CO or CH_2CO , or X_a is CH and L is CONH, CONHCH₂ or CH₂NHCO.

- 20 16. A compound as claimed in Claim 13, in which in which $-L-L_p(D)_n$ is:
(i)



in which q is 1 or 2;


- (a) Q is a direct bond; and R_q is piperidin-4-yl which may bear a C₁₋₃alkyl substituent at the 1-position; or R_q is

NR_aR_b in which each of R_a and R_b independently is hydrogen or C_{1-3} alkyl; or one of R_a and R_b is hydrogen or methyl and the other of R_a and R_b is $-CH_2-R_c$ or $-CH_2-R_d$ in which R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, $CONH_2$, SO_2NH_2 , methylaminosulphonyl, dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and in which R_d is isopropyl or cyclopentyl, or NR_aR_b is pyrrolidino, piperidino, morpholino, piperazino, or tetrahydro-1,4-diazepino in which a pyrrolidino or piperidino may be a 3,4-didehydro derivative and in which a pyrrolidino, piperidino, piperazino, or tetrahydro-1,4-diazepino may bear a methyl group at the 4-position;

(b) Q is $-O-$ or $-NH-$; and R_q is R_c which is defined as above; or

(c) Q is methylene; and R_q is NR_aR_b which is defined as above;


(ii)



in which R_r is $-(CH_2)_c-R_c$, $-CHR_eR_f$, $-CH_2-CHR_eR_f$, or R_g in which c is 1 or 2 and R_c is defined as above; each of R_e and R_f independently is hydrogen or C_{1-3} alkyl; or CHR_eR_f is cyclopentyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position), cyclohexyl (which may bear a methyl, ethyl or hydroxymethyl substituent at the 3- or 4-position), tetrahydropyran-4-yl, tetrahydrothiopyran-4-yl, pyrrolidin-3-yl (which may bear a 1-methyl substituent), piperidin-4-yl (which may bear a 1-methyl substituent), or indan-2-yl; and R_g is 2-methylsulphonylphenyl which may bear

a 4-fluoro substituent or R_g is $\lambda^{6-1,1}$ -dioxobenzo[b]thiophen-7-yl;

(iii)




5 in which q is 1 or 2;

s is 0 or 1; and

R_s is $-(CH_2)_C-R_C$, $-CHReR_f$, or $-CH_2-CHReR_f$ each of which is defined as above;


(iv)



10

in which R_t is piperidin-4-yl, piperidin-3-yl or pyrrolindin-3-yl, any of which may bear a C_{1-3} alkyl substituent at the 1-position (preferably methyl, ethyl or, more preferably, 2-propyl); or R_t is phenyl (which phenyl may bear a fluoro, chloro, C_{1-4} alkyl, methoxy or methylsulphonyl substituent); or

(v)



in which Het is a divalent 5 membered heteroaromatic group containing 1, 2 or 3 heteroatoms selected from O, N and S and having the two ring atoms at which it is connected separated by one ring atom;

h is 0 or 1; and

20

R_h is phenyl which may bear one or more R_3 substituents.

17. A compound as claimed in Claim 16, in which

5 (i) q is preferably 2, and

in (a) R_q is piperidin-4-yl which may bear a (1-3C)alkyl substituent at the 1-position;

and in (b) R_c is pyrid-2-yl, pyrid-3-yl or pyrid-4-yl;

(ii) c is 2 and R_c is pyrid-2-yl, pyrid-3-yl or pyrid-4-yl;

10 (iii) s is 1;


(iv) R_t is piperidin-4-yl which may bear a methyl, ethyl or 2-propyl substituent at the 1-position; and

(v) R_h is phenyl which may bear one or more R_3 substituents independently selected from, for an ortho or a para

15 substituent: C_{1-5} alkyl, fluoro, chloro, difluoromethyl, trifluoromethyl, methoxy, dimethylamino, methylsulphonyl, and C_{1-2} acyl, and for a meta substituent: fluoro, chloro and methyl.

18. A compound as claimed in Claim 17, in which

20 $-L-Lp(D)_n$ is



in which R_h is phenyl which may bear an ortho and/or a para substituent independently selected from, for an ortho: methyl, fluoro, chloro, methylsulphonyl and acetyl, and for
25 a para substituent: methyl, fluoro, chloro, methoxy and dimethylamino;


Z_1 is S, Z_2 is CH, h is 0; or


Z_1 is NH, Z_2 is N, h is 1.



19. A compound as claimed in any one of Claims 13 to 18, in which R_3 is selected from hydrogen, hydroxyl, methoxy, ethoxy, methyl, ethyl, propyl, 2-propyl, butyl, 2-butyl, t-butyl, pentyl, 2-pentyl or 3-pentyl,
- 5 isopropylaminomethyl, dimethylaminomethyl, diethylaminomethyl, dimethylaminoethyl, acetyl, hydroxymethyl, hydroxyethyl, carboxy, methoxymethyl, methoxycarbonyl, ethoxycarbonyl, methylaminocarbonyl, dimethylaminocarbonyl, aminomethyl, aminocarbonyl,
- 10 methylamino, dimethylamino, ethylamino, formylamino, acetylamino, amino, fluoro, chloro, cyano, nitro, thiol, methylthio, methylsulphonyl, ethylsulphonyl, isopropylsulphonyl, methylsulphenyl, 1,2,4-triazol-2-yl, 1,2,4-triazol-4-yl, 1,2,3-triazol-4-yl, 1,3-imidazol-1-yl or
- 15 1,3-imidazol-4-yl, tetrazol-1-yl, tetrazol-5-yl; methylsulphonamido, ethylsulphonamido, propylsulphonamido, methylaminosulphonyl, ethylaminosulphonyl, propylaminosulphonyl, aminosulphonyl, trifluoromethoxy, trifluoromethyl and trichloromethyl.

20

20. A compound as claimed in Claim 13, in which L_p is selected from








where R_8 represents H, OMe, SO_2Me , F, cyano, amido, amino, NO_2 , Cl or OH; and R_1 is hydrogen or (1-6C)alkyl.

- 5 21. A compound as claimed in Claim 13, in which L_p represents



(K)

wherein X_2 is halo, hydrogen, amino, nitro or $CONH_2$.

10

22. A compound as claimed in any one of Claims 1 to 21, in which R_2 represents:

- (i) phenyl optionally being substituted in the 3 and/or 4 position by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, $MeSO_2-$ or R_1 , and optionally substituted at the 6 position by amino, hydroxy, halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio;

- (ii) naphth-2-yl optionally substituted at the 6 or 7 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} and optionally substituted at the 3 position by amino, hydroxy,

halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or

5 benzisoxazol-5-yl optionally substituted at the 3 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(iv) benzimidazol-5-yl or benzothiazol-6-yl optionally substituted at the 2 position by amino;

10 (v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-

15 yl;

(vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

(viii) pyrazol-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, 20 amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(ix) pyrid-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(x) pyrid-3-yl optionally substituted at the 6 25 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(xi) benzofur-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 30 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R₁;

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by alkyl and optionally substituted at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

5 (xiii) indol-6-yl substituted at the 5 position by amino, hydroxy, halo (such as fluoro or chloro), alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio and optionally substituted at the 3 position by halo (such as chloro), haloalkoxy, haloalkyl, cyano, nitro,
10 amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro,
15 amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j}.

23. A compound as claimed in Claim 22, in which R₂ represents:

(i) phenyl optionally being substituted in the 3
20 and/or 4 position by fluoro, chloro, bromo, iodo, nitro, difluoromethoxy, trifluoromethoxy, amino, cyano, trifluoromethyl, methylthio, vinyl, carboxy, acetoxy, MeSO₂-, hydroxy, methoxy, ethoxy, methyl, methoxycarbonyl, methylamino, ethylamino or amido, and optionally substituted
25 at the 6 position by amino, hydroxy, fluoro, methoxycarbonyl, cyano or aminomethyl (preferably phenyl substituted in the 4 position by chloro, amino, vinyl, methylamino, methyl or methoxy, optionally at the 3 position with amino or hydroxy, and optionally at the 6 position with
30 amino or hydroxy);

(ii) naphth-2-yl optionally substituted at the 6, position by hydroxy and optionally substituted at the 3 position by amino or hydroxy;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, 5 indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or benzisoxazol-5-yl optionally substituted at the 3 position by chloro, bromo, amino, methyl or methoxy;

(iv) benzimidazol-5-yl or benzothiazol-6-yl optionally substituted at the 2 position by amino;

10 (v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by methylthio, methyl or acetyl;

(vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;

15 (vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

(viii) pyrazol-2-yl substituted at the 5 position by methyl;

(ix) pyrid-2-yl optionally substituted at the 6 20 position by chloro;

(x) pyrid-3-yl optionally substituted at the 4 position by chloro;

(xi) benzofur-2-yl optionally substituted at the 3 position by chloro, methyl or methoxy, at the 5 or 6 25 position by methyl and at the 6 position by methoxy;

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by methyl and optionally substituted at the 5 or 6 position by fluoro, chloro, bromo, methyl or methoxy;

(xiii) indol-6-yl substituted at the 5 position by 30 chloro, fluoro or hydroxy and optionally substituted at the 3 position by chloro or methyl; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the 3 position by fluoro, chloro or methyl, and optionally substituted at the 5 or 6 position by fluoro, chloro, methyl, hydroxy, or methoxy.

5

24. A compound as claimed in any one of Claim 23, in which R₂ represents indol-6-yl optionally substituted at the 3 position by chloro, bromo, methyl or methoxy or indol-6-yl substituted at the 5 position by chloro, fluoro or hydroxy and optionally substituted at the 3 position by chloro or methyl.

10

25. A compound of formula I as claimed in Claim 1 and named in any one of the Examples herein, or a physiologically tolerable salt thereof.

15

26. A process for the preparation of a compound of formula I as claimed in Claim 1, or a physiologically tolerable salt thereof, substantially as described in any one of the Examples herein.


20

27. A pharmaceutical composition, which comprises a compound as claimed in any one of Claims 1 to 24 together with at least one pharmaceutically acceptable carrier or excipient.

25

Abstract

Compounds of formula (I)



(I)

where R_2 , each X, L, Y, Cy, Lp, D and n are as defined in the specification, are serine protease inhibitors useful as antithrombotic agents.